

FEMSLE 06313

High correlation between Chagas' disease serology and PCR-based detection of *Trypanosoma cruzi* kinetoplast DNA in Bolivian children living in an endemic area

Patrick Wincker ^{*,a}, Marie-France Bosseno ^b, Constança Britto ^a, Nina Yaksic ^c,
Meric Angélica Cardoso ^a, Carlos Médicis Moral ^a and Simone Frédérique Brunière ^b

tion is yet observed) or after specific treatment (when an immune response can persist for years even if the treatment has been successful [2]). These considerations have justified the quest for a more efficient method of parasite detection in chagasic patients. The principal technique that has been tested for *T. cruzi* detection in blood samples is the polymerase chain reaction (PCR). Two main systems, leading respectively to the amplification of kinetoplast minicircle DNA [3–5] or of nuclear satellite sequences [6,7] have been described. Reconstitution experiments have suggested that these techniques are potentially able to detect a single parasite cell in 20 ml of blood [4,8]. However promising, these methods have yet to be validated in clinical samples originating from different geographical regions, due to the great variations in Chagas' disease incidence and clinical manifestations in the Americas. One large-scale study has been conducted in an endemic region of Brazil, showing that PCR can attain a high level of sensitivity in this situation [9]. The present work is aimed at testing the efficacy of PCR diagnosis of Chagas' disease in a radically different situation in Bolivia.

Materials and Methods

Patients and clinical samples

The individuals examined in this study belonged to two different groups. The first was composed of 45 children from the Mizque locality, Campero province, Cochabamba department, Bolivia. They were 5 to 8 years old, and were previously selected by a serological and parasitological (buffy coat) diagnosis for Chagas' disease when they were at school. The second was composed of 8 control individuals from Rio de Janeiro and La Paz who had never lived in endemic areas.

Ten ml of blood were collected from each individual, 5 ml of which were mixed immediately with an equal volume of 6 M guanidine HCl/0.2M EDTA for further processing for the PCR test [4]. The remaining 5 ml were used for serum preparation and buffy coat testing.

Serology and buffy coat tests

Serology was performed by indirect hemagglutination using the HEMAVE kit (Polychaco, Argentina), indirect immunofluorescence (IIF) and enzyme-linked immunosorbent assay (ELISA). A patient was considered serology-positive when his serum reacted in at least two of the three tests. Buffy coat testing was performed in quadruplicate and observed by 2 independent people.

DNA preparation

The guanidine-EDTA-blood lysates were heated for 15 min in boiling water in order to shear the minicircle molecules that constitute most of the kDNA into moderately-sized pieces, and equalise their overall concentration [8]. One hundred μ l was used for DNA preparation. After phenol-chloroform and chloroform extraction, the material was precipitated with ethanol. The pellet was resuspended in 50 μ l of distilled water and stored at -20°C . All these and subsequent steps were performed in a laminar flow hood in a separate room not used for manipulation of amplified products, with dedicated micropipettes and filter-protected tips.

PCR conditions

The amplification reactions were performed in a volume of 75 μ l using the 'hot-start' protocol with a solid paraffin barrier separating the Taq DNA polymerase from the oligonucleotides [10]. The lower phase consisted of 4 μ l of the $10\times$ Taq polymerase buffer (100 mM Tris-HCl, pH 8.3; 500 mM KCl), 1.8 μ l of a 10 mM dNTPs solution, 13.5 μ l of a 25 mM MgCl_2 solution, 200 ng of *T. cruzi*-minicircle specific primers (5'-AAATAATGTACGGG(T/G)GAGATGCATGA 3' and 5' GGTTTCGATTGGGGTTGGTG-TAATATA 3' [11]), and water up to 40 μ l, in a thin-walled reaction tube. An Ampliwax PCR Gem Bead (Perkin-Elmer) was added, melted by placing the tube at 80°C for 5 min, and solidified at room temperature. The upper phase consisted of 7.5 μ l of the DNA sample, 3.5 μ l of $10\times$ Taq DNA polymerase reaction buffer, 2.5 U of Taq DNA polymerase and water up to 35 μ l. The PCR reaction was performed using 2 cycles at 98°C for 1 min and 64°C for 2 min, 33 cycles at

94°C for 1 min and 64°C for 1 min, and one extension step at 72°C for 10 min. The amplified products were electrophoresed in a 2% agarose gel and visualised by ethidium bromide-staining.

Results and Discussion

In the PCR assay used in this study, each blood sample to be tested was subjected to two independent DNA purifications, in order to avoid any problems arising from inhibition in any particular DNA preparation. Each PCR test included 5 samples in duplicate, one DNA preparation from an individual from a non-endemic area (also in duplicate), an amplification reaction without DNA (as a negative control) and an amplification reaction with DNA from a previously characterised chagasic patient (as a positive control). Typical results are shown in Fig. 1. In this way, we analysed 28 blood samples from serology-positive children living in the endemic area. Twenty-six of them (93%) were detected in our PCR assay, while only four of them had parasites detectable in buffy coat testing (Table 1). A previ-

Table 1

Comparison of the PCR results of all individuals with serology and buffy coat test results

Patient status	Patient no.	PCR	
		Positive	Negative
Bolivian children			
Serology-positive with positive buffy coat test	4	4	0
Serology-positive with negative buffy coat test	24	22	2
Serology-negative			
Serology-negative with positive buffy coat test	1	1	0
Serology-negative with negative buffy coat test	16	1	15
Non-chagasic individuals	8	0	8

ous study has indicated the high sensitivity of PCR when compared to xenodiagnosis in detection of *T. cruzi* in blood samples collected in Brazil [9]. Until the advent of PCR, xenodiagnosis was considered to be the most sensitive parasitological technique available for Chag-




Fig. 1. Ethidium bromide-stained agarose gel from a PCR diagnosis experiment performed on 5 Bolivian children and one individual from a non-endemic area. All DNA preparations from blood samples and PCR amplifications were performed in duplicate for each patient. Lane 1, Φ X174 DNA digested with *Hae*III; Lane 2, amplification reaction without added DNA; lanes 3 and 4, amplifications from a control individual; Lanes 5 and 6 to 11 and 12, amplification products from four Bolivian children with positive serology; lanes 13 and 14, amplifications from a serology-negative Bolivian child. Fifteen μ l of the total reaction volume (75 μ l) was loaded on this gel. The arrows indicate the expected products of the minicircle DNA amplification: a 330 bp band corresponding to a single variable region and a 660 bp band corresponding to a dimer.

disease diagnosis [12]. The results of the present study extend these findings to a different geographical area and suggest that PCR may find extensive applications in situations where other methods for parasitological diagnosis of Chagas' disease are currently being used.

Seventeen blood samples from serology-nega-

from the UNDP/World Bank/WHO Special Programme for Research and Training in Tropical Diseases, the INSERM (Réseau Nord-Sud), PAPES-Fiocruz, CNPq and FINEP.

References

- 11 Degraeve, W., Fragoso, S.P., Britto, C., Van Heuverswyn, Kidane, G., Cardoso, M.A., Mueller, R.U., Simpson, L. and Morel, C.M. (1988) Peculiar sequence organization of kinetoplast DNA minicircles from *Trypanosoma cruzi*. *Mol. Biochem. Parasitol.* 27, 63-70.
- 12 Segura, E.L. (1987) *Xenodiagnosis*. In: Chagas' Disease Vectors (Brener, R.R. and Stoka, A.M., Eds.), Vol. 2, pp. 41-45. Boca Raton Fla: CRC Press Inc.
- 13 Valencia Telleria, A. (1990) *Investigación Epidemiológica Nacional de la Enfermedad de Chagas*. Ministerio Prevision Social y Salud Publica. Secretaria Ejecutiva. 480 Titulo III, La Paz, Bolivia.
- 14 Camargo, M.E., Amato Neto, V. (1974) Anti-*T. cruzi* bodies as serological evidence of recent infection. *Inst. Med. Trop. São Paulo* 16, 200-202.
- 15 Brener, Z. (1980) Immunity to *Trypanosoma cruzi*. *Parasitol.* 18, 247-301.

Reprinted from

FEMS

MICROBIOLOGY LETTERS

FEMS Microbiology Letters 124 (1994) 419-424

© 1994 Federation of European Microbiological Societies 0378-1097/94/\$07.00

Published by Elsevier

FEMSLE 06313

High correlation between Chagas' disease serology and PCR-based detection of *Trypanosoma cruzi* kinetoplast DNA in Bolivian children living in an endemic area

Patrick Wincker ^{*a}, Marie-France Bosseno ^b, Constança Britto ^a, Nina Yaksic ^c,
Maria Angélica Cardoso ^a, Carlos Médicis Morel ^a and Simone Frédérique Brenière ^b

^a Laboratório de Biologia Molecular e Doenças Endêmicas, Departamento de Bioquímica e Biologia Molecular, Instituto Oswaldo Cruz, Fundação Oswaldo Cruz, Av. Brasil 4365, 21045-900 Rio de Janeiro, Brazil; ^b UMR CNRS/ORSTOM, Génétique Moléculaire des Parasites et des Vecteurs, CP 9214, La Paz, Bolivia; and ^c Instituto Boliviano de Biología de Altura, Universidad Mayor de San Andrés, La Paz, Bolivia

(Received 17 October 1994; accepted 19 October 1994)

