

New Prenylated Quinones from *Peperomia galioides*

Valérie Mahiou, François Roblot, Reynald Hocquemiller, and André Cavé*

Laboratoire de Pharmacognosie, URA 1843 CNRS (BIOCIS), Faculté de Pharmacie, Université de Paris XI, 92296 Châtenay-Malabry Cedex, France


Antonieta Rojas De Arias, Alba Inchausti, Gloria Yaluff, and Alain Fournet

IICS, Rio de la Plata y la Gerenza, Casilla de Correo 2511, Asunción, Paraguay


Received December 27, 1995[®]

Two new prenylated quinones, piperogalone (**1**) and galopiperone (**2**), and a new prenylated dihydroquinone, hydropiperone (**3**), were isolated from *Peperomia galioides* H.B.K (Piperaceae). Hydropiperone exhibited potent antiparasitic activity against three species of *Leishmania*.


Previous phytochemical studies of a petroleum ether extract of *Peperomia galioides* H.B.K. (Piperaceae) afforded three interesting prenylphenols: grifolin, grifolic acid, and piperogalin.¹ In a continuation of our investigations on this extract, we have isolated three new minor constituents, piperogalone, galopiperone, and hydropiperone, to which structures **1**, **2**, and **3** have been assigned.



1



2



3

We have previously shown¹ that the petroleum ether extract has a significant in vitro activity on three species of *Leishmania*, responsible for leishmaniasis, and on three strains of *Trypanosoma cruzi*, the causative factor for Chagas' disease. The activities of the major isolated compounds have been reported already.¹ One of the new compounds, hydropiperone (**3**), displays a promising activity against promastigote forms of three species of *Leishmania*.

The petroleum ether extract (7% of dried material) was purified by chromatographic methods on Si gel (column chromatography and TLC) and resulted in the isolation of two new prenylated quinones **1** and **2** (piperogalone and galopiperone) and the 2,3-dihydroquinone **3** (hydropiperone).

The HREIMS spectrum of **1** shows a molecular ion peak at m/z 342.2183, corresponding to the molecular formula $C_{22}H_{30}O_3$. The presence of conjugated carbonyl groups is deduced from the IR spectrum (broad absorption at 1641 cm^{-1}) and is substantiated by two $^{13}\text{C-NMR}$ signals at δ 187.3 and 184.9 indicative of a quinone moiety.² A bathochromic shift, obtained in the UV spectrum after addition of base, indicates the presence of a phenolic or enolic function, also deduced from the IR absorption at 3407 cm^{-1} . The $^1\text{H-NMR}$ spectrum shows the singlet of an aromatic methyl group at δ 2.01 and two methylene groups resonating at δ 3.12 and 3.22 as doublets. The quinone nucleus is thus substituted by two alkenyl moieties further identified by a COSY experiment as a geranyl chain and a prenyl group.

The homonuclear $^1\text{H-}^1\text{H}$ and heteronuclear $^1\text{H-}^{13}\text{C}$ correlations clearly indicated the presence of the prenyl group, corresponding with the methylene at δ 3.22, the ethylenic proton at δ 4.97, and two methyl groups at δ 1.65 and 1.74 (Table 1). Other signals in the $^1\text{H-NMR}$ spectrum were attributed to the geranyl moiety, in particular the ethylenic protons at δ 5.15 and 5.05 and the isolated methylene doublet at δ 3.12 bonded with the quinone nucleus. The $^{13}\text{C-NMR}$ chemical shift for the C-10' methyl group indicated the *E*-geometry about the asymmetrically substituted double bond.³

The respective positions of the four substituents of the quinone were then deduced from the long-range $^1\text{H-}^{13}\text{C}$ experiment (HMBC) and NOE (Table 1), which further confirmed the attributions of ^1H and ^{13}C signals of the geranyl and prenyl chains.

The two alkenyl methylene groups at δ 3.12 and 3.22 were both 3J correlated with the C-1 carbonyl group at δ 187.3, thus indicating the position of the geranyl and prenyl chains at C-2 and C-6, respectively. The proton signal at δ 3.12 was also 3J correlated with the OH-bearing carbon at δ 152.4, indicating a C-3 position for the hydroxy group, and subsequently the position of the methyl group at C-5. Further confirmation of this position was brought by the observation of 3J correlations of the methyl protons with C-6 (δ 137.2) and with the second carbonyl group (δ 184.9) located at C-4. The structure of the new prenylated quinone, named piperogalone, is thus established as structure **1**. The spatial proximity between the methyl group and the prenyl methylene was further supported by a NOE experiment.

[®] Abstract published in *Advance ACS Abstracts*, June 15, 1996.




Table 1. NMR Data for Quinones 1 and 2 (in CD₃COCD₃)

position	1			2		
	δ ¹ H ^a	δ ¹³ C ^b	HMBC (H to C) ^a	δ ¹ H ^a	δ ¹³ C ^b	HMBC (H to C) ^a
1		187.3			184.5	
2		120.4			115.6	
3		152.4			151.3	
4		184.9			182.7	
5		145.2			142.9	
6		137.2			139.7	
7 ^d	2.01, s	11.4	C-4, C-5, C-6	1.98 s	11.9	C-4, C-5, C-6
1'	3.12 d (7)	22.6	C-1, C-2, C-3, C-2', C-3'	6.45 d (10)	116.4	C-3, C-3'
2'	5.15 tq (7, 1)	121.6	C-1', C-4', C-10'	5.71 d (10)	129.7	C-1', C-3'
3'		136.5			83.4	
4'	1.96 t (7)	40.4		1.95–2.05 m	42.2	
5'	≈2.00 m	27.3		2.08–2.13 m	23.4	
6'	5.05 m	125.0	C-8', C-9'	5.09 br t (7)	124.9	C-5'
7'		131.4			132.4	
8'	1.61 br s	25.8		1.60 br s	25.9 ^e	C-6', C-7', C-9'
9'	1.55 br s	17.7		1.53 br s	17.8	C-6', C-7', C-8'
10'	1.74 br s	16.2		1.42 s	27.5	C-2', C-3', C-4'
1'' ^d	3.22 d (6)	26.3	C-1, C-5, C-6, C-2'', C-3''	3.20 d (7)	26.1	C-1, C-5, C-6, C-2'', C-3''
2''	≈4.97 m	120.6	C-4'', C-5''	4.96 br t (7)	120.8	
3''		133.7			133.9	
4''	1.65 d (1.3)	25.8		1.64 br s	25.8 ^e	C-2'', C-3'', C-4''
5''	1.74 br s	18.0		1.72 br s	18.1	C-2'', C-3'', C-5''

^a Spectra were recorded at 400 MHz (J values are given in Hz). ^b Spectra were recorded at 50 MHz. ^c The assignments may be interchanged. ^d NOE enhancements were observed between CH₃-7 and CH₂-1'' for 1.

The second compound isolated from *P. galioides*, named galopiperone (2), is another new quinone as shown by the IR absorption at 1668 and 1647 cm⁻¹ and the two carbonyl signals at δ 184.5 and 182.7 in the ¹³C-NMR spectrum. The ¹H- and ¹³C-NMR data of 1 and 2 are very similar (Table 1) and prove the existence of a prenyl group at C-6 and a methyl group at C-5, but the substituents at C-2 and C-3 are modified.

The molecular formula (C₂₂H₂₈O₃) of 2, deduced from the HREIMS, suggests an oxidative cyclization between the geranyl chain and the C-3 OH group. This assertion is confirmed by the disappearance of the conjugated OH group in the UV and IR spectra, and the presence of an oxygen-bearing quaternary carbon at δ 83.4 in the ¹³C-NMR spectrum. The dihydropyran ring is further characterized by the typical coupling constant ($J = 10$ Hz) between C-1' and C-2', and ³ J correlations in HMBC between H-1' and both C-3 and C-3'. The cyclization between the C-3 OH group and the C-3' position of the geranyl chain is also proved by the significant downfield shift (+9.3 ppm) of the C-10' methyl group in the ¹³C-NMR spectrum.

The absence of specific rotation for 2 indicates that the oxidative cyclization of 1 into 2 is not stereoselective and suggests that galopiperone (2) could be an artifact. This hypothesis is confirmed by the spontaneous conversion of piperogalone into galopiperone after a few weeks, even at +4 °C.

Hydropiperone (3) was obtained as a yellow oil, whose elemental composition was determined as C₂₂H₃₂O₃ by HREIMS, two mass units more than 1. The IR spectrum shows characteristic absorption bands for H-bonded OH (ν_{\max} 3389 cm⁻¹) and quinonoid CO (1669 cm⁻¹), confirmed by two ¹³C-NMR resonances at δ 195.6 and 198.7.⁴ In comparison with piperogalone (1), the ¹H- and ¹³C-NMR spectra of 3 (Table 2) exhibited similar signals indicative of a *E*-geranyl chain. The resonances of a prenyl moiety are also observed, but the methylene group appears as part of an AMX system at δ 2.37 and 2.24, indicating the absence of free rotation between the quinonoid nucleus and the prenyl chain. On the EIMS,

the base peak at m/z 275 is presumably due to the loss of the prenyl side chain and confirms the presence of this group.

The reduction of one double bond in the quinone nucleus, suggested by the IR spectrum, the molecular formula, and the downfield shift of the two carbonyl groups in the ¹³C-NMR spectrum, is confirmed by the observation of two high-field resonances at δ 47.0 and 48.0, corresponding to a quaternary carbon and an inequivalent methylene group (δ ¹H 2.65 and 2.75), respectively.

The structure for hydropiperone, therefore, can be proposed as 3. The respective positions of the substituents were determined on the basis of HMBC correlations, which indicated the position of the geranyl chain at C-6, of the OH group at C-5, and of both the methyl group and the prenyl chain at C-3.

The geminal position of the two latter groups was substantiated by NOE enhancements between the methyl group, the prenyl methylene, and the methylene at C-2 (Table 2). The NOE experiments also provided some information about the molecular conformation. Irradiation of the C-7 methyl group resulted in the enhancement of the methylene groups at C-2 (δ 2.75 and 2.65) and C-1'' (δ 2.37 and 2.24), suggesting the equatorial position of the methyl group. Significant enhancements were also observed between the C-5'' methyl group (δ 1.56) and the methylene at C-1'' (δ 2.24 and 2.37), indicating a *cis* position of this methyl.

Hydropiperone was isolated as a racemic mixture, as shown by the absence of specific rotation.

Piperogalone (1), galopiperone (2), and hydropiperone (3) belong to the rare class of prenylated quinones or dihydroquinones.⁵⁻⁷ They are probably derived biogenetically from the prenylated diphenols reported earlier in the same plant,¹ by oxidative procedures accompanied by the introduction of a prenyl chain at C-5 or C-6 of the diphenol precursor.

Despite the very small amount of compounds available, we investigated the antiparasitic activity for two of them, according to Mahiou *et al.*¹, Hocquemiller *et*

Table 2. NMR Data for 2,3-Dihydroquinone 3 (in CD₃COCD₃)

position	δ ¹ H ^a	δ ¹³ C ^b	HMBC (H to C) ^c	NOESY ^d
1		195.6		
2	2.65 d (16) 2.75 d (16)	48.0	C-1, C-3, C-4, C-7, C-1''	H-7
3		47.0		
4		198.7		
5	OH 2.84 s	153.9		
6		125.3		
7	1.22 s	23.2	C-2, C-4, C-1''	H-2, H-1''
1'	3.09 d (7)	21.9	C-1, C-5, C-6, C-2', C-3'	H-2', H-10'
2'	5.14 br t (7)	119.9		H-1', H-4'
3'		135.8		
4'	1.95 t (7)	39.3	C-2', C-3', C-5', C-6', C-10'	H-2', H-6'
5'	≈2.00 m	26.2	C-4', C-6', C-7'	H-6', H-10'
6'	5.07 m	123.9		H-4', H-5'
7'		129.6		
8'	1.62 d (≈1)	24.6 ^c		
9'	1.58 br s	16.5 ^d		
10'	1.72 br s	15.1		H-1', H-5'
1''	2.37 dd (16, 8) 2.24 dd (16, 8)	37.9	C-2, C-4, C-7, C-2'', C-3''	H-7, H-2''
2''	5.03 m	118.5		H-1'', H-4''
3''		135.2		
4''	1.65 br s	24.8 ^c		H-2''
5''	1.56 br s	16.7 ^d		H-1''

^a Spectra were recorded at 400 MHz (*J* values are given in Hz). ^b Spectrum was recorded at 75 MHz. ^{c,d} The assignments may be interchanged.

al.,⁸ and Fournet.⁹ Galopiperone (2) and hydropiperone (3) were inactive against the trypomastigote form of *Trypanosoma cruzi*, responsible for Chagas' disease (gentian violet displayed a total lysis of the parasites at 250 μg/mL).⁹ Concerning the activity against leishmaniasis, hydropiperone (3) displayed a promising toxicity against *Leishmania braziliensis*, *Leishmania donovani*, and *Leishmania amazonensis* at 25 μg/mL, with a total lysis of the parasites at 100 μg/mL. (Pentamidine displayed a total lysis of the parasites at 5 μg/mL and glucantime 40–50% lysis at 100 μg/mL.) These results, however, are somewhat lower than those obtained with the prenylated diphenols precedently reported.¹

Experimental Section

General Experimental Procedures. UV spectra were recorded on a Philips PU 8700 spectrophotometer and IR spectra on a Perkin-Elmer 841 spectrometer. All ¹H- and ¹³C-NMR spectra were recorded in CD₃COCD₃ (δ ppm) on a Bruker AC 200 P spectrometer operating at 200 and 50 MHz, respectively, and a Bruker ARX 400 spectrometer operating at 400 and 100 MHz, respectively. EIMS and CIMS spectra were obtained on a Kratos MS-80 spectrometer.

Plant Material. *P. galioides*, described by A. Bonpland *et al.* in 1815,¹⁰ was collected by A. Fournet in April 1986 along the old road of Chulumani, near the village of Unduavi, Yungas, department of La Paz, Bolivia (altitude, 3000 m). The botanical identification of the species was made by Dr. H. A. Valdebenito, Department of Botany, Ohio State University, Columbus, OH. A voucher specimen (no. AF 615) is deposited in the National Herbarium of Bolivia, La Paz, and in the Ohio State University Herbarium, Columbus, OH.

Extraction and Isolation. The air-dried *P. galioides* (whole plant, 206 g) was extracted in a Soxhlet apparatus with petroleum ether, CH₂Cl₂, and MeOH successively, which gave 14 g, 9 g, and 14 g of crude

extracts, respectively. Si gel GF₂₅₄ was used for TLC. The petroleum extract (7% of dried material) was first purified by a liquid/liquid partition between 1 M NaOH and CH₂Cl₂. The aqueous layer was acidified with 1 M HCl and further extracted with Et₂O. The CH₂Cl₂ extract was submitted to several purifications using flash chromatography on Si gel (0.032–0.063 mm) affording piperogalone (24 mg), galopiperone (12 mg), and hydropiperone (30 mg).

Piperogalone (1): C₂₂H₃₀O₃; yellow oil; UV (EtOH) λ max (log ϵ) 204 (4.43), 277 (3.98); (OH⁻) 232 (3.96), 280 (3.85) nm; IR (dry film) ν max 3407 (br), 1641, 1620, 1450, 1370, 1309 cm⁻¹; EIMS *m/z* 342.2183 (M⁺, 8) [342.2194 calcd], 259 (30), 257 (26), 217 (18), 203 (29), 91 (22), 77 (18), 69 (100); CIMS *m/z* 343 (MH⁺, 100); for ¹H and ¹³C NMR (CD₃COCD₃) data, see Table 1.

Galopiperone (2): C₂₂H₂₈O₃; red oil; [α]_D²⁰ 0 (c 0.07, EtOH); UV (EtOH) λ max (log ϵ) 204 (4.08), 276 (3.52) nm; IR (dry film) ν max 1668, 1647, 1622, 1450, 1370, 1267 cm⁻¹; EIMS *m/z* 340.2022 (M⁺, 9) [340.2038 calcd], 257 (75), 215 (11), 165 (68), 164 (31), 83 (40), 69 (100); CIMS *m/z* 341 (MH⁺, 100); for ¹H and ¹³C NMR (CD₃COCD₃) data, see Table 1.

Hydropiperone (3): C₂₂H₃₂O₃; yellow oil, [α]_D²⁰ 0 (c 0.15, EtOH); UV (EtOH) λ max (log ϵ) 201 (4.17), 242 (3.61), 350 (3.50) nm; (OH⁻) 205 (3.56), 242 (3.71), 352 (3.62); IR (dry film) ν max 3389 (br), 1669, 1377, 1055 cm⁻¹; EIMS *m/z* 344.2365 (M⁺, 4) [344.2351 calcd], 276 (20), 275 (100), 207 (76), 153 (54), 137 (20), 123 (20), 69 (99); CIMS *m/z* 345 (MH⁺, 100); for ¹H and ¹³C NMR (CD₃COCD₃) data, see Table 2.

Biological Assays. In Vitro Study on the Promastigote Form of *Leishmania*.⁸ Hydropiperone (3) was dissolved in DMSO and evaluated against the promastigote forms of *L. amazonensis* (strain LV-79 = IFLA/BR/67/PH8), *L. donovani* (strain PP-75 = MHOM/IN/83/HS70), and *L. braziliensis* (strain 2903 = MHOM/RR/M2903). Microorganisms are deposited in the Instituto de Investigaciones en Ciencias de la Salud (IICS),

Asunción, Paraguay. Each assay was performed three times. The viability of the parasites was estimated by direct observation with an inverted microscope after a 24-h incubation at 28 °C. The positive controls were glucantime (Rhône-Poulenc, France) and pentamidine (May and Baker, UK).

In Vitro Study on the Trypomastigote Form of *T. cruzi*.⁹ Galopiperone (2) and hydropiperone (3) were dissolved in DMSO and evaluated against the trypomastigote forms of *T. cruzi* (strain Y). The microorganism is deposited in the Instituto de Investigaciones en Ciencias de la Salud (IICS), Asunción, Paraguay. Each assay was performed three times. The parasites were counted after 24 h incubation at 4 °C. The positive control was gentian violet.

Acknowledgment. The authors are grateful to J. Mahuteau for NMR measurements and helpful discussions.

References and Notes

- (1) Mahiou, V.; Roblot, F.; Hocquemiller, R.; Cavé, A.; Rojas de Arias, A.; Inchausti, A.; Yaluff, G.; Angelo Barrios, A.; Fournet, A. *J. Nat. Prod.* **1995**, *58*, 324–328.
- (2) Bonny, M. L.; Capon, R. J. *J. Nat. Prod.* **1994**, *57*, 539–540.
- (3) Ampofo, S. A.; Roussis, V.; Wiemer, D. F. *Phytochemistry* **1987**, *26*, 2367–2370.
- (4) Tanaka, H.; Koyama, Y.; Nagai, T.; Marumo, H.; Omura, S. *J. Antibiotics* **1975**, *28*, 868–875.
- (5) Thomson, R. H. *Naturally Occurring Quinones*; Academic Press: London, 1971; pp 93–197.
- (6) Thomson, R. H. *Naturally Occurring Quinones III, Recent Advances*; Chapman and Hall: London, 1987; pp 31–71.
- (7) Howard, B. M.; Clarkson, K. *Tetrahedron Lett.* **1979**, *46*, 4449–4452.
- (8) Hocquemiller, R.; Cortes, D.; Arango, G. J.; Myint, S. H.; Cavé, A.; Angelo, A.; Muños, V.; Fournet, A. *J. Nat. Prod.* **1991**, *54*, 445–452.
- (9) Fournet, A. *Plantes médicinales boliviennes antiparasitaires (leishmaniose et maladie de Chagas): *Galipea longiflora* Krause (Rutaceae), *Pera benensis* Rusby (Euphorbiaceae) et *Ampelocera edentula* Kuhl (Ulmaceae)*. Ph.D. Thesis, Université Paris-Sud, Châtenay-Malabry, France, 1991, pp 83–84.
- (10) Bonpland, A.; Kunth, C.; de Humboldt, A. *Nov. Gen. Sp.* **1815**, *71i*, t17.

NP960148P