


Seafloor Mapping in the West, Southwest and South Pacific: Foreword

Thirty years ago the West and Southwest Pacific Oceans (Figure 1) were quite virgin regarding seafloor mapping. Conventional soundings started to be more intensively carried out around the 70's. Major breakthroughs in seafloor mapping technology occurred in the 70 and 80's allowing new mapping tools to be developed which simultaneously acquire detailed swath bathymetry and reflectivity of the seafloor. This new generation of mapping tools was designed to rapidly investigate the morphology and nature of large

about 1 million square kilometers of seafloor have been mapped.

The considerable amount of new results obtained by swath mapping during the last ten years encouraged us to organize the "Seafloor mapping in the West and SW Pacific: Results and applications" Workshop at Nouméa (New Caledonia) in November 1994.

The papers presented in this Special Issue resulted from this workshop. They cover the South and Southwest Pacific area from the Antarctica up to the Okin-



levels of complexity of the spreading ridge geometry and evolution.

One paper by Exon *et al.* deals with the structures of the western margin of Tasmania, which is a key area for the reconstruction of the initial break-up of the eastern part of Gondwana.

Two papers, Auzende *et al.* and Pelletier and Auzende, are devoted to the study of the "fossil" boundary between the Australia and Pacific plate along the so-called Vityaz Trench Lineament.

The deformation of the Australia-Pacific plate boundary is described and discussed in three papers. Lafoy *et al.* emphasize the present-day deformation due to the collision between New Caledonia and New Hebrides, Collot *et al.* analyze the effects of the oblique subduction of the oceanic Hikurangi Plateau on the Southern Kermadec forearc and Hikurangi continental margin offshore North Island, New Zealand, and Delteil *et al.* study the transition from the Fiordland subduction and southern extension of the Alpine fault to the Puysegur trench and ridge south of New Zealand. Herzer and Mascle more precisely consider the continent backarc Veining Meinesz Fracture Zone in the northwestern part of New Zealand.

The Gracia *et al.* paper on Bransfield Basin is the only one dealing with the opening of a back-arc basin in the very low latitudes of the Pacific Ocean close to the Antarctica.

One theme of the Nouméa Workshop was the applications of swath-mapping. One of these applications

is the use of detailed bathymetric and reflectivity maps for the precise location of fishing grounds in order to evaluate fish concentration around seamounts and shallow banks. In their paper Lehodey and Grandperin, demonstrate the use of swath mapping for fishing deep species around New Caledonia.

The two last papers by Augustin *et al.* and Bourillet *et al.*, are technical papers related to the processing of swath mapping data and their merging with other geophysical or geological data for the use by the scientific community.

In conclusion we hope that the considerable amount of new data presented in this Special Issue will be useful to both the scientific and fishing communities working in the South, West and Southwest Pacific domain. This Special Issue is also the opportunity to thank the captains and crews of all the ships from different countries and institutions that were involved in the data acquisition. We also wish to acknowledge the invaluable contribution made by the 45 referees to ensure the scientific quality of the papers.

JEAN-MARIE AUZENDE

Ifremer/CB, BP 70, 29280 Plouzané, France (Now at Laboratoire de Géologie-Géophysique, Centre ORSTOM de Nouméa, BP A5, UR14, 98848 Nouméa cedex, Nouvelle-Calédonie)

JEAN-YVES COLLOT

ORSTOM, BP 48, 06320 Villefranche/ Mer, France (Now at IGNS, PO Box 1320, Wellington, New Zealand)