

HIGH ANNUAL AND SEASONAL VARIATIONS IN MALARIA TRANSMISSION BY
ANOPHELINES AND VECTOR SPECIES COMPOSITION IN DIELMO, A
HOLOENDEMIC AREA IN SENEGAL

DIDIER FONTENILLE, LAURENCE LOCHOUARN, NAFISSATOU DIAGNE, CHEIKH SOKHNA,

Laboratoire de Zoologie Medicale, Institut Francais de Recherche Scientifique pour le Developpement en Cooperation (ORSTOM), Dakar, Senegal; Institut Pasteur, Dakar, Senegal; Departement de Biologie Animale, Universite Cheikh Anta Diop, Dakar, Senegal

Abstract. We conducted a three-year entomologic study in Dielmo, a village of 250 inhabitants in a holoendemic area for malaria in Senegal. Anophelines were captured on human bait and by pyrethrum spray collections. The mosquitoes belonging to the *Anopheles gambiae* complex were identified using the polymerase chain reaction. Malaria vectors captured were *An. funestus*, *An. arabiensis*, and *An. gambiae*. *Anopheles funestus* was the most abundant mosquito captured the first year, *An. arabiensis* in the following years. The annual entomologic inoculation rates calculated by enzyme-linked immunosorbent assay were 238, 89, and 150 for the first, second, and third years, respectively. Each year there was a peak of transmission at the end of the rainy season, but transmission occurred




TABLE 1
Number and percentage of malaria vectors caught from April 1992 to March 1995 by different methods in Dielmo, Senegal*

	Feeding†		Resting		Total
	Outdoors	Indoors	Bedrooms	Storehouses	
No. of <i>An. funestus</i> captured	1,783	2,099	1,630	232	5,744
No. of <i>An. gambiae</i> s.l. captured	5,286	5,039	930	371	11,626
Total number of vectors captured	7,069	7,138	2,560	603	17,370
No. of <i>An. gambiae</i> s.l. PCR-tested	898	773	461	205	2,337
% <i>An. funestus</i>	25.2	29.4	63.7	38.5	33.1
% <i>An. gambiae</i>	9.4	11.6	9.2	24.4	10.8
% <i>An. arabiensis</i>	65.4	59.0	26.9	36.3	56.1
% <i>An. melas</i>	0	0.0‡	0.2	0.8	0.1
Total %	100	100	100	100	100

* *An.* = *Anopheles*; PCR = polymerase chain reaction.
 † Caught on human bait.
 ‡ Only one female was captured on human bait indoors (0.01%).

Variation of the HBR. The number and the relative proportion of each of the vector species strongly varied during the three year follow-up study (Figure 2). Among captures on human bait, *An. funestus* was much more abundant than the other vectors during the first year, while *An. arabiensis* dominated the two following years. *Anopheles gambiae* was the least captured vector each year.

Figure 3 shows the variations of the HBR, rainfall, and temperature for the three-year period. *Anopheles arabiensis* was captured during all 36 months, but showed a peak of abundance each year from June to October. It was very abundant in 1993 and 1994, with its HBR reaching 103 bites per person per night in July 1993 and 164 in August 1994. Unlike *An. arabiensis*, more *An. gambiae* females were captured the first year. The maximum HBR for this species was observed in September 1992 with 25 bites per person per night. Despite the presence of apparently favorable larval development sites, this species was rare or absent during each dry season (December to June).


Anopheles funestus was present throughout the survey, but showed great variation. It had a classic peak of abundance each year during the dry season (December to March).¹³ However, the first year of the survey it was also very abundant in July and August, reaching 50 bites per person per night in July 1992. This species became less common in the

second year but the number of specimens captured increased again at the end of the 1994 dry season.

Vector infection rates. The CS protein rate was calculated for each species each month. The ELISA method detected 1.9 times more *Plasmodium*-positive mosquitoes than did dissection. *Plasmodium falciparum* was found in 92% of the positive mosquitoes, *P. malariae* in 7.3%, and *P. ovale* in 1.7%. One *An. arabiensis* was positive for *P. falciparum* and *P. malariae*, and one was positive for *P. ovale* and *P. malariae*. The infection rate for each species varied greatly, not only according to the season, as it classically does, but also according to the year.

Each year *An. funestus* had the highest infection rate for *P. falciparum* (Table 2). Its mean rate calculated over the three years was 2.62%; it decreased by a factor of 2.5 between the first and the third year. These differences are highly significant ($P = 0.001$, by χ^2 test). The mean rate for *An. gambiae* was 1.46%. It was higher the first year and lower the second year. *Anopheles arabiensis* had a mean rate of 0.6% and decreased between the first and the third year. However, for *An. gambiae* and *An. arabiensis* the variations in the infection rate observed, depending on the year, were not significant ($P = 0.18$ and $P = 0.73$, respectively).

Variation of the EIR. The *P. falciparum* total EIR, calculated as described in the Materials and Methods, for the






FIGURE 3. Rainfall, temperature (temp.), monthly human biting rate, and monthly entomologic inoculation rate calculated by enzyme-linked immunosorbent assay for all three *Plasmodium* species for each vector from April 1992 to March 1995 in Diégo Verde.

TABLE 2

Infection rate, calculated by enzyme-linked immunosorbent assay from the head-thoraces of mosquitoes captured on human bait, for each *Plasmodium* species and each vector species by year*

	First year			Second year			Third year					
	No. tested	<i>P.f.</i>	<i>P.m.</i>	<i>P.o.</i>	No. tested	<i>P.f.</i>	<i>P.m.</i>	<i>P.o.</i>	No. tested	<i>P.f.</i>	<i>P.m.</i>	<i>P.o.</i>
<i>An. gambiae</i>	618	2.10	0.16	0	509	0.78	0	0	314	1.27	0.32	0
<i>An. arabiensis</i>	797	0.75	0.13	0.13	2,993	0.63	0.10	0.03	4,410	0.54	0.09	0
<i>An. funestus</i>	1,977	3.49	0.1	0.05	363	2.20	0	0	1,277	1.40	0.08	0

* *P.f.* = infection rate for *P. falciparum*; *P.m.* = infection rate for *P. malariae*; *P.o.* = infection rate for *P. ovale*; *An.* = *Anopheles*.

three-year period was 437 infected bites per person: 223 the first year, 79 the second year, and 135 the third year. Despite the low infection rate for *P. malariae* and *P. ovale*, their mean annual EIRs were 10.8 and 2.5 infected bites per person, respectively (Table 3). Table 4 shows the total EIR for *P. falciparum* for each of the 12 quarters, and the confidence interval calculated following the Poisson distribution.¹⁷ The maximum EIR observed was approximately 95 (95% confidence interval = 69–133) infected bites during the rainy season from July to September 1992, while the EIR was 0 from April to June 1994 during the dry season when no ELISA-positive mosquitoes were found. For each season, large variations in the EIR were observed during the different years.

Figure 3 shows the variation in the EIR by month for each of the three vectors. The EIR was calculated by ELISA for each vector species. During the first year, transmission was higher, reaching 53 infected bites per person in July 1992. The comparative role of each vector according to each month is clearly identified. The first year, *An. funestus* was the main vector, accounting for 76% of all transmission, with 180 infected bites per person. At the end of the rainy season, *An. gambiae* also played a prominent role, whereas *An. arabiensis* was only a secondary vector. In contrast, during the following two years, *An. arabiensis* was responsible for the most transmission, sometimes with sudden variations from one month to another, as in June and July 1994, when the monthly EIR for *An. arabiensis* increased from 0 to 44 (Figure 3).

DISCUSSION

This study shows the high annual and seasonal variations of malaria transmission in Dielmo. Mosquitoes were present throughout the year because of the permanent stream and the presence of different kinds of anopheline larval development sites. One aim of the study was to learn as accurately as possible the rate of transmission of malaria by the differ-

ent vectors to determine the relationship between the variations in transmission and the variations in clinical incidence, parasitemia, and immunologic parameters. Great care must be taken in the analysis of these variations in clinical and biological data due to the great variation in transmission found at four different levels: 1) the relative vector frequency according to the place and method of capture, 2) the abundance and relative frequency of vectors according to the month, as is classically observed, but also according to the year, 3) the infection rate of each vector by year, and 4) the number of infected bites for all vectors, and for each species, for the year.

The identification of all mosquito specimens captured, particularly those caught on humans, is necessary. Since 1990, a PCR that identifies five species of the *An. gambiae* complex from individual dried mosquitoes has been used.^{12,16} This technique makes it possible to determine the species of every specimen collected, particularly those caught on humans, and not just the half-gravid ones, as is the case with cytogenetic methods. Prior to this study, we demonstrated that the PCR technique gave the same identification as cytogenetics on West African mosquitoes of the *An. gambiae* complex.¹⁸ Previous cytotoxic studies have shown two main cytotypes in the *An. gambiae* specimens of this region: savanna and bissau.¹⁹

Comparative studies on transmission using only cytogenetics on half-gravid females captured as indoor resting mosquitoes for the determination of specimens of the *An. gambiae* complex are often biased because this type of sample leads to an overrepresentation of *An. gambiae*, which is well known for its endophilic behavior, as our results clearly show. Among specimens from the *An. gambiae* complex, only 14.4% of the mosquitoes captured on humans were *An. gambiae*, whereas this rate was 25.4% and 39.6% for mosquitoes collected by spraying in bedrooms and storehouses, respectively.

There are few studies on the transmission of malaria that have analyzed differences in transmission according to the

TABLE 3

Annual entomologic inoculation rate calculated by enzyme-linked immunosorbent assay for the three *Plasmodium* species by vector species*

	First year			Second year			Third year		
	<i>P.f.</i>	<i>P.m.</i>	<i>P.o.</i>	<i>P.f.</i>	<i>P.m.</i>	<i>P.o.</i>	<i>P.f.</i>	<i>P.m.</i>	<i>P.o.</i>
<i>An. gambiae</i>	32.5	2.5	0	10.1	0	0	8.4	1.7	0
<i>An. arabiensis</i>	17.5	2.5	2.5	48.2	7.6	2.6	81.0	10.7	0
<i>An. funestus</i>	172.8	5	2.5	20.2	0	0	45.8	2.5	0
Total	222.8	10	5	78.5	7.6	2.6	135.2	14.9	0

* *P.f.* = *P. falciparum*; *P.m.* = *P. malariae*; *P.o.* = *P. ovale*; *An.* = *Anopheles*.

TABLE 4

Entomologic inoculation rate for *Plasmodium falciparum* calculated by enzyme-linked immunosorbent assay with 95% confidence intervals (values in parentheses), by quarter from April 1992 to March 1995 in Dielmo, Senegal

	April to June	July to Sept	Oct to Dec	Jan to March	Total
April 1992 to March 1993	27.5 (14-50)	95.3 (69-133)	47.5 (29-75)	52.5 (33-80)	222.8 (179-274)
April 1993 to March 1994	5.2 (1-18)	23.1 (11-44)	40.1 (23-66)	10.1 (3-26)	78.5 (53-111)
April 1994 to March 1995	0 (0-9)	68.8 (49-93)	25.7 (15-43)	40.7 (23-65)	135.2 (106-173)

month by the different vector species. In Africa, the only published work is that of Taylor and others,²⁰ but the determination of the species of the *An. gambiae* complex was done only on anophelines caught resting indoors by radio-labeled DNA probes. In our study, only 12 *An. melas* were caught. The larval development sites of this halophilic and zoophilic species are located near the mangrove forest 6 km west of Dielmo, and probably very few specimens reached the village. The infection rate was calculated by ELISA because this method allowed the processing of all mosquitoes and the comparison of the results using the same supplies and technique. To limit the overestimation of the sporozoite index by this method, only the head-thoraces were tested. While dissection in the field was done by different people and sensitivity might have varied, the results of the ELISA always remained the same throughout the survey and made possible the comparison from month to month. In our study, the ELISA detected 1.9 times more positive mosquitoes than dissection. This higher value is similar to the results of other studies.^{8, 21}

Using a standardized protocol for collecting and analyzing mosquitoes for three years, we observed very large variations in malaria transmission in Dielmo. As is classically observed in Africa, the EIR for the three vectors varied

the dry season and it ensured continuous transmission throughout the year. For these reasons, this species probably plays an important role in maintaining immunity in Dielmo. In the second and the third years, *An. arabiensis*, which is generally considered to be a less efficient vector, was the major vector due to its very high HBR, despite its low infection rate. *Anopheles arabiensis* accounted for 10%, 66%, and 61%, respectively, of the transmission over the three-year period. Transmission by *An. gambiae* was the lowest except in the first year, during which its EIR was twice that of *An. arabiensis*. Over the three-year period, *An. gambiae* was responsible for only 12% of the transmission.

These data differ from those obtained in other regions of West Africa, where it is generally considered that *An. gambiae* and *An. funestus* are more efficient vectors than *An. arabiensis*.²² In the Gambia, 30 km south of Dielmo, studies carried out during the rainy season between 1979 and 1981 and in 1988 showed that *An. gambiae* was largely dominant.^{6, 19} This was also the case in Burkina Faso in the region of Bobo Dioulasso.²³ To our knowledge, no study has found such an important role for *An. funestus* in West Africa, as was found in Dielmo. Some years, for as yet unidentified reasons, *An. funestus* can be very abundant and transmission can be higher during the dry season than during the rainy

greatly according to the month, with a peak of transmission during and at the end of the rainy season from July to September. More surprising were the great variations in the entomologic components of transmission, such as the HBR and the mosquito infection rate, as well as the number and relative proportion of the three vectors over the three years.

From April to June 1994, no mosquitoes carrying CS protein were found in 179 captured, whereas two years before during the same season, 28 infected bites were recorded. Likewise, transmission was higher from January to March 1993 in the dry season than during the following rainy season from July to September. These data show that in Dielmo, where larval development sites are permanent, there is no justification for dividing the year into only two seasons (dry and rainy) to assess transmission and to correlate it with clinical and immunologic data because there are too many variations between the years.

The role of each vector species was different for each of the three years. The first year, *An. funestus* had a much great-

season. This species is, however, present in the entire Sudanese and Guinean climatic region, and is an important vector.²⁴ In East Africa and Madagascar, it can play a major role.^{8, 20, 25}

During the three-year study, we were unable to find any explanation, for example, temperature or rainfall, that could predict the variation of the observed transmission and the involvement of different vector species.

These detailed data on transmission are necessary to correlate with data on parasitemia, incidence of clinical attacks, associated immune responses, human genetic factors, and variation of *Plasmodium* polymorphism. Recent studies show that several different *Plasmodium* genotypes exist in Dielmo.^{26, 27} These genotypes could have varying pathogenicities and may not be transmitted with equal efficiency by the different vector species. For these reasons, only a very thorough study of transmission would provide useful data for associated research.

of Cooperation (grant 93837), the Institut Francais de Recherche Scientifique pour le Developpement en Cooperation (ORSTOM), and the Institut Pasteur de Dakar.

Authors' addresses: Didier Fontenille, Laurence Lochouarn, Nafisatou Diagne, Cheikh Sokhna, Jean Jacques Lemasson, Mathurin Diatta, Farba Faye, and Jean-Francois Trape, ORSTOM, BP 1386, Dakar, Senegal. Lassana Konate, Departement de Biologie Animale, Universite Cheikh Anta Diop, Dakar, Senegal. Christophe Rogier, Institut Pasteur, BP 220, Dakar, Senegal.

Reprint requests: Didier Fontenille, ORSTOM, BP 1386, Dakar, Senegal.

REFERENCES

1. Hviid L, Theander TG, 1993. Seasonal changes in human immune responses to malaria. *Parasitol Today* 9: 26-27.
2. Beier JC, Oster CN, Onyango FK, Bales JD, Sherwood JA, M, Legros F, Molez JF, Trape JF, 1995. Les anopheles du Senegal. Liste commentee et illustree. *Bull Soc Pathol Exot* 87: 267-277.
12. Paskewitz SM, Collins FH, 1990. Use of polymerase chain reaction to identify mosquito species of the *Anopheles gambiae* complex. *Med Vet Entomol* 4: 367-373.
13. Gillies MT, DeMeillon B, 1968. The Anophelinae of Africa South of the Sahara. Second edition. Johannesburg: South African Institute of Medical Research.
14. Burkot TR, Williams JL, Schneider I, 1984. Identification of *Plasmodium falciparum*-infected mosquitoes by a double antibody enzyme linked immunosorbent assay. *Am J Trop Med Hyg* 33: 783-788.
15. Wirtz RA, Zavala F, Charoenvit Y, Campbell GH, Burkot TR, Schneider I, Esser KM, Beaudoin RL, Andre RC, 1987. Comparative testing of monoclonal antibodies against *Plasmodium falciparum* sporozoites for ELISA development. *Bull World Health Organ* 65: 139-145.
16. Scott JA, Brogdon WG, Collins FH, 1993. Identification of single specimens of the *Anopheles gambiae* complex by poly-

Page 2

ISSN = 0002-9637

VOLUME 56

MARCH 1997

NUMBER 3

The American Journal of
**TROPICAL
MEDICINE &
HYGIENE**

ORSTOM
CENTRE DE DOCUMENTATION
Montpellier

22 MAI 1997

