

Systematic Numbering of Vegetative Compatibility Groups

TABLE 1. Vegetative compatibility groups (VCGs) of *Fusarium oxysporum* listed by forma specialis

forma specialis	Host	forma specialis code	Assigned numbers for VCG	Number of VCGs	Literature citations
<i>albedinis</i>	<i>Phoenix dactylifera</i>	017-	0170	1	32
<i>apii</i>	<i>Apium</i>	001-	0010 to 0012	3	5
<i>asparagi</i>	<i>Asparagus</i>	100-	1001 to 1008	8 + ^a	7
<i>basilici</i>	<i>Ocimum</i>	016-	0160	1	16
<i>betae</i>	<i>Beta</i>	... ^b	...	7 +	12
<i>chrysanthemi</i>	<i>Chrysanthemum</i>	005-	0050 to 0051	2	27
<i>ciceris</i>	<i>Cicer</i>	1	25
<i>conglutinans</i>	<i>Brassica</i>	010-	0101	1	20
<i>cubense</i>	<i>Musa</i>	012-	0120 to 0126, 0128 to 01220	21 ? ^c	26
<i>cucumerinum</i>	<i>Cucumis sativus</i>	018-	0180	1 +	15
<i>cyclaminis</i>	<i>Cyclamen</i>	015-	0151 to 0153	3	36
<i>dianthi</i>	<i>Dianthus</i>	002-	0020 to 0022, 0025, 0027, 0028	6	1
<i>elaeidis</i>	<i>Elaeis</i>	001- ^{*,d} , 014-	0011 to 0015, 0140 to 0141	5	6, 10
<i>erythroxyli</i>	<i>Erythroxyllum</i>	1	31
<i>gladioli</i>	<i>Gladiolus</i>	034-	0340 to 0343	4	24
<i>lactucum</i>	<i>Lactuca</i>	1	13
<i>lilii</i>	<i>Lilium</i>	4	22
<i>lupini</i>	<i>Lupinus</i>	2 ?+	28, 29
<i>lycopersici</i>	<i>Lycopersicon</i>	003-	0030 to 0033	4 +	23
<i>mathioli</i>	<i>Mathiola</i>	010-	0103	1	20
<i>medicaginis</i>	<i>Medicago</i>	004-	0040 to 0041	2	27
<i>melongenae</i>	<i>Solanum melongena</i>	017-	0170	1	15
<i>melonis</i>	<i>Cucumis melo</i>	013-	0130 to 0138	8 +	14, 17
<i>niveum</i>	<i>Citrullus</i>	008-	0080 to 0082	3	21
<i>phaseoli</i>	<i>Phaseolus</i>	016-	0161 to 0165	5	35
<i>pisi</i>	<i>Pisum</i>	007-	0070	6 ?+	2, 34
<i>radicis-lycopersici</i>	<i>Lycopersicon</i>	009-	0090 to 0098	8 +	18, 30
<i>raphani</i>	<i>Raphanus</i>	010-	0102	1	20
<i>spinaciae</i>	<i>Spinacia</i>	3	9
<i>tracheiphilum</i>	<i>Vigna</i>	006-	0060	1	27
<i>tuberosi</i>	<i>Solanum tuberosum</i>	6	33
<i>vasinfectum</i>	<i>Gossypium</i>	011-	0111 to 01110	10	8

^a + = Additional isolates are incompatible with the established VCGs.

^b ... = VCG code and numbers not assigned.

^c ? = Two or more of these VCGs may overlap.

^d * = forma specialis code previously given to forma specialis *apii* by Puhalla (27).

are actually members of *F. redolens* f. sp. *dianthi*. In cases such as this, VCG numbers for isolates no longer considered to belong to *F. oxysporum* should be removed from use without replacement. A similar four-digit code as used for VCGs in *F. oxysporum* has been proposed for VCGs and formae speciales in *F. redolens* (1). The latter species causes wilt and rot diseases in numerous host plants including pea, carnation, and spinach (11). Other VCGs currently placed in *F. oxysporum* may eventually prove to belong to *F. redolens* or other species of *Fusarium*.

As a rule, nonpathogenic isolates of *F. oxysporum* should not be given four-digit VCG codes. Since tests for vegetative compatibility are unlikely to be done among the many nonpathogenic isolates from different laboratories, we discourage the routine assignment of formal VCG numbers for these isolates to prevent an overabundance of VCG categories. In some cases, however, these isolates will be sufficiently important to warrant classification of their VCG. VCG numbers may be assigned when (i) isolates belonging to the same VCG are recovered from multiple locations, (ii) isolates belonging to the same VCG compose a large fraction (>5%) of the population at a single site (sample size > 100 independent isolates), or (iii) isolates belonging to the same VCG are of particular utility, e.g., biological control isolates. Nonpathogenic isolates vegetatively compatible with pathogenic isolates also should be given the VCG number of their pathogenic counterparts. Four-digit VCG codes for nonpathogens will be assigned by the coordinator as done for pathogenic VCGs. If the following recommendations for numbering are followed, duplications and inconsistencies in the system can be kept to a minimum.

To address the problems outlined here, and with the endorsement of the *Fusarium* Committee of the International Society of Plant Pathology, we hereby propose the creation of a "VCG numbering coordinator." This person should be contacted by investi-

gators prior to submission of new VCG numbers for publication. If requested by the investigator, communications with the VCG numbering coordinator will be kept strictly confidential prior to publication. Talma Katan (Department of Plant Pathology, ARO, Volcani Center, P.O. Box 6, Bet Dagan 50250, Israel; fax: 972-3-968-3543) has accepted our nomination to serve as the initial VCG numbering coordinator until the 6th International Fusarium Workshop (to be held at the International Mycological Institute from 17 to 21 August 1998), at which time a new VCG coordinator can be selected or T. Katan can be renominated to continue as coordinator.

In the coming year, representative isolates of all known VCGs of *F. oxysporum* should be deposited for storage at two or more internationally recognized fungal culture collections. Individual investigators will be contacted shortly to assemble this collection. The dual utility of such a centralized collection is that it will be able to provide reference cultures of all of the major phytopathogenic VCGs and insure long-term strain maintenance. Puhalla (27) noted that "(to) standardize the assignment of VCG, a great deal of cooperation between *Fusarium* workers will be necessary." It is our hope that the efforts begun here will promote the cooperation required to make VCG standardization possible.

LITERATURE CITED

1. Baayen, R. P., van Dreven, F., Krijger, M. C., and Waalwijk, C. 1997. Genetic diversity in *Fusarium oxysporum* f.sp. *dianthi* and *Fusarium redolens* f.sp. *dianthi*. Eur. J. Plant Pathol. 103:395-408.
2. Bodker, L., Lewis, B. G., and Coddington, A. 1993. The occurrence of a new genetic variant of *Fusarium oxysporum* f.sp. *pisi*. Plant Pathol. 42: 833-838.
3. Booth, C. 1971. The Genus *Fusarium*. Commonwealth Mycological Institute, Kew, Surrey, United Kingdom.

4. Correll, J. C. 1991. The relationship between formae speciales, races, and vegetative compatibility groups in *Fusarium oxysporum*. *Phytopathology* 81:1061-1064.
5. Correll, J. C., Klittich, C. J. R., and Leslie, J. F. 1987. Nitrate nonutilizing mutants of *Fusarium oxysporum* and their use in vegetative compatibility tests. *Phytopathology* 77:1640-1646.
6. Dossa, C., Pando-Bahuon, A., Renard, J. L., and Boisson, C. 1991. Determination of vegetative compatibility groups in African *Fusarium oxysporum* strains isolated from vascular wilt-infected oil palms. *Oleagineux* 46:145-147.
7. Elmer, W. H., and Stephens, C. T. 1989. Classification of *Fusarium oxysporum* f. sp. *asparagi* into vegetatively compatible groups. *Phytopathology* 79:88-93.
8. Fernandez, D., Assigbetse, K., Dubois, M. P., and Geiger, J. P. 1994. Molecular characterization of races and vegetative compatibility groups in *Fusarium oxysporum* f.sp. *vasinfectum*. *Appl. Environ. Microbiol.* 60:4039-4046.
9. Fiely, M. B., Correll, J. C., and Morelock, T. E. 1995. Vegetative compatibility, pathogenicity, and virulence diversity of *Fusarium oxysporum* recovered from spinach. *Plant Dis.* 79:990-993.
10. Flood, J., Whitehead, D. S., and Cooper, R. M. 1992. Vegetative compatibility and DNA polymorphisms in *Fusarium oxysporum* f.sp. *elaeidis* and their relationship to isolate virulence and origin. *Physiol. Mol. Plant Pathol.* 41:201-215.
11. Gerlach, W., and Nirenberg, H. I. 1982. The genus *Fusarium*—A pictorial atlas. *Mitt. Biol. Bundesanst. Land-Forstwirtschaft. Berlin-Dahlem.* 209:1-406.
12. Harveson, R. M., and Rush, C. M. 1997. Genetic variation among *Fusarium oxysporum* isolates from sugar beet as determined by vegetative compatibility. *Plant Dis.* 81:85-88.
13. Hubbard, J. C., and Gerik, J. S. 1993. A new wilt disease of lettuce incited by *Fusarium oxysporum* f. sp. *lactucum* forma specialis nov. *Plant Dis.* 77:750-754.
14. Jacobson, D. J., and Gordon, T. R. 1990. Further investigations of vegetative compatibility within *Fusarium oxysporum* f.sp. *melonis*. *Can. J. Bot.* 68:1245-1248.
15. Katan, T. 1996. Vegetative-compatibility groups in populations of *Fusarium oxysporum* in Israel. (Abstr.) *Phytoparasitica* 24:139.
16. Katan, T., Gamliel, A., and Katan, J. 1996. Vegetative compatibility of *Fusarium oxysporum* from sweet basil in Israel. *Plant Pathol.* 45:656-661.
17. Katan, T., Katan, J., Gordon, T. R., and Pozniak, D. 1994. Physiologic races and vegetative compatibility groups of *Fusarium oxysporum* f. sp. *melonis* in Israel. *Phytopathology* 84:153-157.
18. Katan, T., Zamir, D., Sarfatti, M., and Katan, J. 1991. Vegetative compatibility groups and subgroups in *Fusarium oxysporum* f. sp. *radicis-lycopersici*. *Phytopathology* 81:255-262.
19. Kistler, H. C. 1997. Genetic diversity in the plant-pathogenic fungus *Fusarium oxysporum*. *Phytopathology* 87:474-479.
20. Kistler, H. C., Momol, E. A., and Benny, U. 1991. Repetitive genomic sequences for determining relatedness among strains of *Fusarium oxysporum*. *Phytopathology* 81:331-336.
21. Larkin, R. P., Hopkins, D. L., and Martin, F. N. 1990. Vegetative compatibility within *Fusarium oxysporum* f.sp. *niveum* and its relationship to virulence, aggressiveness, and race. *Can. J. Microbiol.* 36:352-358.
22. Löffler, H. J. M., and Rumine, P. 1991. Virulence and vegetative compatibility of Dutch and Italian isolates of *Fusarium oxysporum* f.sp. *lilii*. *J. Phytopathol.* 132:12-20.
23. Marlatt, M. L., Correll, J. C., Kaufmann, P., and Cooper, P. E. 1996. Two genetically distinct populations of *Fusarium oxysporum* f. sp. *lycopersici* race 3 in the United States. *Plant Dis.* 80:1336-1342.
24. Mes, J. J., van Doorn, J., Roebroek, E. J. A., van Egmond, E., van Aartrijk, J., and Boonekamp, P. M. 1994. Restriction fragment length polymorphisms, races and vegetative compatibility groups within a worldwide collection of *Fusarium oxysporum* f.sp. *gladioli*. *Plant Pathol.* 43:362-370.
25. Nogales-Moncada, A. M., Pérez-Artés, E., and Jiménez-Díaz, R. M. 1993. VCG diversity within *Fusarium oxysporum* f.sp. *ciceris*. (Abstr.) Page 169 in: *Abstr. Int. Congr. Plant Pathol.*, 6th. National Research Council Canada, Ottawa.
26. Ploetz, R. C. 1994. Panama disease: Return of the first banana menace. *Int. J. Pest Manage.* 40:326-336.
27. Puhalla, J. E. 1985. Classification of strains of *Fusarium oxysporum* on the basis of vegetative compatibility. *Can. J. Bot.* 63:179-183.
28. Rataj-Guranowska, M. 1992. *Fusarium oxysporum* f.sp. *lupini* in Wielkopolska-genetic interrelationships. Pages 133-139 in: *IHAR Radzikow (Eur. Fusarium Semin., 3rd)*, Poznan, Poland.
29. Rataj-Guranowska, M., Paszewski, A., and Walkowiak, I. 1991. *Fusarium oxysporum* forms heterokaryons with *Fusarium redolens*. *J. Phytopathol.* 132:294-302.
30. Rosewich, U. L., Pettway, R. E., Kistler, H. C., and Katan, T. 1997. Morphological and molecular characterization of isolates of *Fusarium oxysporum* f.sp. *radicis-lycopersici* from Florida. (Abstr.) Page 83 in: *Abstr. Fungal Genet. Conf.*, 19th. Fungal Genetics Stock Center, Kansas City, KS.
31. Sands, D. C., Ford, E. J., Miller, R. V., Sally, B. K., McCarthy, M. K., Anderson, T. W., Weaver, M. B., Morgan, C. T., Pilgeram, A. L., and Darlington, L. C. 1997. Characterization of a vascular wilt of *Erythroxylum coca* caused by *Fusarium oxysporum* f. sp. *erythroxyli* forma specialis nova. *Plant Dis.* 81:501-504.
32. Tantaoui, A., Ouinten, M., Geiger, J.-P., and Fernandez, D. 1996. Characterization of a single clonal lineage of *Fusarium oxysporum* f. sp. *albedinis* causing Bayoud disease of date palm in Morocco. *Phytopathology* 86:787-792.
33. Venter, S. L., Theron, D. J., Steyn, P. J., Ferreira, D. I., and Eicker, A. 1992. Relationship between vegetative compatibility and pathogenicity of isolates of *Fusarium oxysporum* f. sp. *tuberosi* from potato. *Phytopathology* 82:858-862.
34. Whitehead, D. S., Coddington, A., and Lewis, B. G. 1992. Classification of races by DNA polymorphism analysis and vegetative compatibility grouping in *Fusarium oxysporum* f. sp. *pisi*. *Physiol. Mol. Plant Pathol.* 41:295-305.
35. Woo, S. L., Zoina, A., Del Sorbo, G., Lorito, M., Nanni, B., Scala, F., and Noviello, C. 1996. Characterization of *Fusarium oxysporum* f. sp. *phaseoli* by pathogenic races, VCGs, RFLPs, and RAPD. *Phytopathology* 86:966-973.
36. Woudt, L. P., Neuvel, A., Sikkema, A., van Grinsven, M. Q. J. M., de Milliano, W. A. J., Campbell, C. L., and Leslie, J. F. 1995. Genetic variation in *Fusarium oxysporum* from cyclamen. *Phytopathology* 85:1348-1355.

