

QUATERNARY STATE OF STRESS IN THE NORTHERN ANDES AND THE RESTRAINING BEND MODEL FOR THE ECUADORIAN ANDES

Frédéric EGO (1), Michel SEBRIER (1), Alain LAVENU (2), Hugo YEPEZ (3), Arturo EGUEZ (3)

(1) URA 1369 CNRS, bât 509, Université de Paris Sud, 91405 Orsay cedex, France.

(2) ORSTOM, 213 rue La Fayette, 75480 Paris cedex10, and, LGMBS, UPPA, 64000 Pau, France.

(3) Escuela Politecnica Nacional, Quito, Ecuador.

Résumé : L'inversion des mécanismes au foyer superficiels des Andes septentrionales, et l'analyse néotectonique des Andes d'Equateur, montrent un champ de contrainte homogène ($\sigma_1 \pm E-W$). Celui-ci induit un mouvement dextre sur les failles N30-35°E et du raccourcissement sur les failles N-S, (relais compressif nord équatorien). Les vitesses de déplacement des failles dextres majeures sont calculées et discutées.

Key Words : Sismotectonics, State of Stress, Quaternary, Andes, Ecuador, Kinematics.

INTRODUCTION

The subduction of the Nazca Plate beneath South America results from a N80°E trending convergence (Fig. 1). This convergence is thus oblique along the N30°E, 1500-km long northern Andes. This obliquity is accommodated in the upper plate by NNE trending, active dextral faults. In Ecuador, two major active strike-slip faults are known. (1) The Pallatanga Fault (PF), that is located in the Cordillera Occidental of southern Ecuador, is well studied (Soulas, 1988; Winter, 1990; Soulas et al., 1991; Winter et al., 1993). (2) The Rio Chingual-la Sofia Fault (CSF), that is located in the Cordillera Real (CR) of northern Ecuador (Soulas et al., 1988; 1991; Tibaldi & Ferrari, 1992), is still poorly known. In addition, several Quaternary, N-S striking folds and reverse faults have been reported in northern Ecuador (Winter, 1990; Ego et al., 1993); They are interpreted as a restraining bend between the PF and the CSF (Ego et al., 1993). In order to address this problem, we have analyzed the stress pattern in northern Andes inverting shallow focal mechanism data. Then, we calculated slip rates on the major strike-slip and reverse faults.

INVERSION OF FOCAL MECHANISMS

In the literature, 113 shallow focal mechanisms, registered by world seismic networks, are available (e.g., Pennington 1981; CMTS by Dziewonski 1981-1992 and USGS catalog 1977-1988). They are mainly distributed along the Andean subduction zone and in the Sub-Andean region; in contrast, few events are located in the Andean mountains. We have defined 8 zones, each one corresponding to a specific structural province. Then, data inversion (same method as Carey-Gailhardis & Mercier, 1987) was performed in each zone in order to calculate its mean state of stress (Fig. 2). The most significant result is that the state of stress is quite homogeneous in northern Andes, σ_1 trending roughly E-W, except nearby the Caribbean.

CSF SLIP RATE DETERMINATION AND IAV SHORTENING ESTIMATE

In order to constrain the CSF slip rate, a detailed fault mapping was performed using LANDSAT imagery coupled with aerial photo analysis and field control. Numerous dextral offsets are seen along the 70 km long, N35°E striking trace of the CSF in Ecuador. The rio Chingual, a major river of the CR, exhibits a dextral offset ranging between 7.5km and 10.5km. Other streams of minor importance show offsets between 50m and 500m. All these offset streams, that are from different ages, allow us to conclude that the fault has been active in a dextral sense for a long time (>1My). These offsets could be used to calculate the slip rate of the fault; however, lacking of precise stream ages, we determined the CSF slip rate using the Soche lava





Figure 1: Geodynamic setting and major structures in northern Andes and restraining bend model for the northern Ecuadorian Andes. CSF: Rio Chingual-la Sofia Fault, CPFS: Cauca-Patia Fault System, IAV: Inter-Andean Valley and PF: Pallatanga Fault.



flow dated at 9670 yrBP (Hall & Beate, 1991). Since this lava flow exhibits an offset ranging between 60m and 70m, and assuming a 5% error on the age, the "long term" slip rate is 7 ± 1 mm/yr. In contrast, the N30°E trending PF has a slip rate of 4 ± 1 mm/yr (Winter (1990; Winter et al 1993). The marked difference in slip rate between PF and CSF suggests a northward increase in dextral slip rate along northern Andes.