

NEOGENE TO PRESENT TECTONIC EVOLUTION AND STRESS FIELD IN ECUADOR

Alain LAVENU (1), Frédéric EGO (2), Christophe NOBLET (3), Thierry WINTER (4)

(1) ORSTOM, 213 rue La Fayette, 75480 Paris cedex 10, France and Laboratoire de Géodynamique et Modélisation des Bassins Sédimentaires, UPPA, Avenue de l'Université, 64000 Pau, France.

(2) Laboratoire de Géologie Dynamique Interne, URA CNRS, Géophysique et Géodynamique Interne, Université de Paris-Sud, Centre d'Orsay, Bat. 509, 91405 Orsay cedex, France.

(3) BHP Minerals, 550 California street, San Francisco, California 94104-1020, USA.

(4) Laboratoire de Tectonique et Mécanique de la Lithosphère, IGP, 4 place Jussieu, 75252 Paris cedex 05, France.

Résumé : L'analyse de la microtectonique, celle des déformations synsédimentaires et celle de la morphologie des bassins néogènes et quaternaires de la zone intracordillère des Andes d'Equateur montrent une persistance de la tectonique compressive NNE-SSW à E-W du Miocène inférieur à l'Actuel. Des modifications locales du champ de contrainte peuvent être dues à des conditions particulières.

Key Words : Synsedimentary deformations, stress, Neogene, Andes, Ecuador.

INTRODUCTION

Strong continental sedimentation had developed during the Neogene (Bristow, 1973; Baldock, 1982), in basins formed along major crustal faults in the Ecuadorian Andes. This concerns particularly the basins of Cuenca, Giron, Nabon, Loja, Malacatos and Zumba, in south Ecuador and the Interandean Depression (IAD) with the Quaternary basins of Quito and Latacunga-Ambató in central and northern Ecuador (Figure). These basins are located between the Eastern Cordillera composed essentially of Paleozoic and Mesozoic rocks and the Western Cordillera represented by a series of Cretaceous (and some Cenozoic) volcanic rocks. In southern Ecuador, the Neogene basins are limited by two families of N-S and NE-SW trending faults. In central Ecuador, the Interandean Depression is characterized by an uppermost Pliocene - Quaternary basin, developed alongside great regional N-S faults.

STRATIGRAPHY

In south Ecuador, the Tertiary rocks can be regionally grouped into three superposed lithostratigraphic units (Noblet et al., 1988).

- a lower, essentially volcanic unit (U1, Saraguro Fm.) which comprises the substratum of the basins, and consists of intermediate and acidic lava and pyroclastic deposits (35.3 to 26.8 Ma; 1000 m thick).

- an intermediate sedimentary unit (U2) corresponding to most of the infill. This unit lies in discordance with unit U1 and correspond to the Neogene infill of the basins (22 to 8 Ma; 1500 to 4000 m thick). The unit can be divided into two megasequences: M1 and M2.

- an upper volcano-sedimentary unit (U3, Llacao and Tarqui Fm.) which disconformably overlies the previous units.

The IAD infilling consist of Pliocene to Recent volcanoclastic continental, fluvial and lacustrine sediments. In recent studies we have shown that these deposits consist of four distinct sedimentological units (Lavenu et al., 1992).

- The oldest unit, U1, is essentially composed of lahars, pyroclastic levels, interstratified andesitic flows (400 m thick; 1.4 to 1.8 Ma).

- Unit U2 forms a progressive unconformity over the previous unit (fluvial and lacustrine deposits, 50 to 80 m thick).

These two units represent the Latacunga Formation.

- Unit U3 lies unconformably over the previous units (Chalupas Fm.; 1.2 Ma; >50 m thick).

- Unit U4 is composed of recent pyroclastic tuffs (Cangagua Formation).

NEOGENE SOUTH ECUADORIAN BASINS

Synsedimentary folding deformations affected the basin sediments (Lavenu and Noblet, 1989).

The M1 megasequence deposits (lower Miocene) display sedimentary wedges which indicate the presence of a normal component in the N20°-N40°E faults during sedimentation. An additional folding deformation corresponds to a NNE-SSW shortening direction, perpendicular to the NW-SE tensional direction. This tensional deformation characterizes the basin's opening.

In megasequence M2 a synsedimentary folding deformation with a N60°E trending shortening direction affects the lower to middle Miocene deposits. The development of these folds is kinematically compatible with a right-lateral component of the movement along the N20°E faults.

The upper levels of the megasequence M2 (upper Miocene) show a series of progressive unconformities related to synsedimentary N-S folding with a mean E-W trending shortening.

A microtectonic study evidences an ongoing compressive stress regime during at least the Miocene period.

Lower sediments (M1) and substratum (U1) of the basins display a first direction of the major stress component σ_1 during lower Miocene, between N23°E and N64°E. Then, in the deposits of middle to upper Miocene (M2) and substratum, the direction of major stress component σ_1 is between N71°E to N107°E. A relative chronology of the striations is coherent with a NE-SW compression followed by an E-W compression.

PLIOCENE-QUATERNARY INTERANDEAN BASINS

The IAD is characterized by a large late Pliocene - Quaternary basin situated between N-S reverse basement faults. The formation of the IAD probably began in the Upper Miocene. An E-W shortening drives the reverse faults since upper Pliocene. The shortening became more pronounced during and after deposition of unit U1 (Latacunga Fm.). Analysis of folds and flexures demonstrates that at least between 1.7 and 1.2 Ma, the southern part of the IAD was subject to major shortening along an E-W direction. This deformation began while U1 sediments were being deposited, and continued and intensified during the deposition of U2. The morphology (deviation of rivers, landslides etc...) reveals the continuation of the shortening during the recent Quaternary. The seismicity and focus mechanisms highlight the present-day persistence of the E-W shortening. All of these arguments are in favor of a "push-down" basin interpretation.

CONCLUSION

From a sequential analysis of infill deposits in the South Ecuadorian basins, a phase of deepening of the lower Miocene basins, followed by infilling in the middle to upper Miocene has evidenced. Synsedimentary deformations affected the entire Miocene. They reveal a normal component on N20°E-N40°E trending faults during M1 and compressive tectonics during M2. Shortening directions deduced from fold axes of synsedimentary folds




Figure: Schematic tectonic map of the mean recent and active faults. Location of the Neogene basins.

IAD: Interandean Depression; NB: Neogene basins; Black triangles, volcanoes: A= Atacazo, Al= Altar, C= Casitagua, Co= Chimborazo, Cx= Cotopaxi, Cz= Corazon, I= Iliniza, Ig= Iguazata, Q= Quilotoa, P= Pichincha, S= Sagoatota, T= Tungurahua. Focal mechanism solutions of three shallow-focus (>30 km) earthquakes (lower hemisphere projection). P axis in white quadrant. At the present time Andes of central Ecuador are affected by an E-W trending shortening.

vary between N60°E at the base of M2 to N80°E-N130°E at the top of M2. Microtectonic analysis has evidenced a state of compressive stress with σ_1 following a NNE-SSW direction in the lower Miocene. The lower Miocene deepening is the consequence of right-lateral strike-slip movements along N-S faults and right-lateral-normal movements along N20°E-N40°E faults. The stress field in the middle and upper Miocene is characterized by a major stress σ_1 approximately E-W. Results of the microtectonic analysis, the sedimentological study and the analysis of synsedimentary deformations, suggest that the basin infilling is due to movements with a strong reverse component along N-S and N20°E faults.

The E-W shortening observed in the IAD is kinematically compatible with the present right-lateral kinematics (with a slight reverse component) of the Pallatanga Fault (Winter and Lavenu, 1989; Winter et al., in press). It is also compatible with the pull-apart basin interpretation given for the Gulf of Guayaquil and related to right-lateral movements since the Miocene along the D-G M fault.

The central part of the Ecuadorian IAD between Quito and Ambato is characterized by structures (folds, flexures, reverse faults) which are due to E-W shortening.

To the south of this area, active Quaternary structures have reverse-right strike slip movements, of NE-SW direction which are also due to the E-W compression (Pallatanga Fault).

To the north of this area, in the south of Colombia, a system of active NE-SW right-lateral faults is also due to E-W compression.

The behavior of the Quito-Ambato zone can be illustrated as a compressive N-S relay in a system of large right strike-slip structures.

So, sedimentological, synsedimentary folding deformation and microtectonic analyses show a persistent and continuous compressional stress regime in the Interandean basins of Ecuador, from Miocene times to Present-day.

REFERENCES

- BALDOCK J.W. 1982. Geology of Ecuador: Explanatory Bulletin of the National Geological Map of the Republic of Ecuador. Esc. 1:1 000 000, Min. Rec. Nat. Energ., Quito, 70 p.
- BRISTOW C.R. 1973. Guide to the geology of the Cuenca Basin, southern Ecuador. Ecuadorian Geological and Geophysical Society, 54 p.
- LAVENU A. and NOBLET C. 1989. Synsedimentary tectonic control of Andean Intermontane strike-slip basins of South Ecuador (South America). International Symposium on Intermontane Basins: Geology and Resources: 306-317, Chiang Mai, Thailand.
- LAVENU A., NOBLET C., BONHOMME M.G., EGUEZ A., DUGAS F., and VIVIER G. 1992 New K/Ar age dates of Neogene and Quaternary volcanic rocks from the Ecuadorian Andes : Implications for the relationships between sedimentation, volcanism, and tectonics. J. of S. Amer. Earth Sci., 5, 3/4 : 309-320.
- NOBLET Ch., LAVENU A. and SCHNEIDER F. 1988. Etude géodynamique d'un bassin intramontagneux tertiaire sur décrochements dans les Andes du sud de l'Equateur: l'exemple du bassin de Cuenca. Géodynamique, 3, (1-2): 117-138.
- WINTER T., AVOUAC J.P., and LAVENU A. 1993. Holocene kinematics of the Pallatanga strike-slip fault (Central Ecuador) from topographic measurements of displaced morphological features. Geophys. J. Intern. (in press).
- WINTER Th. and LAVENU A. 1989b. Morphological and microtectonic evidence for a major active right lateral strike-slip fault across central Ecuador (South America). Annales Tectonicae, III, 2: 123-139.