

CO₂ evolution from soil and physical protection of soil organic carbon in a young pasture on Vertisol

CHEVALLIER T.¹, BLANCHART E.¹, ALBRECHT A.² AND FELLER C.³

¹ IRD-BOST, BP 8006, 97259 Fort-de-France cedex, Martinique (FWI)

e-mail: chevali@versailles.inra.fr

² IRD/ICRAF, P.O. Box 30677, United Nations Avenue, Gigiri, Nairobi (Kenya)

³ CENA-USP/IRD, Caixa Postal 96, 13400-970 Piracicaba, SP, Brazil

Key words: Vertisols, re-grassed pasture, soil aggregates, field CO₂ evolution, organic C bioavailability

Introduction

Long periods of continuous cropping were recognised to affect soil organic matter (SOM) content and to increase soil erosion. For example, in Martinique several years of intensive farming on vertisols led to an important decrease in SOM and soil aggregation and to an important increase in soil erosion. There is now a need to restore the soil organic matter in order to preserve the soil resources. In literature, pastures seem to be one of the best agrosystems for enhancing the SOM stock in cultivated soils. In Martinique, planting a *Digitaria decumbens* pasture on a cultivated vertisol increased the soil organic carbon (SOC) stock from 31 to 37 MgC ha⁻¹ (0-20 cm) within 5 years (Chevallier et al., 2000). This high storage value could be explained partly by the heavy clayey soil and the high plant inputs.

Nevertheless, we did not know so far if this increase of SOC stock was a result only of an increase of annual organic carbon (OC) inputs from the *D.decumbens* pasture or also of a decrease of OC outputs by mineralisation. **The first objective** of this study was to show by field CO₂ measurements if that the SOC storage in vertisols could be explained by a relative decrease of SOM mineralisation.

This decrease in SOM mineralisation could be due to a chemical stabilisation of SOM or a physical protection of SOM in soil aggregates to biodegradation. That protection effect was already mentioned in the literature, but few data are available for tropical soils (Beare et al. 1994). In Martinique, Albrecht et al. (1992) observed that soil C promoted the vertisol aggregation as a binding agent. **Our second objective** was to verify throughout laboratory measurements if SOC protection to mineralisation exists in the aggregates of vertisol under pasture.

Materials and methods

The experiment was located in the south-eastern part of Martinique, French West Indies (14°25'N / 60°53'W). The area is characterised by a humid tropical climate. The soil was a smectitic Leptic Hapludert (USDA classification) or Eutric Vertisol (FAO-UNESCO classification) developed on andesite. It has about 50-60% of clay. The soil was irrigated (rain plus irrigation amounted to about 120 mm month⁻¹).

Comparative measurements of CO₂ in field and in laboratory were conducted on several plots under different land use systems as long term market gardening (18 years), re-grassed pasture after fifteen years of market gardening and long term pasture (18 years), providing a range of soil carbon content.

1-Field CO₂ evolution measurement

In the field, CO₂ evolutions were measured by the closed chamber method with an infrared gas analysis. Different replicates in time and space were assumed. Each chamber location was defined by its SOC content measured with a Carbon Nitrogen Sulfur Analyser, NA 1500, (Carlo Erba). The contribution of root respiration in total soil respiration was estimated by comparison of CO₂ emissions between untreated soil and soil treated with systemic herbicide, applied 15 days before measurement.

To test the bioavailability of SOC to microorganisms, we defined an index (I_{CO₂}) of daily CO₂ emission relative to the SOC content. For simplification we only use the SOC content of the upper 10 cm. This index (gCO₂-C kgC⁻¹ day⁻¹) was calculated as followed:

$I_{CO_2} = (C - CO_{2\text{ annual}} / 365) / C$, where $C - CO_{2\text{ annual}}$ is the C-CO₂ annual evolution and C is the soil C.

2-Laboratory CO₂ measurements

Soil was sampled, fragmented in clods (diameter 1-2 cm), and air-dried. The air-dried clods of soil were either not crushed or crushed and sieved to 200 µm. Then the soil samples were incubated in a closed vessel at 28 °C during 21 days. The CO₂ produced is absorbed in sodium hydroxide and quantified by titration.

Results and discussion

1- CO₂ evolved from soil measured in the field.

The annual CO₂ flux increased with the duration of the pasture from 14 Mg CO₂-C ha⁻¹ yr⁻¹ in market gardening to 18-20 Mg CO₂-C ha⁻¹ yr⁻¹ in five years old pasture and to 25 Mg CO₂-C ha⁻¹ yr⁻¹ in a long-term pasture. Data for soil respiration rates from tropical pasture in literature (4.7 to 21 Mg C ha⁻¹ yr⁻¹) were generally inferior or equal to these values. This could be explained by lower soil moisture in these soils compared to the regularly irrigated soil in our study.

Combining the data of each chamber location, we found a significantly positive correlation ($r^2=0.42$, $p<0.01$) between SOC stock and CO₂ fluxes (Fig.1). Since soil respiration is the sum of microorganisms respiration and root respiration, the high CO₂ evolution in the richer SOC soils could be related to a higher root respiration according to the presence of more roots. We compared the soil respiration between plots without living roots (herbicide application) and plots with living roots, and assumed that in this pasture the contribution of root respiration was about 12 %, to the total soil respiration. This value did not explain all the extra CO₂ evolved from richer SOC soils. A larger SOC content in soils gave a larger quantity of OC substrates for soil microorganisms metabolism and then a larger CO₂ emissions from soils. This was even verified by many laboratory measurements but more sparsely by field measurements.


The index (I_{CO_2}) of daily CO₂ emission relative to the SOC content decreased with the duration of the pasture and with the C content ($r^2= 0.40$, $p< 0.01$, Figure 2). In spite of continuous larger C inputs in the re-grassed soil, the bioavailability of the SOC decreased with the pasture establishment. These results illustrate a protection of SOC against mineralisation and subsequently a storage of OC in soils with the pasture establishment. This process of SOC protection against mineralisation could be a chemical stabilization of organic compounds, or a limited access of the SOM to microorganisms (physical protection). The second part of this study was to verify or not this latter hypothesis.

2- Bioavailability of SOC characterised in standard conditions

In standard conditions, as well as in the field, the CO₂ evolved was higher in richer SOC content soil, from 0.26 to 1.04 gC kg soil⁻¹ for clod samples (Figure 3). These values were higher when the samples were crushed (Figure 3), there was then a "de-protection" of some organic materials when the soil has been crushed before incubation. This amount of organic carbon protected in soil structure (200 µm – clods) was higher when the SOC content was higher (Figure 3). Moreover in the long term pasture soil drying-rewetting treatments, which are known to affect intimately the soil structure, also increase C mineralisation more than in the market gardening soil (Figure 3).

These results were comparable to Beare et al. (1994) and Gijsman and Sanz (1998) results, where experiences of tropical soil incubation were conducting in quite the same conditions as in our study (crush to 250 µm and un-crushed samples and soil incubation of 20 and 41 days) (Figure 4). Nevertheless, the increase of CO₂ emission after crushing seemed to be generally lower in temperate conditions than in tropical conditions and larger in cultivated soils than in virgin soils. But few data were available to really conclude.

The second hypothesis mentioned after field CO₂ measurement was verified. In pasture, no soil disturbance plus the development of root enmeshment is promoting soil aggregation. Indeed, Angers (1992) and Jastrow (1996) in temperate re-grassed soil observed an increase of soil aggregation before the OC accumulation in the soil. The SOC can then promote the vertisol aggregation as a binding agent (Albrecht et al. 1992) and in positive feedback, the soil aggregation can promote the SOC storage. Recent C inputs in regrassed soil from D.decumbens could be then incorporated in soil aggregates (Golchin et al. 1994) and then be partly physically protected from mineralisation (Beare et al. 1994).



...uts in five years were st

Correlation between ICO_2 and SO_2


ods (cm)

Crushed and sieved to 200 μm

- Clod without drying-rewetting treatment

50

...SOC content in standard conditions on clods and crush samples.



● Our study

● Litterature (tropical pasture)

○ Litterature (tropical culture)

— Linéaire (Our study)

— Linéaire (Litterature (tropical culture))

...between **SOC** content and amount of protected **SOC** against mineralisation in our study and other (Gijssels et al. 1994, 20 days of incubation; Gijssman and Sanz 1998, 41 days of incubation).

References:

- Albrecht A., Rangon L. and Barret P. (1992) Effet de la matière organique sur la stabilité structurale et la détachabilité d'un vertisol et d'un ferrisol (Martinique). *Cahiers ORSTOM, série Pédologie* 27, 121-133.
- Beare M. H., Cabrera M. L., Hendrix P. F. and Coleman D. C. (1994) Aggregate-Protected and Unprotected Organic Matter Pools in Conventional-Tillage and No-Tillage Soils. *Soil Science Society of America Journal* 58, 3, 787-795
- Chevallier T., Blanchart E., Girardin C., Mariotti A., Albrecht A., Feller C. The role of biological activity (roots, earthworms) in medium-term C dynamics in vertisol under a *D. decumbens* pasture. Accepted in *Applied Soil Ecology* (June 2000).
- Golchin A., Oades J. M., Skjemstad J. O. and Clarke P. (1994) Soil Structure and Carbon Cycling. *Australian Journal of Soil Research* 32, 5, 1043-1068.
- Gijsman A. J. and Sanz J. I. (1998) Soil organic matter pools in a volcanic-ash soil under fallow or cultivation with applied chicken manure. *European Journal of Soil Science* 49, 427-436.