

SEDIMENTATION MODIFIED BY WIND INDUCED RESUSPENSION IN A SHALLOW TROPICAL LAGOON (COTE D'IVOIRE).

ROBERT ARFI, DANIEL GUIRAL and MARC BOUVY

KEY WORDS: sedimentation; resuspension; trade winds; waves; lagoon.

ABSTRACT

In shallow environments, under certain conditions of fetch, wind velocity, bathymetry and bottom characteristics, resuspension can be generated by wind induced waves. In the tropical Ebrié lagoon, austral trade winds are dominant almost all year long, and their velocity shows a marked diel pattern with maximum speed between noon and midnight. Only austral trade winds with a speed $> 3 \text{ m s}^{-1}$ allow particle resuspension which is effective for depths $< 1.5 \text{ m}$. In these areas, significantly higher values of chlorophyll biomass and mineral seston are noted during the windy sequences. Granulometric and mineralogical analyses showed that only the surficial sediment (0-3 cm) was involved in resuspension. This process induces several effects: 1) an increase of the suspended matter concentration in the water and thus a light attenuation due to a higher turbidity, 2) a redistribution in the whole water column of nutrients from the pore water and 3) a removal of the finer fractions from the superficial sediment. On the contrary, for depths $> 1.5 \text{ m}$, particle sinking is permanent in depressions which are spontaneously transformed into anoxic systems. At the lagoon scale, sedimentation is significantly modified by wind induced resuspension. According to the bathymetry and the distance from a river, three sedimentary facies are recognized. Their grain size distributions are parabolic in areas where resuspension occurs, logarithmic in areas where no resuspension is possible and hyperbolic in the hollows and the main channels. Finally, a large part of the allochthonous inputs (from drainage and rivers) and autochthonous pelagic production is trapped into the Ebrié lagoon and less than 10% of the particles entering the lagoon are exported toward the Atlantic Ocean.

INTRODUCTION

In shallow environments, resuspension can be generated by wind induced waves under certain conditions of fetch, wind velocity, bathymetry, and bottom characteristics (KENNISH, 1986; PRESS and SIEVER, 1986). This process is a major factor controlling the spatial and temporal distribution of particles (CARPER and BACHMANN, 1984; FLODERUS and PIHL, 1990; LUETTICH *et al.*, 1990). Resuspension can influence algal productivity (HELLSTRÖM, 1991) as well as the sedimentation process (BENGTSSON *et al.*, 1990). In the tropical Ebrié lagoon (average depth: 4.8 m), austral trade winds are dominant, and, in favorable conditions, they can induce sediment resuspension (ARFI *et al.*, 1993). A survey was con-

ducted at a shallow water station in order to understand the impact of this process in the pelagic ecosystem. The aims of the present work were to analyze (1) the change of seston and chlorophyll biomass in the water column and (2) the modification of the sedimentary interface using granulometric and mineralogical analyses. The fate of the suspended materials and the sedimentary zonation related to the wind induced resuspension are also discussed at a lagoon scale.

MATERIAL AND METHODS

The study was conducted in a sampling site ($5^{\circ}20 \text{ N}$, $4^{\circ}20 \text{ W}$; mean depth 1 m) located on the

Fonds Documentaire ORSTOM



010015567

427

Fonds Documentaire ORSTOM

Cote : B* 15567 Ex : 1




Fig. 1. Time series of wind velocity and water level during 3 days of an austral trade winds sequence and 3 days of a boreal trade winds (harmattan) sequence.

north shore of the Ebrié lagoon (Côte d'Ivoire) in December 1991 (details in ARFI *et al.*, 1993). Location and geographical characteristics of study area and sampling site are reported in BOUVY *et al.* (1994). During 23 days, the wind velocity was integrated each minute and averaged over 5 min. Water level and wave height were obtained using a surface height gauge. The effective fetch was calculated according to CARPER and BACHMANN (1984) and estimated as 6800 m for the predominant southern winds. The minimum wind velocity needed to generate a wave action reaching the sediment at the sampling station was calculated according to DENMAN and GARGETT (1983) and DEMERS *et al.* (1987). Water samples were collected every 3 h using a peristaltic pump integrating from 10 cm below the surface to 10 cm above the bottom. Seston content was estimated by gravimetric determination and the mineral fraction was determined as loss on ignition. Chlorophyll biomass (corrected for phaeopigments) was measured by




Fig. 2. Average diurnal pattern of hourly wind speed and wave height observed during the trade wind periods 4-15 and 19-26 December 1991.

fluorometry after methanol extraction of pigments.

The suspended materials were sampled using traps deployed for 24 h and the mineral fractions were concentrated by acidification (pH 2.0). The organic fractions were destroyed using hydrogen peroxide with the dried samples.

At the sampling site, sediment was sampled manually with a PVC corer and frozen. Prior to their analysis, the frozen cores were cut into 3 cm thick slices. After lyophilization and decarbonation, the organic fraction was destroyed with hydrogen peroxide. Granulometric analyses were performed on suspended and sedimented particles using a Sedigraph after these preliminary treatments. A qualitative study (nature of particles) was performed using X-ray analysis on total and <math> < 50 \mu\text{m}</math> samples (Siemens 500 diffractometer, Cu anticathode). The mineralogical clay fractions were studied by X-ray analysis according to three processes (oriented slide, glycerol and heating tests).

RESULTS AND DISCUSSION

Wind and resuspension

During the study, three sequences were distinguished according to the wind direction: a period of trade winds (SW direction) from 4 to 15 December, a period of harmattan (NE direction) from 16 to 18 December and another period of trade winds (19-26 December). The trade wind periods were characterized by maximum wind speeds (>3 m s⁻¹) recorded between 14 and 17 h; after sunset, their velocity decreased slowly. At the station studied, there was sufficient fetch for these winds to generate waves, and the wave height varied according to the wind velocity (Fig. 1). A close link between the trade wind speed and the wave height




Fig. 3. Average diel cycle of chlorophyll *a* and seston concentrations observed during the trade wind periods 4-15 and 19-26 December 1991.

was observed, with marked diurnal patterns (Fig. 2). A marked increase of wave height (4 to 6 cm) was noted for wind speeds ranging from 1.5 to 3.5 m s^{-1} , and higher values (7-10 cm) were observed for wind velocity above 3.5 m s^{-1} . Due to the geographical configuration of the site, wind induced resuspension was possible only when the wind speed was above 3 m s^{-1} . During the harmattan period, no waves occurred in spite of a significant wind speed (Fig. 1), due to an insufficient fetch in the wind direction.

Effects of resuspension on seston and bed characteristics

A marked diel cycle closely associated with the wind speed variations was observed both for the mineral seston and chlorophyll biomass during the trade wind periods (Fig. 3). No such pattern was noted during the harmattan period. For both parameters, significant average differences (Student *t*-test after normality assessment) were observed according to wind velocity and direction. When the wave height was sufficient during the trade wind periods, the turbulence induced by winds enhanced the distribution of living and inert particles in the whole water column.

This periodic redistribution of phytoplankton into the euphotic layer induced by the physical forcing (minimum chlorophyll concentrations between 6 and 9 h, maximum between 15 and 18 h) is different from that observed in areas of the Ebrié lagoon not influenced by resuspension (fetch not effective and/or water column >3 m). In the latter case, the minimum chlorophyll concentrations were observed between midnight and 4 h, and the maximum between 10 and 14 h (TORRÉTON *et al.*, 1994). This discrepancy can be explained by variations of the phytoplankton abundance when physical control

occurs, but can also be attributed to grazing processes in relation with the zooplankton photophobia. Other processes, such as variations of the cellular chlorophyll content, in relation with the diurnal light cycle might explain this phenomenon.


Suspended mineral particles were essentially clays (mainly kaolinite and illite) corresponding to the finest fractions from the surficial bed sediments (ARFI *et al.*, 1993). Owing to the wave action, these finest fractions were removed from the surficial bed sediment and distributed in the water column. However, some of the suspended particles had also an allochthonous origin (probably from the near-by Agnéby river) since vermiculite was present in the suspended assemblages but not in the bed sediment.

Only the surficial sediment (0-3 cm) was affected by the removal of the finest fractions. Below this level, the vertical heterogeneity of the sediment (granulometry, mineralogy) was conserved, indicating that erosion related to wind induced resuspension was rapidly attenuated. In the area studied, the alternation of diurnal resuspension and nocturnal sedimentation induced several effects: (1) an increase of suspended matter concentration in the water and thus a light attenuation due to a higher turbidity, (2) a distribution in the whole water column of nutrients from the interstitial water, and (3) a removal of the finest fractions from the sediment interface.

Modification of sedimentation by wind induced resuspension

At the scale of the Ebrié lagoon, the sedimentation is significantly modified by wind induced resuspension. The local bathymetry, the distance from a river (main source of suspended particles) and the eolian control (through the fetch related to the austral trade winds) are at the basis of the zonation of the Ebrié lagoon sediment. The characteristics of a given sedimentary facies are thus dependent on these 3 components of the sedimentary dynamic.

The riverine inputs and the shoreline run-off are essentially carrying clay and sand eroded from the high plateau on the north, sedimentary beach ridge on the south. Once entering the lagoon, these particles settle rapidly, either after discharge into low energy environments (in case of sands) or after flocculation (in case of clays). As a consequence of these processes, the river outfalls feature progradation areas, permanently reworked by the local hydrodynamics. These deposits contribute to the building of shoals or sand-banks, blocking the



Hyperbolic facies



average grain diameter $<2 \mu\text{m}$, Fig. 4), with a high proportion of organic matter are accumulated, thus creating anoxic environments. The deepest parts of the lagoon ($>5 \text{ m}$) are then the ultimate catchment basins of the particles entering the lagoon, directly from the riverine inputs or indirectly, after several phases of settling and resuspension. Areas where particle settling is predominant (hollows) represent 57% of the lagoon surface. In these depressions, particles are no more reworked by hydrodynamics, explaining that less than 10% of particles entering the lagoon are exported toward the Atlantic Ocean (TASTET and GUIRAL, 1994).

REFERENCES

- ARFI, R., D. GUIRAL and M. BOUVY, 1993. Wind induced resuspension in a shallow tropical lagoon. *Estuar. Coast. Shelf Sci.*, 36: 587-604.
- BOUVY, M., ARFI, R. and D. GUIRAL, 1994. Short term variations of seston characteristics in a shallow tropical lagoon: effects of wind induced resuspension. *Neth. J. Aquat. Ecol.*, 28: 433-440.
- BENGTSSON, L., T. HELLSTRÖM and L. RAKOCZI, 1990. Redistribution of sediments in three swedish lakes. *Hydrobiologia*, 192: 167-181.
- CARPER, G.L. and R.W. BACHMANN, 1984. Wind resuspension of sediments in a prairie lake. *Can. J. Fish. Aquat. Sci.*, 41: 1763-1767.
- DEMERS, S., J.-C. THERRIault, E. BOURGET and A. BAH, 1987. Resuspension in the shallow sublittoral zone of a macrotidal estuarine environment: Wind influence. *Limnol. Oceanogr.*, 32: 327-339.
- DENMAN, K.L. and A.E. GARGETT, 1983. Time and space scales of vertical mixing and advection of phytoplankton in the upper ocean. *Limnol. Oceanol.*, 28: 801-815.
- FLODERUS, S. and L. PIHL, 1990. Resuspension in the Kattegat: impact of variation in wind climate and fishery. *Estuar. Coast. Shelf Sci.*, 31: 487-498.
- GUIRAL, D., 1992. L'instabilité physique, facteur d'organisation et de structuration d'un écosystème tropical saumâtre peu profond: la lagune Ebrié. *Vie Milieu*, 42: 73-92.
- HELLSTRÖM, T., 1991. The effect of resuspension on algal production in a shallow lake. *Hydrobiologia*, 213: 183-190.
- HOPKINSON, C.S. Jr, 1985. Shallow water benthic and pelagic metabolism: evidence of heterotrophy in the near-shore Georgia Bight. *Mar. Biol.*, 87: 19-32.
- KENNISH, M.J., 1986. Ecology of estuaries. Volume 1: Physical and chemical aspects. CRC Press, Boca Raton, 254 pp.
- LUETTICH, R.A., D.R.F. HARLEMAN and L. SOMLYODY., 1990. Dynamic behavior of suspended sediment concentrations in a shallow lake perturbed by episodic wind events. *Limnol. Oceanogr.*, 35: 1050-1067.
- PRESS, F. and R. SIEVER, 1986. Earth. 4th ed. W.H. Freeman and Co., New York.
- TASTET, J.P. and D. GUIRAL, 1994. Géologie et sédimentologie du système lagunaire Ebrié. In: *Environnement et ressources aquatiques de Côte d'Ivoire. Les milieux lagunaires*. Editions de l'ORSTOM: 35-55.
- TORRÉTON, J.P., M. BOUVY and R. ARFI, 1994. Diel fluctuations of bacterial abundance and productivity in a shallow eutrophic tropical lagoon. *Arch. Hydrobiol.*, 131: 79-92.

Address of the authors:

Centre de Recherches Océanologiques, ORSTOM, BP V18, Abidjan, Côte d'Ivoire

