

P. Paris
4

Regeneration of the tropical legume *Aeschynomene sensitiva* Sw. from root explants

Claudine Nef-Campa, Clémence Chaintreuil-Dongmo & Bernard Louis Dreyfus
Lab. de Microbiologie des Sols, Centre ORSTOM de Bel-Air, BP 1386. Dakar, Senegal

Received 11 April 1995; accepted in revised form 9 November 1995

Key words: Agar effects, direct shoot regeneration, Legume, stem nodule, water availability

Abstract

Regeneration of *Aeschynomene sensitiva* Sw. after callogenesis was obtained from small (2–5 mm long) root explants of 30-day-old seedlings aseptically cultivated on Murashige and Skoog medium supplemented with various concentrations of growth regulators. After 4 weeks, the best results were observed with 0.54 μM α -naphthaleneacetic acid and 2.22 μM benzyladenine. On this medium, the rate of regeneration depended on seedling age and agar concentration. The highest number of shoots per explant was obtained with small cuttings from 30-day-old seedlings grown on a medium containing 8 g l⁻¹ of agar. Regeneration success was also dependent on explant size. When longer explants (7–20 mm) were cut from the main root, direct regeneration was obtained in two weeks. These cuttings also generated shoots through callogenesis in four weeks but always in lower quantities than with direct regeneration, whatever the seedling age. Here also, the best regeneration was obtained with cuttings from 30-day-old seedlings maintained on a medium with 8 g l⁻¹ of agar. Regenerants were rooted on growth-regulator-free Murashige and Skoog medium and then acclimatized in a greenhouse. A better survival to transplantation was observed when plantlets were inoculated with the photosynthetic *Bradyrhizobium* strain ORS 278. Stem and root nodules developed on the inoculated plantlets and were able to fix nitrogen.

Abbreviations: BA – benzyladenine; 2,4-D – 2,4-dichlorophenoxyacetic acid; NAA – α -naphthaleneacetic acid; MS – Murashige & Skoog (1962) medium

Introduction

The genus *Aeschynomene* contains about 160 species of tropical legumes (Kretschmer & Bullock, 1980) spread throughout America and Africa, with a few species occurring in Southeast Asia and in the Pacific Islands. Some species are able to form both stem and root nitrogen-fixing nodules with *Rhizobia* (Dreyfus *et al.*, 1984; Loureiro *et al.*, 1994). The stem infection always occurs at predetermined sites which correspond to root primordia (Dreyfus & Dommergues, 1981). Most of the *Bradyrhizobium* strains capable of inducing stem nodules are of great significance because of their unusual ability to produce bacteriochlorophylla (Lorquin *et al.*, 1993) and to realize photosynthesis during heterotrophic growth (Eaglesham *et al.*, 1990). Moreover, these stem-nodulating legumes are

used as green manure in paddy-fields because of their high nitrogen accumulating potential, their tolerance to flooding and their fast growing ability (Alazard & Becker, 1987). No techniques of propagation have so far been described for this genus and particularly for *Aeschynomene sensitiva*. The paper reports an efficient system of tissue culture for rapid plantlet regeneration from root explants.

Fonds Documentaire ORSTOM

Cote: Bx 16787 Ex: 1

Materials and methods

Seeds of *Aeschynomene sensitiva* Sw. obtained from Casamance (South of Senegal) were surface sterilized (30 min in concentrated sulfuric acid followed by 5 rinses with distilled water) and then aseptically germinated by dropping them in test tubes on a filter paper




Table 1. Effect of combinations of 2,4-D and kinetin on callus development and shoot regeneration of small root cuttings from 30-day-old seedlings of *Aeschynomene sensitiva* Sw.

2,4-D (μM)	Kinetin (μM)	Explants with callus (%)	Calluses with shoots (%)	Number of shoot explant ⁻¹
0.45	0.46	13a ¹	0a	0a
	2.32	9a	0a	0a
2.26	0.46	100d	0a	0a
	2.32	82c	28b	0.64c
4.53	0.46	100d	91c	1.91d
	2.32	37b	25b	0.34b

¹Each value represents the mean of 3 replicates each with 10–12 root cuttings (2–5 mm). Callus formation was scored after 2 weeks and shoot regeneration after 4 weeks. Means within the same column followed by different letters are significantly different ($p \leq 0.05$) according to one way ANOVA analysis.

maintained at the surface of distilled water. Cultures were then incubated with a 14-h photoperiod ($77 \mu\text{mol m}^{-2} \text{s}^{-1}$ from Osram L36W/11 day-light fluorescent lamps) at $27\text{--}23^\circ\text{C}$ ($\pm 1^\circ\text{C}$) day-night temperatures. After 20 days of culture, the main root reached 3 to 6 cm in length. At this stage, both main and adventitious roots were cut into 2–5 mm sections referred to hereafter as small cuttings. After 30 to 45 days of culture, the main root reached 6 to 9 cm in length. The upper part of the main root (4 cm under the collar) was cut into 7–20 mm sections referred to hereafter as long cuttings. The lower part of this main root and the adventitious roots were treated as those from 20-day-old seedlings (small cuttings). The effect of some plant growth regulator combinations (2,4-D in combination with kinetin and NAA in combination with BA) on callogenesis and regeneration was investigated on small cuttings from 30-day-old seedlings. These explants were then cultivated in Petri dishes containing MS medium supplemented with agar at 8 g l^{-1} (Agar-Agar Prolabo). The growth regulators were tested at different concentrations as indicated in Tables 1 and

Table 2. Effect of combinations of NAA and BA on callus development and shoot regeneration of small root cuttings from 30-day-old seedlings of *Aeschynomene sensitiva* Sw.

NAA (μM)	BA (μM)	Explants with callus (%)	Calluses with shoots (%)	Number of shoot explant ⁻¹
0.54	0.44	0a ¹	0a	0a
	2.22	96d	83c	3.6c
2.68	0.44	21b	79c	1.25b
	2.22	8a	0a	0a
5.37	0.44	89d	28b	1.05b
	2.22	48c	9ab	0.2a

¹Each value represents the mean of 3 replicates each with 10–12 root cuttings (2–5 mm). Callus formation was scored after 2 weeks and shoot regeneration after 4 weeks. Means within the same column followed by different letters are significantly different ($p \leq 0.05$) according to one way ANOVA analysis.

4, 6, 8 and 10 g l^{-1} . Experiments were carried out in three replicates containing 10–12 explants each. After 2 and 4 weeks of culture, explants were observed for callogenesis and regeneration.

The regenerated plantlets were transferred to growth regulator free medium for rooting. When the rooted shoots reached the 7–8 leaf stage, one hundred (50 were obtained from small cuttings of 30 day-old seedlings and 50 were obtained directly from long cuttings) were distributed into three pots containing a mix of sterilized sand and vermiculite (1:1). Two pots were inoculated after 3 days of acclimatization with 1 ml of a 5-day-old photosynthetic *Bradyrhizobium* ORS 278 liquid culture containing 10^8 bacteria ml^{-1} (Vincent, 1970). The last pot was a control containing 30 non-inoculated plantlets. All plants were covered with polypropylene screw caps during the first week of culture. After transferring the plants into soil, the ability of stem and root nodules to fix nitrogen was estimated by the acetylene reduction assay (Hardy *et al.*, 1973).

2. The effect of explant age was evaluated by comparing shoot regeneration of small cuttings from 20-, 30- and 45-day-old seedlings grown on MS medium containing NAA and BA at 0.54 and 2.22 μM , respectively (NB medium). In addition, the influence of explant size

Results

Influence of growth regulators on shoot regeneration

Various combinations of growth regulators were tested

Table 3. Effect of seedling age on shoot regeneration from small root cuttings (2–5 mm) of *Aeschynomene sensitiva* Sw. after 4 weeks of culture on NB medium.

Seedling age (days)	Explants with shoots (%)	No of shoot explant ⁻¹
20	69 ± 18 ¹	1.46 ± 0.26
30	83 ± 11	3.60 ± 0.70
45	41 ± 9	1.48 ± 0.43

¹Means ± standard deviation of 3 replicates each with 10–12 root cuttings.




Fig. 2. Influence of the agar concentration on shoot production from small cuttings (2–5 mm) of root explants from 20 and 30-day-old seedlings of *Aeschynomene sensitiva* Sw. after 4 weeks of culture on NB medium. Means within the same series mentioned by different letters are significantly different ($p \leq 0.05$) according to one way ANOVA analysis.

ture (Fig. 1A, B and C). The highest callogenesis and shoot regeneration frequencies (100 and 91%, respectively) were observed on medium containing 2,4-D at 4.52 μM and kinetin at 0.46 μM (Table 1). The highest level of regeneration, in terms of shoot number per explant, was obtained on medium containing NAA at 0.54 μM and BA at 2.22 μM (Table 2). The average shoot number per callus was 3,6 which was 2-fold higher than for any other regenerating medium. This latter medium (called NB medium) was considered to be the most suitable for regeneration.

Effect of seedling age and agar concentration

The comparison of regeneration rates between small cuttings obtained from 20, 30 and 45-day-old seedlings cultivated on NB medium supplemented with 8 g l⁻¹ of agar showed that the explants from 30-day-old seedlings had the best ability to regenerate shoots.

Table 4. Effect of seedling age on shoot regeneration from long cuttings (7–20 mm) of main root of *Aeschynomene sensitiva* Sw. after 4 weeks of culture on NB medium.

Seedling age (days)	Direct regeneration (No. of shoot explant ⁻¹)	Regeneration through callogenesis (No. of shoot explant ⁻¹)
30	2.70 ± 0.51 ¹	0.84 ± 0.29
45	2.10 ± 0.40	0.41 ± 0.27

¹Means ± standard deviation of 3 replicates each with 8–10 root cuttings.




Fig. 3. Evolution of the number of shoots per explant obtained directly or after callogenesis on long cuttings (7–20 mm) of the main root from 30-day-old seedlings of *Aeschynomene sensitiva* Sw. cultivated on NB medium. Bar indicates the standard deviation of the mean.

These cuttings presented the highest rate of callogenesis (2-fold higher than for explants from 45-day-old seedlings) and the highest number of shoots per explant (Table 3). The regeneration rate was also dependent on the agar concentration of the culture medium. The best results were obtained with 6 and 8 g l⁻¹ of agar for cuttings from 20 and 30-day-old seedlings (Fig. 2). However, these results were significantly different ($p \leq 0.05$) only for 30-day-old seedlings.

Effect of explant size

The culture on the NB medium of long cuttings from the main root gave direct shoot formation on the root axis after the second week of culture (Fig. 1D, E). These explants were also able to form calluses located at the cut ends. On these calluses, bud formation was noticed after 3 weeks of culture and this allowed shoot isolation after 4 weeks. The seedling age seemed to have a low impact on direct regeneration (Table 4). However, as observed with small cuttings, the rate of




Fig. 4. Influence of the agar concentration on shoot production by direct regeneration or after callogenesis on long cuttings (7–20 mm) of the main root from 30-day-old seedlings of *Aeschynomene sensitiva* Sw. after 4 weeks of culture on NB medium. Means within the same series mentioned by different letters are significantly different ($p \leq 0.05$) according to one way ANOVA analysis.

regeneration after callogenesis decreased significantly when explants were cut from 45-day-old seedlings; the number of shoots formed per explant was 2-fold lower than with 30-day-old seedlings. As observed during the culture of small cuttings, the modification of the agar concentration induced variations on the rate of regeneration (Fig. 4). Direct regeneration was more effective with an agar concentration of 6 g l⁻¹ (average

Discussion

This study showed the capacity of *Aeschynomene sensitiva* Sw. to regenerate plantlets from root explants. Using NB medium (MS medium supplemented with NAA at 0.54 μ M and BA at 2.22 μ M), these root explants were able to develop shoots either directly or through callogenesis. Explant size and agar concentration seemed to be determining factors on the mode of regeneration. As for shoot regeneration from petunias leaf discs (Beck & Camper, 1991), a minimum size of explant seemed to be required to obtain direct regeneration. In this work, root cuttings shorter than 5 mm gave regenerated shoots only after callus production. Longer cuttings regenerated shoots directly on the root axis but for these explants, high agar concentrations promoted regeneration through callogenesis at the expense of direct regeneration. As the gelling agent concentration acted essentially on the water availability of the medium, with a matric potential 20% higher at 4 than at 10 g l⁻¹ (Owens & Wozniak, 1991), these results showed that a certain water availability and a certain explant length were needed to obtain direct regeneration. In return, the highest levels of regeneration after callogenesis were obtained for small cuttings from 30-day-old seedlings on media presenting a low water availability. The explant age also seemed to be a factor interfering with the level of regeneration. The best results for direct or indirect regeneration were obtained with root explants from 30-day-old seedlings. The root system of 20-day-old seedlings was not actually well developed. The small

with the *Bradyrhizobium* strain ORS 278. This could be attributed to a higher disponibility for the plant in nitrogen source, the symbiotic fixation constituting another nitrogen source for nitrogen metabolism. Root explants of *Aeschynomene sensitiva* Sw. seem to be suitable material for shoot regeneration and could be an interesting tool for plant transformation.

References

- Alazard D & Becker M (1987) *Aeschynomene* as a green manure for rice. *Plant and Soil*. 101: 141–142
- Beck MJ & Camper ND (1991) Shoot regeneration from petunia leaf discs as a function of explant size, configuration and benzyladenine exposure. *Plant Cell Tiss. Org. Cult.* 26: 101–106
- Chraïbi KMB, Latche A, Roustan JP & Fallot J (1991) Stimulation of shoot regeneration from cotyledons of *Helianthus annuus* by the ethylene inhibitor, silver and cobalt. *Plant Cell Rep.* 10: 204–207
- Dreyfus BL & Dommergues YR (1981) Nitrogen fixing nodules induced by rhizobium on the stems of the tropical legume *Sesbania rostrata*. *FEMS Microbiol. Lett.* 10: 313–317
- Dreyfus BL, Alazard D & Dommergues YR (1984) New and unusual microorganisms and niches. Stem-nodulating rhizobia. In: Klug MJ & Reddy CA (eds) *Current Perspectives in Microbial Ecology* (pp 161–169). American Society for Microbiology, Washington, DC
- Eaglesham ARJ, Ellis JM, Evans WR, Fleischman DE, Hungria M & Hardy RWF (1990) The first photosynthetic N₂-fixing *Rhizobium*: characteristics. In: Gresshoff PM, Roth LE, Stacey G & Newton WL (eds) *Nitrogen fixation: Achievements & Objectives*, (pp 805–811). Chapman & Hall New York.
- Hardy RWR, Burns RC & Holsten RD (1973) Application of the acetylene-ethylene assay for measurement of N₂ fixation. *Soil Biol. Biochem.* 5: 47–81
- Kretschmer AE Jr & Bullock RC (1980) *Aeschynomene* spp.: distribution and potential use. *Soil Crop Sci. Soc. Fla. Proc.* 39: 145–152
- Lorquin J, Molouba F, Dupuy N, Ndiaye S, Alazard D, Gillis M & Dreyfus B (1993) Diversity of photosynthetic *Bradyrhizobium* strains from stem nodules of *Aeschynomene* species. In: Palacios R, Mora J & Newton WE (eds) *New Horizons in Nitrogen Fixation* (pp 683–689). Kluwer Academic Publishers, Dordrecht, Boston, London
- Loureiro MF, Defaria SM, James EK, Pott A & Franco AA (1994) Nitrogen-fixing stem nodules of the legume, *Discolobium pulchellum* Benth. *New Phytol.* 128: 283–295
- Murashige T & Skoog F (1962) A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plant.* 15: 473–497
- Owens LD & Wozniak CA (1991) Measurement and effects of gel matrix potential and expressibility on production of morphogenic callus by cultured sugarbeet leaf discs. *Plant Cell Tiss. Org. Cult.* 26: 127–133
- Palmer CE (1992) Enhanced shoot regeneration from *Brassica campestris* by silver nitrate. *Plant Cell. Rep.* 11: 541–545
- Vincent JM (1970) A manual for the practical study of root nodule bacteria. *International Biological Handbook N° 15*. Blackwell Scientific Publications Ltd, Oxford

For information about current subscription rates and prices for back volumes for *Plant Cell, Tissue and Organ Culture*, ISSN 0167-6857

please contact one of the customer service departments of Kluwer Academic Publishers or return the form overleaf to:

Kluwer Academic Publishers, Customer Service, P.O. Box 322, 3300 AH Dordrecht, the Netherlands, Telephone (+31) 78 524 400, Fax (+31) 78 183 273, Email: services@wkap.nl

or

Kluwer Academic Publishers, Customer Service, P.O. Box 358, Accord Station, Hingham MA 02018-0358, USA, Telephone (1) 617 871 6600, Fax (1) 617 871 6528, Email: kluwer@world.std.com

Call for papers

Authors wishing to submit papers related to any of the themes or topics covered by *Plant Cell, Tissue and Organ Culture* are cordially invited to prepare their manuscript following the 'Instructions for Authors'. Please request these instructions using the card below.

Author response card

Plant Cell, Tissue and Organ Culture

I intend to submit an article on the following topic:

Please send me detailed 'Instructions for Authors'.

NAME : _____

INSTITUTE : _____

DEPARTMENT : _____

ADDRESS : _____

Telephone : _____

Telefax : _____

Email : _____

Library Recommendation Form

Route via Interdepartmental Mail

To the Serials Librarian at: _____

From: _____ Dept./Faculty of: _____

Dear Librarian,

I would like to recommend our library to carry a subscription to **Plant Cell, Tissue and Organ Culture**, ISSN 0167-6857 published by Kluwer Academic Publishers.

Signed: _____ Date: _____

Request for information about current subscription rates and prices for back volumes of
Plant Cell, Tissue and Organ Culture, ISSN 0167-6857

Please fill in and return to:

Kluwer Academic Publishers, Customer Service, P.O. Box 322, 3300 AH Dordrecht, the Netherlands

Kluwer Academic Publishers, Customer Service, P.O. Box 358, Accord Station, Hingham MA 02018-0358, USA

- Please send information about current program and prices
 Please send a free sample copy

NAME : _____
INSTITUTE : _____
DEPARTMENT : _____
ADDRESS : _____
Telephone : _____
Telefax : _____
Email : _____



REF. OPC

STAMP

Plant Cell, Tissue and Organ Culture
Kluwer Academic Publishers,
P.O. Box 17
3300 AA Dordrecht
The Netherlands



REF. OPC

TO : The Library
FROM: _____

V I A I N T E R D E P A R T M E N T A L M A I L