

091578

Evaporation and annual evolution of chloride concentrations in the water of a shallow Sahelian lake: Lake of Guiers (Senegal)

François-Xavier Cogels and Jean Yves Gac

Introduction

Lake Guiers is the only lake in Senegal. Located 50 km from the Atlantic coast (latitude 15°10' N; longitude 16°08' W), it is a narrow cavity 50 km long and 7 km wide, situated at 2 m b.s.l. (Fig. 1). The lake constitutes an important fresh water reserve and is connected to the lower Senegal River in a straight line by the Taoué canal. A system of sluice gates allows control of water exchange between the lake and the river.

The configuration of the lake changes considerably through time and space, under the effects of the river inflows, losses through evaporation and pumping. In its

average state, the lake has a depth of 1.5 m, an area of 225 km² and a volume of 350 millions m³.

The climate is sahelian and the data recorded by the Richard Toll station (north of the lake) show:

- an annual rainy season (July–September) with an average rainfall of 215 mm (1976–1991);
- an annual average air temperature of 27.6 °C with a minimum of 22 °C in January and a maximum of 30.3 °C in June;
- a low relative humidity (at 40%), a maximum deficit in the vapor tension in March, April and May and a minimum during the rainy season;
- an average windspeed at 2.4 m/s.





Fig. 1. Lake Guiers in West Africa.

Fonds Documentaire ORSTOM



010016941

Fonds Documentaire ORSTOM

0 100 km Cole: Bx 16941 Ex:

0368-0770/93/0025-0115 \$ 1.00

© 1993 E. Schweizerbart'sche Verlagsbuchhandlung, D-70176 Stuttgart




Fig. 2. Average monthly evaporation ($\text{mm} \cdot \text{d}^{-1}$) and standard deviation (s): 1976–1991.

Guier's hydrology was studied over the period 1976 to 1991 (COGELS et al. 1990), and is regulated by two major periods of the year:

- a *filling period* between August and October, with inflow from the Senegal river to the lake;
- a nine month *isolation period* without convergence between lake and river. The lake's water level decreases due to evaporation and the pumping out of water.

The lake's evaporation, methodology

The lack of data on the climate hinders direct calculation of lake's evaporation by the Penman's equation. The first stage of work was to ascertain the water balance of Guiers with maximum precision to determine the unknown quantity of the budget: evaporation. The monthly hydrological results were calculated over the 16-year period, from 1976 to 1991. All of the quantifiable parameters were measured directly, i.e. the volumes pumped (irrigation and drinkable water production) and released (by drains from irrigated cultures) and the pluviometric input. The surface and volume of the lake were calculated from the water level.

The evaporation is calculated separately during the two periods of the hydrological year:

- during the isolation period, evaporation is the only unknown parameter of the water balance and can be directly deduced.
- during the filling period (August and September) the lack of limnometric equipment on the Taoué canal hinders to quantify the river's inflow; two parameters, evaporation and river's inflow, are simultaneously the unknown terms of the water balance. Lake's evaporation in August and September is then estimated with reference to "Class A tank".

Results

On average, evaporation represents 80% of the annual water losses from the lake, various pumpings 15 to 18% and infiltrations 1.5%. 85% of the water supply come from the river Senegal, 8% from the pluviometry and 6 to 7% from the drains of the irrigated areas.

Fig. 2 shows the average monthly evaporation (192 months) calculated by dividing monthly evaporated volumes (from water balances) by the average surface areas of Lake Guiers at the same month. The evaporation follows a regular cycle, each year repeated, with a minimum in Decem-




Fig. 3. The evolution of the dissolved chlorides during the hydrological year of reference 1985–86.

ber, and a progressive rise up to May. As the rainy season approaches it gradually diminishes until September. The end of the rainy season is well characterized by a secondary maximum in October. The average of the total annual evaporation is 2.27 m (6.25 mm/day) with a standard deviation of 0.11 m (5%) and some extremes calculated figures of 2.14 m in 1979 and 2.58 m in 1983.

The data recorded for Guier's Lake are similar to those measured from other sahelian lakes: Lake Bam (POUYAUD 1986) and the sahelian pound of Oursi in Burkina Faso (CHEVALLIER et al. 1985), and Lake Chad (GAC 1980). The annual cycle is comparable with a simple seasonal decline due to the difference in longitude.

Evaporation and the annual change in water quality

The alternation of the 2 periods of the hydrological year induces an annual cycle in the water quality with:

- a phase of *mineralisation* (9 to 10 months) during the isolation period, due not only to the evaporation but also to the salted water released in the lake by the drainage system from salted soils of the irrigated areas.
- a phase of *dilution* (2 to 3 months) by the river inflow during the filling period.

The hydrological year 1985–86 can be considered as example to demonstrate the phenomenon repeated every year with more or less intensity according to the evaporation and the annual evolution of the lake's volume. This hydrological year of reference begins July 29, 1985 with the filling phase until September 23rd. The isolation period follows, and lasts until July 21, 1986.

Dissolved chlorides are analysed weekly at the NGnith station (on the west bank). This station can be considered as representative of the average quality of lake's water (COGELS et al. 1992). Fig. 3 shows the importance of the annual variations of dissolved chlorides.

During the lake's filling phase, the water dilution rate is 6.0, almost identical to that recorded during the isolation period (5.6). From one year to another, the figures vary in accordance with the importance of the water balance's parameters and with the evolution of the lake's volume during the 2 periods. The exact role of the evaporation in the annual process of mineralisation cannot be quantified directly since the salted drainage waters contribute equally to it. However it can be estimated monthly by dividing the volume evaporated by the mean volume of the lake during the same month. For the year 1985–86, and during the isolation period, the concentration rate of the water as a result of evaporation's influence was thus 3.85.

From 1976 to 1991, the average annual concentration rate of dissolved chloride was near 4.2 during the isolation period of the lake. During this period, evaporation induces an average concentration rate of 3.6, as 85 % of the total.

Conclusions

In sahelian region the total annual evaporation of a lake is near 2.25 m. In general, sahelian lakes are shallow lakes and are affected not only in terms of the volume of available water, but in terms of annual evolution of their physico-chemical quality too.

In the case of Guiers, the water quality shows very strong annual variations in mineralization, accentuated here by the high rate of annual water's renewal by River Senegal inflow and by the water from the drainage of the heavily salted cultures. Evaporation occurs on an average of 80 % during the mineralization process of solutions.

Paradoxically, environmental disturbances, hydrological as much as physicochemical, have had a beneficial effect in controlling the development of aquatic vegetation up to the present. However modifications in the lake's functioning have been recently engendered by the construction of the Diama Dam on Senegal River in 1985. The new hydrological river's conditions allow a more stable annual water level in the lake. A better dilution of the lake's water and the flush of the mineralized water of the Guier's southern region to the dead valley of Ferlo in the south cause a sensible reduction in the water salinity, especially since 2 years. The new conditions of the lake's environment would appear to be more favourable to the growth of the floating and rooted aquatic vegetation. Such pessimistic hypotheses have been confirmed by the excessive and recently recorded development of *Typha australis*, *Pistia stratiotes* and

Sphaenoclea zeylanica. The consequences on the lake's hydrology, hydrobiology and public health could be very harmful.

Acknowledgements

This research was financially supported by the Commission of the European Community (DG XII) and is a part of the project "Environment and Water Quality of the Senegal river" (EQUESEN) from the Science and Technology for Development Programme.

References

- CHEVALLIER, P., CLAUDE, J., POUYAUD, B. & BERNARD, A., 1985: Pluies et crues au Sahel. Hydrologie de la mare d'Oursi (Burkina Faso). – *Coll. Trav. et Doc. de l'ORSTOM* 190, 251 pp.
- COGELS, F.-X., GAC, J. Y., APPAY, J. L., EVORA, N. & LABROUSSE, B., 1990: Fonctionnement et bilans hydrologiques du lac de Guiers de 1976 à 1986. – *Doc. ORSTOM-Dakar*, 60 pp.
- COGELS, F.-X., NIANG, A. & GAC, J. Y., 1992: Définition d'une station de référence pour le suivi qualitatif des eaux du lac de Guiers. – *Doc. ORSTOM-Dakar*, 25 pp.
- GAC, J. Y., 1980: Géochimie du bassin du lac Tchad. – *Trav. et Doc. de l'ORSTOM* 123, 251 pp.
- POUYAUD, B., 1986: Contribution à l'évaluation de nappes d'eau libre en climat tropical sec. Exemples du lac de Bam, de la mare d'Oursi (Burkina Faso), du lac Tchad et d'acudes du Nordeste brésilien. – *Coll. Etudes et Thèses de l'ORSTOM*, Paris, 254 pp.

Authors' address:

ORSTOM, BP 1386, Dakar, Senegal.

This is a reprint of

Verhandlungen – Proceedings – Travaux

of the International Association for Theoretical and Applied Limnology, Congress in Barcelona 1992

If you are a member of this association you receive these "Proceedings" against your yearly membership fee of Swiss Francs 50.–

Please notice also the

Archiv für Hydrobiologie

an official organ of the International Association for Theoretical and Applied Limnology (SIL), and its Supplements.

Edited by: Prof. Dr. H. KAUSCH, Hamburg, Germany and
Prof. Dr. W. LAMPERT, Plön/Holstein, Germany.

Editorial Board: Prof. Dr. CAROLYN W. BURNS, Dunedin, New Zealand; Prof. Dr. Z. MACIEJ GLIWICZ, Warszawa, Poland; Prof. Dr. J. OVERBECK, Plön, Germany; Prof. Dr. ERIC PATÉE, Villeurbanne, France; Dr. COLIN S. REYNOLDS, Ambleside, Great Britain; Prof. Dr. ROBERT G. WETZEL, Tuscaloosa, Alabama, USA.

As a member of SIL you are entitled to receive this important journal at a special membership price.

True to its tradition this periodical serves freshwater research in the widest sense, including treatment of problems of brackish and seawater as far as they bear a relationship to limnology. It is the editors' aim to devote increased attention to ecology in association with experimental, above all physiological works corresponding to the more recent developments in limnology. Finally, it is intended that the "Archiv" should continue to form a bridge between theoretical and applied water research.

For details please write to the Publishers E. Schweizerbart'sche Verlagsbuchhandlung (Nägele u. Obermiller), Johannesstraße 3A, D-70176 Stuttgart, Germany.

If you are interested in

Archiv für Hydrobiology, Supplements

and in the special issues

Ergebnisse der Limnologie/Advances in Limnology

these are available also against a special membership price; for details please ask the Publishers (for address see above).

If you are interested in being a member of the International Association for Theoretical and Applied Limnology, please write to the General Secretary-Treasurer:

Prof. Dr. R. G. WETZEL, Department of Biological Sciences,
The University of Alabama, Tuscaloosa, Alabama 35487-0344/USA

how long
was the
case