

Clostridium peptidivorans sp. nov., a peptide-fermenting bacterium from an olive mill wastewater treatment digester

Tahar Mechichi,¹ Marie-Laure Fardeau,¹ Marc Labat,¹ Jean-Louis Garcia,¹ Frédéric Verhé¹ and Bharat K. C. Patel²

Author for correspondence: Bharat Patel. Tel: +61 417 726671. Fax: +61 7 38757800.
e-mail: bharat@genomes.sci.gu.edu.au

¹ Laboratoire IRD de Microbiologie, Université de Provence, CESB/ESIL, Case 925, 163 Avenue de Luminy, 13288 Marseille cedex 9, France

² School of Biomolecular and Biomedical Sciences, Faculty of Science, Griffith University, Brisbane, Queensland 4111, Australia

A new peptide-degrading, strictly anaerobic bacterium, designated strain TMC4^T, was isolated from an olive mill wastewater treatment digester. Cells of strain TMC4^T were motile, rod-shaped (5–10 × 0.6–1.2 µm), stained Gram-positive and formed terminal to subterminal spores that distended the cells. Optimal growth occurred at 37 °C and pH 7 in an anaerobic basal medium containing 0.5% Casamino acids. Arginine, lysine, cysteine, methionine, histidine, serine, isoleucine, yeast extract, peptone, Biotrypcase, gelatin and crotonate also supported growth, but not carbohydrates, organic acids or alcohols. The end-products of degradation were: acetate and butyrate from lysine and crotonate; acetate, butyrate, H₂ and CO₂ from Biotrypcase, gelatin and peptone; acetate, alanine, H₂ and CO₂ from cysteine; acetate, H₂ and CO₂ from serine, cysteine and yeast extract; acetate and formate from histidine; propionate from methionine; methyl 2-butyrate, H₂ and CO₂ from isoleucine; acetate and ethanol from arginine; and acetate, propionate, butyrate, methyl 2-butyrate, H₂ and CO₂ from Casamino acids. The DNA G+C content of strain TMC4^T was 31 mol%. Phylogeny based on 16S rRNA sequence analysis showed that strain TMC4^T was a member of the low-G+C-content Gram-positive genus *Clostridium*, with the closest relative being *Clostridium pascui* (sequence similarity of 96 %). Due to considerable differences in genomic and phenotypic properties between strain TMC4^T and those of its nearest relative, strain TMC4^T is proposed as a new species of the genus *Clostridium*, *Clostridium peptidivorans* sp. nov. Strain TMC4^T has been deposited in the DSMZ as strain DSM 12505^T.

Keywords: *Clostridium peptidivorans*, peptides, amino acids, olive mill wastewater, digester

INTRODUCTION

Olive mill wastewater is a dark liquid that contains high concentrations of biopolymers such as proteins, polysaccharides and polyphenols and is a cause of severe pollution in the Mediterranean basin. Agro-industrial wastewater has been shown to be treated successfully in anaerobic methanogenic digesters (Lettinga, 1995). Hamdi (1996) has proposed the use of similar systems to treat olive mill wastewater. In

order to understand the anaerobic digestion processes occurring in olive oil waste treatment, we have recently isolated and characterized several new species of anaerobic microbes that degrade aromatic compounds (Mechichi *et al.*, 1999). In this paper, we report the isolation and characterization of a new peptide-fermenting bacterium, *Clostridium peptidivorans* sp. nov.

Fonds Documentaire IRD

METHODS

Cote: B* 21588 Ex: 7

Sample source: Strain TMC4^T was enriched from samples collected from an anaerobic methanogenic digester (Sfax, Tunisia) fed with olive mill wastewater. The samples were

The GenBank accession number for the 16S rDNA sequence of strain TMC4^T is AF156796.



collected anaerobically using a N₂-flushed syringe and transferred directly to a sterile anaerobic serum bottle, which was maintained at room temperature until use. The digester temperature was set at 37 °C with no pH control.

Culture medium. The basal medium used for enrichment, isolation and cultivation contained (l⁻¹ distilled water): 1 g yeast extract, 1 g NH₄Cl, 0.3 g K₂HPO₄, 0.3 g KH₂PO₄, 0.6 g NaCl, 0.1 g CaCl₂·2H₂O, 0.2 g MgCl₂·6H₂O, 0.1 g KCl, 1.5 ml trace element solution (Widdel & Pfennig, 1981) and 1 mg resazurin. The pH was adjusted to 7 with 10 M KOH solution and the medium was made anaerobic using techniques described previously (Hungate, 1969; Macy *et al.*, 1972; Miller & Wolin, 1974).

Enrichment, isolation and cultivation. Enrichment, isolation and cultivation were performed in 5 ml basal medium containing 0.5% Casamino acids followed by incubation at 37 °C, as described previously (Mechichi *et al.*, 1999). Pure cultures were obtained by picking single well-isolated colonies from dilution series by using the roll-tube method (Hungate, 1969). Studies of optimum growth conditions and electron acceptor utilization were performed with 0.5% Casamino acids as the substrate.

Substrate utilization. Substrate utilization studies were performed in basal medium containing complex substrates at 0.5% (Biotrypcase, yeast extract, gelatin and peptone), carbohydrates at 20 mM (glucose, fructose, xylose, ribose, sorbose, sorbitol, sucrose, melibiose, raffinose, galactose, *myo*-inositol, lactose, cellobiose, mannitol, mannose, arabinose, arabitol, cellulose and xylan), organic acids at 20 mM (formate, acetate, propionate, butyrate, valerate, crotonate, fumarate, malonate, malate, lactate, citrate and succinate) or amino acids at 10 mM (arginine, alanine, proline, histidine, serine, aspartate, glycine, threonine, glutamate, glutamine, leucine, isoleucine, aspartate, asparagine, valine, methionine, lysine, tyrosine, phenylalanine and tryptophan), as described by Mechichi *et al.* (1999).

Cellular studies. Gram staining was performed using a kit purchased from Sigma. Light microscopy and electron microscopy were performed as described by Fardeau *et al.* (1997).

Analytical methods. Growth was determined by measuring changes in optical density at 580 nm by inserting anaerobic Hungate tubes directly into the cuvette holder of a Shimadzu model UV 160A spectrophotometer. Acetate, propionate, butyrate, formate, isobutyrate, ethanol and methyl 2-butyrate were measured by HPLC (Spectra Series 100 model; Thermo Separation Products) equipped with an Aminex HPX-87X 300 × 7.8 (inside diameter) mm column (Bio-Rad) connected to a differential refractometer (RID-6A; Shimadzu). Analysis was performed using a CR-6A Shimadzu integrator. The mobile phase was 0.0025 M H₂SO₄ at a flow rate of 0.6 ml min⁻¹ and the column temperature was 35 °C. The volume of the injection loop was 20 µl. Amino acid concentrations were determined by HPLC according to Moore *et al.* (1958). Sulfide was measured photometrically as colloidal CuS by using the method of Cord-Ruwisch (1985).

H₂ and CO₂ were measured using a gas chromatograph GC-8A (Shimadzu) equipped with a thermal conductivity detector, a C-R6A integrator (Shimadzu) and a Chromosorb WAW 80/100 mesh sp100 column (Alltech). N₂ at a pressure of 100 kPa was used as the carrier gas. The detector and the injector temperature was 200 °C, the column temperature was 150 °C.

Determination of G+C content. The G+C content of DNA was determined at the DSMZ by HPLC (Mesbah *et al.*, 1989). Non-methylated lambda DNA (Sigma) was used as the standard.

DNA extraction, amplification and sequencing of the 16S rDNA gene. DNA extraction, 16S rDNA amplification and sequencing were performed as described previously (Redburn & Patel, 1993; Andrews & Patel, 1996; Mechichi *et al.*, 1999). The universal primers Fd1 and Rd1 were used to obtain a PCR product of approximately 1.5 kb corresponding to base positions 28–1539 of the numbering of *Escherichia coli* 16S rDNA (Winker & Woese, 1991). The new sequence data that were generated, consisting of 1481 nucleotides, were aligned and an almost full-length consensus 16S rRNA gene sequence was assembled and checked for accuracy manually using the ae2 alignment editor (Maidak *et al.*, 1999). These were compared with other sequences in the GenBank database (Benson *et al.*, 1993) by using BLAST (Altschul *et al.*, 1997) and in the Ribosomal Database Project, version 7.0, by using SIMILARITY-RANK and SUGGEST-TREE (Maidak *et al.*, 1999). Reference sequences most closely related to our newly generated sequences were extracted from these databases and aligned. Positions of sequence and alignment uncertainty were omitted from the analysis. Pairwise evolutionary distances based on 1128 unambiguous nucleotides were computed by using DNADIST (Jukes and Cantor option) and neighbour-joining programs that form part of the PHYLIP suite of programs (Felsenstein, 1993). TREECON was used extensively for bootstrap analysis (Van de Peer & De Wachter, 1993).

RESULTS

Morphology

Several pure cultures were obtained and one of the cultures was designated strain TMC4^T and studied further. As strain TMC4^T did not grow on glucose, basal medium containing glucose was routinely used to check culture purity. Cells of strain TMC4^T were rod-shaped (0.6–1.2 × 5–10 µm). Spores were oval and were located terminally or subterminally, and they distended the cells (Fig. 1a). The cells stained Gram-positive and possessed a thick Gram-positive-type cell wall ultrastructure (Fig. 1b). Cells were motile and possessed peritrichous flagella.

Physiology

Strain TMC4^T was a strictly anaerobic and chemo-organotrophic bacterium. Growth occurred between 20 and 42 °C with the optimum temperature for growth at 37 °C. The optimum pH for growth was 7 and no growth occurred below pH 6.0 or above pH 9.0. NaCl was not required for growth, but was tolerated up to 4%.

Substrate utilization and fermentation end-products

Arginine, lysine, cysteine, methionine, histidine, serine, isoleucine, Biotrypcase, yeast extract and Casamino acids were fermented but carbohydrates, organic acids and other single amino acids (alanine, proline,

Fig
Tra
me

Table 1.

Amino a
asparagi
glutamin

Amino
Isoleuc
Histidi
Methic
Lysine
Argini
Cystei:
Serine
Casam

aspart:
leucine
alanin:
lowing
serine,
sine, 1
was n:
cultu:
histidi
valine
end-p:
acids
produ:
butyr:
CO₂
acetat

Interni




Fig. 1. (a) Phase-contrast micrograph of cells of strain TMC4^T in the exponential growth phase. Bar, 10 μm. (b) Transmission electron micrograph of the cell wall ultrastructure of strain TMC4^T, showing the presence of a cytoplasmic membrane (cm) and a thick, electron-dense cell wall layer (cw). Bar, 0.1 μm.


Table 1. Fermentation of amino acids by strain TMC4^T

Amino acids tested but not used were threonine, aspartate, asparagine, leucine, alanine, valine, glutamate, glycine, glutamine, phenylalanine, tyrosine, tryptophan and proline.

Amino acid	Fermentation end-products	Maximum OD ₅₈₀
Isoleucine	Methyl 2-butyrate, H ₂ , CO ₂	0.30
Histidine	Acetate, formate	0.32
Methionine	Propionate	0.40
Lysine	Acetate, butyrate	0.28
Arginine	Acetate, ethanol	0.41
Cysteine	Acetate, alanine, H ₂ , CO ₂	0.36
Serine	Acetate, H ₂ , CO ₂	0.37
Casamino acids	Acetate, butyrate, methyl 2-butyrate, H ₂ , CO ₂	0.55

aspartate, glycine, threonine, glutamate, glutamine, leucine, aspartate, asparagine, valine, tyrosine, phenylalanine and tryptophan) were not utilized. The following amino acids were utilized from Casamino acids: serine, glycine, leucine, phenylalanine, histidine, tyrosine, lysine, arginine and isoleucine. Strain TMC4^T was not able to perform the Stickland reaction when cultured on the following mixtures of amino acids: histidine/aspartate, histidine/proline, valine/cysteine, valine/proline and leucine/proline. The fermentation end-products from single amino acids and Casamino acids are shown in Table 1. The fermentation end-products from other substrates were: acetate and butyrate from crotonate; acetate, butyrate, H₂ and CO₂ from Biotrypcase, gelatin and peptone; and acetate, H₂ and CO₂ from yeast extract.

The carbon balance from cysteine fermentation was:



Electron acceptor utilization

During Casamino acid degradation, thiosulfate, but not sulfate, elemental sulfur, nitrate or fumarate, was reduced to H₂S and an increase was observed in the concentrations of the end-products acetate, propionate, butyrate, methyl 2-butyrate, H₂ and CO₂.

DNA base composition

The DNA G+C content of strain TMC4^T was 31 mol%, as determined by HPLC.

Phylogeny

Analysis of the 16S rDNA sequence showed that strain TMC4^T was a member of the low-G+C-content Gram-positive bacteria and a member of cluster I (Collins *et al.*, 1994) and, hence, was not related to most of the previously described peptide-fermenting bacteria (Baena *et al.*, 1998, 1999a, 1999b). Cluster I is similar to the rRNA group I of Johnson & Francis (1975), consisting of some 78 species, including *Clostridium pascui* and *Clostridium acetireducens*, and is currently the largest group of the low-G+C-content Gram-positive clostridia. Further in-depth phylogenetic analysis indicated that *C. pascui* (sequence similarity of 96%), rather than *C. acetireducens* (sequence similarity of 93%), was the closest relative of strain TMC4^T. A dendrogram showing the relationship of strain TMC4^T to *C. pascui* and other representative members of cluster I is depicted in Fig. 2.




Fig. 2. Unrooted phylogenetic dendrogram based on 16S rRNA sequence data indicating the position of *C. peptidovorans* strain TMC4^T within the radiation of representative members of the genus *Clostridium* (cluster I) according to Collins *et al.* (1994). All sequences used in the analysis were obtained from the Ribosomal Database Project, version 7 (Maidak *et al.*, 1999). The triangle indicates representative members of cluster II (*Clostridium histolyticum* ATCC 19401^T, *Clostridium limosum* ATCC 25620^T and *Clostridium proteolyticum* ATCC 49002^T). Evolutionary distances (based on 1128 unambiguous nucleotides) and bootstrap analysis (100 data sets) were computed using programs that form part of the PHYLIP package (Felsenstein, 1993). Only values greater than 90% were considered significant and are therefore reported. Bar, 5 nucleotide substitutions per 100 nucleotides.

DISCUSSION

The trait of degrading proteinaceous compounds but not carbohydrates is a characteristic shared by strain TMC4^T with only a few members of the low-G+C-content Gram-positive bacteria, including *C. pascui*, *C. acetireducens*, *Clostridium litorale*, *Clostridium sticklandii*, *Clostridium aminophilum*, *Clostridium hydroxybenzoicum*, *Peptostreptococcus anaerobius*, *Acidaminobacter hydrogenoformans*, *Eubacterium acidaminophilum*, *Anaeromusa acidaminophila* (formerly '*Selenomonas acidaminophila*'), *Dethiosulfovibrio peptidovorans*, *Aminobacterium colombiense* and *Aminomonas paucivorans*. However, with the exception of *C. pascui*, with which it shares a sequence similarity of 96%, all the other bacteria are phylogenetically distantly related. In addition, strain TMC4^T and *C. pascui* share the ability to ferment histidine but not carbohydrates, have similar optimum conditions for growth (37 °C and pH 7), form spores and are motile. However, numerous phenotypic differences also exist between the two taxa (Table 2). *C. pascui* stained

Gram-negative and fermented glutamate, whereas strain TMC4^T stained Gram-positive and did not ferment glutamate. Strain TMC4^T is a proteolytic bacterium and utilized a large number of proteinaceous substrates, whereas *C. pascui* is not proteolytic and used only amino acids. The G+C content of TMC4^T is 31 mol%, whereas that of *C. pascui* is much lower (27 mol%). The habitats from which these two strains were isolated are also different; strain TMC4^T was isolated from an anaerobic digester whereas *C. pascui* was isolated from intestinal tracts of donkeys. A different spectrum of end-products from histidine fermentation was produced by strain TMC4^T (acetate and formate) when compared with *C. pascui* (acetate, butyrate, ethanol, CO₂ and H₂).

On the basis of the phenotypic, genotypic and phylogenetic evidence presented and based on the current taxonomic guidelines, we propose that strain TMC4^T should be designated as a new member of cluster I, which encompasses the true members of the genus *Clostridium* (Collins *et al.*, 1994).

Table 2. Characteristics of strain TMC4^T and *C. pasculi*Data for *C. pasculi* were taken from Wilde *et al.* (1997).

Characteristic	Strain TMC4 ^T	<i>C. pasculi</i>
Habitat	Olive mill wastewater digester	Intestinal tract of donkey
Morphology	Rods, 5–10 × 0.6–1.2 µm	Rods, 3.2–8 × 0.75–1 µm
Spores	Oval, terminal to subterminal; distend the cell	Elliptical, subterminal; distend the cell
Temperature for growth (°C):		
Range	20–42	10–43
Optimum	37	37–40
pH for growth:		
Range	6–9	5.5–9
Optimum	7.0	6.4–7.8
Substrates utilized	Arginine, lysine, cysteine, methionine, histidine, serine, isoleucine, Casamino acids, gelatin, yeast extract, peptone, Biotrypcase	Glutamate, histidine, ribose (weakly)
End-products of histidine fermentation	Acetate, formate	Acetate, <i>n</i> -butyrate, ethanol, CO ₂ , H ₂
DNA G+C content (mol%)	31	27

Description of *Clostridium peptidivorans* sp. nov.

Clostridium peptidivorans (pep.ti.di.vo'rans. N.L. n. *peptidum* peptide; L. v. *vorare* to devour; N.L. part. adj. *peptidivorans* peptide consuming).

Cells are motile, stain Gram-positive, are rod-shaped (0.6–1.2 × 5–10 µm) and form spores. Spores are terminal to subterminal and oval and distend the cell. Strictly anaerobic and grows optimally at 37 °C over a temperature range of 20–42 °C. No growth is observed at or below 15 °C or above 45 °C. The pH range for growth is between 6 and 9 with an optimum at 7.0. Grows in the presence of NaCl concentrations less than 4%. Grows on Biotrypcase, yeast extract, Casamino acids, gelatin, peptone, arginine, lysine, cysteine, methionine, histidine, serine, isoleucine and crotonate but not on carbohydrates (glucose, fructose, xylose, sorbose, sorbitol, sucrose, melibiose, raffinose, galactose, *myo*-inositol, sucrose, lactose, cellobiose, mannitol, mannose, arabinose, arabitol, raffinose, cellulose or xylan ribose), organic acids (formate, acetate, propionate, *n*-butyrate, valerate, fumarate, malonate, malate, lactate, citrate or succinate) or other amino acids (alanine, proline, aspartate, glycine, threonine, glutamate, glutamine, leucine, aspartate, asparagine, valine, tyrosine, phenylalanine or tryptophan). Thiosulfate is reduced but not sulfate, elemental sulfur, nitrate or fumarate. The fermentation end-products are: acetate and butyrate from lysine and crotonate; acetate, butyrate, H₂ and CO₂ from Biotrypcase, gelatin and peptone; acetate, alanine, H₂ and CO₂ from cysteine; acetate, H₂ and CO₂ from serine, cysteine and yeast extract; acetate and formate from histidine; propionate from methionine; methyl 2-butyrate, H₂ and CO₂ from isoleucine; acetate and

ethanol from arginine; and acetate, propionate, butyrate, methyl 2-butyrate, H₂ and CO₂ from Casamino acids. The G+C content of DNA is 31 mol%, as determined by HPLC. The type strain is TMC4^T (= DSM 12505^T). Isolated from an olive mill wastewater treatment digester.

ACKNOWLEDGEMENTS

Financial assistance in part to B. K. C. P. from the Australian Research Council is gratefully acknowledged.

REFERENCES

- Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W. & Lipman, D. J. (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25, 3389–3402.
- Andrews, K. T. & Patel, B. K. C. (1996). *Fervidobacterium gondwanense* sp. nov., a new thermophilic anaerobic bacterium isolated from nonvolcanically heated geothermal waters of the Great Artesian Basin of Australia. *Int J Syst Bacteriol* 46, 265–269.
- Baena, S., Fardeau, M.-L., Labat, M., Ollivier, B., Thomas, P., Garcia, J.-L. & Patel, B. K. C. (1998). *Aminobacterium colombiense* gen. nov. sp. nov., an amino acid-degrading anaerobe isolated from anaerobic sludge. *Anaerobe* 4, 241–250.
- Baena, S., Fardeau, M.-L., Woo, T. H. S., Ollivier, B., Labat, M. & Patel, B. K. C. (1999a). Phylogenetic relationships of three amino-acid-utilizing anaerobes, *Selenomonas acidaminovorans*, '*Selenomonas acidaminophila*' and *Eubacterium acidaminophilum*, as inferred from partial 16S rDNA nucleotide sequences and proposal of *Thermanaerovibrio acidaminovorans* gen. nov., comb. nov. and *Anaeromusa acidaminophila* gen. nov., comb. nov. *Int J Syst Bacteriol* 49, 969–974.
- Baena, S., Fardeau, M.-L., Ollivier, B., Labat, M., Thomas, P., Garcia, J.-L. & Patel, B. K. C. (1999b). *Aminomonas paucivorans*

- gen. nov., sp. nov., a mesophilic, anaerobic, amino-acid-utilizing bacterium. *Int J Syst Bacteriol* 49, 975–982.
- Benson, D., Lipman, D. J. & Ostell, J. (1993). GenBank. *Nucleic Acids Res* 21, 2963–2965.
- Collins, M. D., Lawson, P. A., Willems, A., Cordoba, J. J., Fernandez-Garayzabal, J., Garcia, P., Cai, J., Hippe, H. & Farrow, J. A. E. (1994). The phylogeny of the genus *Clostridium*: proposal of five new genera and eleven new species combinations. *Int J Syst Bacteriol* 44, 812–826.
- Cord-Ruwisch, R. (1985). A quick method for the determination of dissolved and precipitated sulfides in cultures of sulfate-reducing bacteria. *J Microbiol Methods* 4, 33–36.
- Fardeau, M.-L., Ollivier, B., Patel, B. K. C., Magot, M., Thomas, P., Rimbault, A., Rocchiccioli, F. & Garcia, J.-L. (1997). *Thermotoga hypogea* sp. nov., a xylanolytic, thermophilic bacterium from an oil-producing well. *Int J Syst Bacteriol* 47, 1013–1019.
- Felsenstein, J. (1993). PHYLIP (Phylogenetic Inference Package) version 3.51c. Distributed by the author. Department of Genetics, University of Washington, Seattle, WA, USA.
- Hamdi, M. (1996). Anaerobic digestion of olive mill wastewaters. *Process Biochem* 31, 105–110.
- Hungate, R. E. (1969). A roll-tube method for the cultivation of strict anaerobes. *Methods Microbiol* 136, 194–198.
- Johnson, J. L. & Francis, B. S. (1975). Taxonomy of the clostridia: ribosomal ribonucleic acid homologies among the species. *J Gen Microbiol* 88, 229–244.
- Lettinga, G. (1995). Anaerobic digestion and wastewater treatment systems. *Antonie Leeuwenhoek* 67, 3–28.
- Macy, J. M., Snellen, J. E. & Hungate, R. E. (1972). Use of syringe methods for anaerobiosis. *Am J Clin Nutr* 25, 1318–1323.
- Maidak, B. L., Cole, J. R., Parker, C. T., Jr & 11 other authors (1999). A new version of the RDP (Ribosomal Database Project). *Nucleic Acids Res* 27, 171–173.
- Mechichi, T., Labat, M., Patel, B. K. C., Woo, T. H. S., Thomas, P. & Garcia, J.-L. (1999). *Clostridium methoxybenzovorans* sp. nov., a new aromatic *o*-demethylating homoacetogen from an olive mill wastewater treatment digester. *Int J Syst Bacteriol* 49, 1201–1209.
- Mesbah, M., Premachandran, U. & Whitman, W. B. (1989). Precise measurement of the G + C content of deoxyribonucleic acid by high-performance liquid chromatography. *Int J Syst Bacteriol* 39, 159–167.
- Miller, T. L. & Wolin, M. J. (1974). A serum bottle modification of the Hungate technique for cultivating obligate anaerobes. *Appl Microbiol* 27, 985–987.
- Moore, S., Spackman, D. H. & Stein, W. H. (1958). Chromatography of amino acids on sulfonated polystyrene resins: an improved system. *Anal Chem* 30, 1158–1190.
- Redburn, A. C. & Patel, B. K. C. (1993). Phylogenetic analysis of *Desulfotomaculum thermobenzoicum* using polymerase chain reaction-amplified 16S rRNA-specific DNA. *FEMS Microbiol Lett* 113, 81–86.
- Van de Peer, Y. & De Wachter, R. (1993). TREECON: a software package for the construction and drawing of evolutionary trees. *Comput Appl Biosci* 9, 177–182.
- Widdel, F. & Pfennig, N. (1981). Studies on dissimilatory sulfate-reducing bacteria that decompose fatty acids. I. Isolation of new sulfate-reducing bacteria enriched with acetate from saline environments. Description of *Desulfobacter postgatei* gen. nov., sp. nov. *Arch Microbiol* 129, 395–400.
- Wilde, E., Collins, M. D. & Hippe, H. (1997). *Clostridium pascui* sp. nov., a new glutamate-fermenting sporeformer from a pasture in Pakistan. *Int J Syst Bacteriol* 47, 164–170.
- Winker, S. & Woese, C. R. (1991). A definition of the domain *Archaea*, *Bacteria* and *Eucarya* in terms of small subunit ribosomal RNA characteristics. *Syst Appl Microbiol* 13, 161–165.