

STARFISH SAPONINS, XIII. OCCURRENCE OF NODOSOSIDE IN THE STARFISH
ACANTHASTER PLANCI AND *LINCKIA LAEVIGATA*¹

L. MINALE,* C. PIZZA, R. RICCIO, F. ZOLLO

Istituto di Chimica Biorganica, Università, Via L. Rodinò, 22, I-80138 Naples, Italy

J. PUSSET,

Laboratoire des Plantes Médicinales, CNRS, B.P. 643, Nouméa, New Caledonia

and P. LABOUTE

Centre ORSTOM, B.P. A5, Nouméa, New Caledonia
with the technical collaboration of M. Marly, Centre ORSTOM,
and M. Pusset, Laboratoire de Plantes Médicinales, CNRS, Nouméa

Starfish have been a rich source for discovery of marine steroidal glycosides with new structures (1). Recently, we have described a novel group of steroidal glycosides that are composed of polyhydroxylated (five to six hydroxyl) sterol aglycones and a carbohydrate portion glycosidally attached at C-24 of the steroid, from the starfish *Protoreaster nodosus* (2) and *Hacelia attenuata* (3,4). In particular, *Protoreaster nodosus* has yielded 24-O-[2-O-methyl-β-D-xylopyranosyl-(1→2)-α-L-arabinofuranosyl]-5α-cholestane-3β,5,6β,8,15α,24ξ-hexol, which we named nodososide (2).

Continuing with our investigation of biologically active marine steroidal glycosides from echinoderms, we have been working on saponins of the Pacific starfishes, *Acanthaster planci* (L.), whose major asterosaponin, thornasteroside A, was fully characterized a few years ago by Kitagawa and Kobayashi (5), and *Linckia laevigata* (L.), and have isolated from both sources, in small amounts, a steroidal glycoside identical with nodososide.

EXPERIMENTAL

ANIMAL COLLECTION AND EXTRACTION.—*L. laevigata* was collected in March 1981, and *A. planci* in September 1982, off Nouméa, New Caledonia. Starfishes were chopped and extracted (5 h) with H₂O and the extracts lyophilized. *L. laevigata* (5 kg, fresh animals) gave 129 g of crude extract; *A. planci* (5 kg) gave 850 g of crude extract.

ISOLATION OF THE NODOSOSIDE.—*L. laevigata*: The aqueous lyophilized extract was extracted with light petroleum (bp 40–70°), then with CHCl₃, followed by MeOH-CHCl₃ (1:9), MeOH, and H₂O-MeOH (1:1). The MeOH-CHCl₃ extract gave 23 g of residue, which was separated by a silica gel short-column chromatography (300 g SiO₂). Fractions (250 ml) were collected; fractions 1–8 were eluted with MeOH-CHCl₃ (1:9), fractions 9–11 with MeOH-CHCl₃ (2:8), and fractions 12–13 with MeOH-CHCl₃ (3:7). Fraction 9 gave 187 mg of residue that was chromatographed by hplc on a μ-bondapak C-18 column (7.8 mm×30 cm, flow rate 5.5 ml/min⁻¹) using 35% H₂O in MeOH. The major peak collected between 18 and 21 min after injection contained 8 mg of nodososide.

A. planci: The same procedure was used to yield 9 mg of nodososide.

Nodososide, [α]_D²⁰ = -20.3° (from *L. laevigata*), -19.5° (from *A. planci*) (lit. [α]_D²⁰ = -21.3°, 2), was identified by 270 MHz pmr and authentic sample comparison by hplc and SiO₂-tlc.

Full details of the identification of the compound are available on request to the senior author.

ACKNOWLEDGMENTS

We thank the divers, G. Bargibant, J.L. Menou, and P. Tirard of the Centre ORSTOM de Nouméa for collection of the animals.

LITERATURE CITED

1. L. Minale, C. Pizza, R. Riccio, and F. Zollo, *Pure Appl. Chem.*, **54**, 1935 (1982).
2. R. Riccio, L. Minale, C. Pizza, F. Zollo, and J. Pusset, *Tetrahedron Lett.*, 2899 (1982).
3. L. Minale, C. Pizza, R. Riccio, and F. Zollo, *Experientia*, **39**, 567 (1983).
4. L. Minale, C. Pizza, R. Riccio, and F. Zollo, *Experientia*, **39**, 569 (1983).
5. I. Kitagawa and M. Kobayashi, *Chem. Pharm. Bull.*, **26**, 1864 (1978).

Received 21 June 1983

¹This contribution is part of the Progetto Finalizzato "Chimica fine e secondaria," CNR, Rome, Italy.

