

**AN OVERVIEW OF THE 1997 ACTIVITIES OF THE  
SEMI-ARID LAND-SURFACE-ATMOSPHERE (SALSA) PROGRAM**

D. C. Goodrich (1)\*, A. Chehbouni (2), B. Goff (1), B. MacNish (3), T. Maddock (3), S. Moran (4), W.J. Shuttleworth (3), D.G. Williams (3), J.J. Toth (1), C. Watts (12), L.H. Hipps (5), D.I. Cooper (6), J. Schieldge (14), Y.H. Kerr (7), H. Arias (12), M. Kirkland (6), R. Carlos (6), W. Kepner (17), B. Jones (17), R. Avissar (19), A. Bégué (8), G. Boulet (2), B. Branat (18), I. Braud (15), J.P. Brunel (2), L.C. Chen (9), T. Clarke (4), M.R. Davis (13), J. Dautat (8), H. DeBruin (10), G. Dedieu (7), W.E. Eichinger (9), E. Elguero (2), J. Everitt (13), J. Garatuza-Payan (3), A. Garibay (12), V.L. Gempko (3), H. Gupta (3), C. Harlow (3), O. Hartogensis (10), M. Helfert (4), C. Holfield (4), D. Hymer (2), A. Kahle (14), T. Keefer (1), S. Krishnamoorthy (9), J-P. Lhomme (2), D. Lo Seen (8), D. Luquet (8), R. Marsett (1), B. Monteny (2), W. Ni (4), Y. Nouvellon (8), R. Pinker (20), C. Peters (3), D. Pool (16), J. Qi (4), S. Rambal (11), H. Rey (8), J. Rodriguez (12), E. Sano (3), S.M. Schaeffer (3), M. Schulte (3), R. Scott (3), X. Shao (6), K.A. Snyder (3), S. Sorooshian (3), C.L. Unkrich (1), M. Whitaker (3), I. Yucel (3)

(1) USDA-ARS, Tucson, AZ; (2) ORSTOM/IMADES, Hermosillo, Sonora, Mexico; (3) Univ. of Arizona, Tucson, AZ, USA; (4) USDA-ARS Water Cons. Lab., Phoenix, Arizona, USA; (5) Utah St. Univ., Logan, UT, USA; (6) Los Alamos Nat. Lab., Los Alamos, NM, USA; (7) CESBIO, Toulouse, France; (8) CIRAD, Montpellier, France; (9) Univ. of Iowa, Iowa City, IA USA; (10) AUW, Wageningen, The Netherlands; (11) CEFE, CNRS, Montpellier, France; (12) IMADES, Hermosillo, Sonora, Mexico; (13) USDA-ARS, Weslaco, TX, USA; (14) Jet Prop. Lab., Pasadena, CA, USA; (15) LTHE, BP 53, 38041 Grenoble, France; (16) USGS-WRD, Tucson, AZ, USA; (17) US-EPA, Las Vegas, NV, USA; (18) Audubon Research Ranch, Elgin, AZ USA; (19) Rutgers Univ., New Brunswick, NJ, USA; (20) Univ. of Maryland, College Park, MD, USA

## 1. INTRODUCTION

The primary objective of the Semi-Arid Land-Surface-Atmosphere (SALSA) Program is to understand, model and predict the consequences of natural and human-induced change on the basin-wide water balance and ecological diversity of semiarid regions at event, seasonal, interannual, and decadal time scales. SALSA is a long-term program whose current research and integrated measurement efforts are focused on the San Pedro River basin which originates in northern Sonora, Mexico and flows north into southeastern Arizona. This paper provides an overview of the 1997 SALSA Program elements and data collection activities. It serves as an introduction to a number of other, more detailed, SALSA papers presented in the special session on "Integrated Observations of Semi-Arid Land-Surface-Interactions."

## 2. 1997 SALSA SCIENTIFIC OBJECTIVES

1997 SALSA activities are part of a longer term (3 to 10 year effort) to address the primary objective developed from initial program goals (Goodrich, 1994) and an international workshop (Wallace, 1995). The 1997 SALSA objectives, focused on priorities established at the workshop and within logistical and monetary constraints are:

- 1) To improve the diagnosis of surface fluxes used in atmospheric models with grid spacings of several kilometers and compare remote and *in-situ* observations with real-time model runs;
- 2) To initiate the development and validation of a coupled soil-vegetation-atmosphere transfer (SVAT) and vegetation growth model for semi-arid regions

that will assimilate remotely sensed data with several years of observed data;

- 3) To conduct *in-situ* and remote measurements to: a) quantify and develop models for groundwater, surface water, and evapotranspiration interactions on a seasonal basis; b) identify plant water sources; and c) identify plant function and atmospheric controls on a semi-arid riparian system consisting of mesquite, sacaton, and cottonwood/willow vegetation communities.
- 4) To develop and validate aggregation schemes with data over very highly heterogeneous surfaces; and,
- 5) To develop a multi-scale system of landscape pattern indicators using remotely sensed data to estimate current status, trend and changes in ecological condition.

Other factors involved in the selection of the above objectives include the desire to maintain and initiate long term observations and multi-disciplinary research in the San Pedro Basin, and to assist in addressing a number of pressing socio-economic concerns in the region. Long-term monitoring and research is essential to capture a range of both seasonal and interannual climatic variations and their associated impacts on basin water resources and ecology. It is because of the long history of monitoring and multidisciplinary experimentation in the San Pedro Basin and the long-term nature of SALSA that the NASA Earth Observing System ASTER (Adv. Spaceborne Thermal Emission and Reflection Radiometer) team has selected the San Pedro as a regional study and validation area. Research supporting development and validation of ASTER aircraft simulator data products was also undertaken as part of 1997 SALSA objectives.

\* David C. Goodrich, USDA-ARS, 2000 E. Allen Rd., Tucson, AZ 85719; e-mail:goodrich@tucson.ars.ag.gov



Fonds Documentaire IRD

Cote : B\* 23310 Ex : X

*unkrich*

### 3. THE STUDY AREA

The Upper San Pedro Basin (USPB) was identified as the focal area for initial SALSA research during the 1995 workshop. The riparian system in the U.S. portion of the USPB is the first Congressionally designated National Riparian Conservation Area. The basin embodies a variety of characteristics which make it an exceptional outdoor laboratory to address a large number of scientific challenges in arid and semi-arid hydrology, meteorology, ecology, and the social and policy sciences. The area represents a transition between the Sonoran and Chihuahuan deserts with significant topographic and vegetation variation, and a highly variable climate. It is an international basin spanning the Mexico-United States Border with significantly different cross border legal and land use practices. The upper and middle portions of the basin depicted in Figure 1 have a drainage area of 7610 km<sup>2</sup> at the U.S. Geological Survey gaging station at Reddington, Arizona with approximately 1800 km<sup>2</sup> in Mexico. Elevations range from roughly 1100 to 2900 m. A color Landsat MSS image illustrating the basin's heterogeneity and cross-border land use differences is available on the SALSA home page ([http://www.hwr.arizona.edu/salsasalsa\\_1.html](http://www.hwr.arizona.edu/salsasalsa_1.html))

The annual rainfall ranges from around 300 mm to 750 mm. Approximately 65% of this typically occurs during the July through September monsoon season from high intensity air-mass convective thunderstorms. Roughly 30% comes from less intense winter frontal systems. Potential evapotranspiration is estimated at more than ten times annual rainfall at lower elevations (Renard et al., 1993). Interannual climate variability is also high with a demonstrated linkage to the El Niño-Southern Oscillation (Woolhiser et al., 1993). Major vegetation communities include desert shrub-steppe, riparian, grasslands, oak savannah, and ponderosa pine. In portions of the basin all of these vegetation types are contained within a 20 km span. The USPB supports the second highest known number of mammal species in the world and the riparian corridor provides habitat for more than 300 bird species.

From a socio-economic perspective, great concern exists regarding the long-term viability of the San Pedro riparian system and ranching in the face of continued population growth. Groundwater sustains the riparian system in the United States and also much of the ranching industry in the Mexican portion of the San Pedro.

The threat of excessive groundwater pumping to this riparian system has prompted the first application of international environmental law within the U.S. via the North American Free Trade Agreement (see Arias et al., 1998 - this issue).

### 4. METHODS

To address the *first objective* the Regional Atmospheric Modeling System (RAMS) mesoscale atmospheric model was run using variable surface parameters appropriate for the region in a diagnostic mode in near real-time for all of 1997 with a 4 x 4 kilometer grid spacing over the entire San Pedro Basin (see Figure 1). Analyses and 6-hour forecasts from the operational meso-Eta model were used to generate




Figure 1: Upper San Pedro Basin

boundary conditions and nudging tendencies for RAMS. For comparative purposes, model output was saved every 15 minutes for the grid cells closest to the experimental surface observations shown in Figure 1. In addition, all model fields were saved every 3 hours so assessments

could be made of the value of integrating and assimilating both remotely sensed and *in-situ* observations (Toth et al., this issue).

Experimental surface observations were obtained from a variety of existing data collection networks. These include intensive monitoring at the 148 km<sup>2</sup> USDA-ARS Walnut Gulch Watershed (85 raingages, 30 runoff measurement sites, two sites with energy and CO<sub>2</sub> flux, meteorology, vegetation and soil moisture measurements; Renard et al., 1993), meteorological data collection by the US Army at Ft. Huachuca, and five USGS stream gaging stations.

In addition, several new intensive research and data collection sites were established during 1997 in areas having sparse desert grass cover in Mexico at Morelos and Zapata (see Figure 1). A wide variety of *in-situ* and remotely sensed data are being collected at these sites to address *objective two*. These measurements include meteorology, energy and CO<sub>2</sub> fluxes, soil moisture, vegetation sampling, and surface reflectances (Chehbouni et al., this issue). Stable isotopes of water were also collected over a board transect to examine the competition for water between grass and shrubs so that the potential extension of shrub invasion into grasslands may be estimated. Extensive large scale measurements, on a scale commensurate with the resolution of the RAMS model, of sensible heat flux are also being taken at these relatively uniform sites with a scintillometer (Chehbouni et al., this issue).

In addition to the broader scale measurements noted above, an intensive set of groundwater, surface water, isotope, energy flux and plant transpiration measurements were concentrated in the Lewis Springs section of the San Pedro Riparian corridor to address *objective three*. This stream section is a "gaining reach" where groundwater contributes water to the stream. The vegetation type and structure in the Lewis Springs area is relatively representative of vegetation throughout the 50 kilometer long San Pedro National Riparian Conservation Area. This riparian system typically consists of a narrow (20 - 100m) cottonwood/willow forest gallery surrounded on either side by riparian grasses and mesquite thickets. These grow upon the ancient flood plain which varies in width from several hundred to between one and two kilometers. A schematic of a typical section of riparian corridor geometry is illustrated in the lower insert of Figure 2. A primary goal is to quantify the surface water, groundwater, and evapotranspiration fluxes into and out of the control volume as depicted in this figure.

The network of instrumentation installed at Lewis Springs to address this goal and the other aspects of *objective three* is illustrated in the main portion Figure 2 (Note: An illustration of the vegetation classification for this area is presented in Moran et al., this issue). Continuous measurements of water levels in the deep wells, meteorology and fluxes (Bowen Ratio) over the riparian grass and mesquite thicket (Scott et al., this issue), and stream stage at section three were made for all of 1997. Measurements were made of gravimetric soil moisture, temperature, and soil tension measurements in the near-stream bank trenches to investigate bank storage-stream conductance relationships (Whitaker et al., this issue) for the last half of 1997.

In addition to the continuous measurements, intensive synoptic *in-situ* and remote measurement campaigns of 32 to 56 hours were conducted in March, April, June, August, and October. This period spans the pre-green up and growing season to allow for characterization of the seasonal variations in evapotranspiration, and surface water - groundwater interactions. Measurements taken during the synoptic runs included hourly stream stage and water levels from five river sections and the piezometer network, stream discharge measurements determined by current metering, dye-dilution, and an in-stream flume for the June campaign (see summary by Maddock et al, this issue). Neutron probe measurements of soil moisture were made during synoptic runs and other periods of dynamic change. Treesap flow, water potential, stomatal conductance, and water sources using stable oxygen and hydrogen isotope ratios were determined for mesquite, cottonwood, and willow during each of the synoptic campaigns to capture variations in transpiration demand as a function of atmospheric demand and surface moisture availability. A subset of the vegetation and water source measurements was also made in a both an intermittent losing and ephemeral riparian stream reach (at Boquillas and Escapule locations, respectively - see Figure 1). This design enabled the proportion and magnitude of surface water use by the gallery trees as a function groundwater availability to be evaluated (see overview by Williams et al, this issue).

Concurrent with these *in-situ* measurements, ground-, aircraft-, and satellite-based remotely-sensed measurements were taken. Plant and soil reflectance, temperature and radar backscatter remote measurements were made during each synoptic run and at other strategic times during 1997. These data will be used to develop remote-sensing approaches for monitoring regional and riparian corridor vegetation productivity and evapotranspiration (see remote sensing summary in Moran et al, this issue).

During the August campaign a greater concentration of instrumentation, including two additional aircraft, were deployed at the Lewis Springs site from Aug. 8-19. Additional instrumentation deployed at this time included an array of eddy correlation flux instrumentation (see ET summary by Hipps et al., this issue). A scintillometer, and the Las Alamos National Laboratory Raman LIDAR system (Cooper et al., this issue). During the first portion of the August campaign an intercomparison of flux instruments was conducted over a homogeneous region of the sacaton grass while the LIDAR was deployed on the east bank of the San Pedro River. After several days the LIDAR was moved to the west bank and the flux instruments were redeployed across the highly heterogeneous vegetation types under the path of the scintillometer. *Objective four* will be addressed with this data due to the wide variation in vegetation type, height and stress resulting from differential access to groundwater in the compact riparian area.

Finally, ground-based remotely sensed observations of atmospheric moisture derived from interferometric methods (Shao et al., this volume) were also taken roughly 400 m south of the Lewis Springs site. Like the scintillometer measurements, these were taken on a scale



measurements to estimate the evaporative water loss over the USPB riparian corridor. These evapotranspiration losses had a significant impact on both stream discharge and near stream groundwater levels (MacNish et al., this issue). Prior to green-up and inputs from storm flows, discharge at both Lewis Springs and the Charleston gage were very stable. After green-up of the cottonwoods and willows, stream discharge decreased and a distinct diurnal pattern in stream discharge was apparent.

Initial results also indicate that diurnal and seasonal variations in stomatal conductance and transpiration in cottonwood and willow were strongly influenced by radiation inputs and vapor pressure deficit (VPD). Transpiration fluxes at Lewis Springs were about 12% lower during the June observation period compared to that in August, which was proportional to differences in VPD. Water potential gradients between the rhizosphere and leaf during maximum transpiration, however, were higher in June than in August, suggesting that hydraulic conductivity of trees may have increased over the growing season. Stable isotope analysis revealed that cottonwood along the river at the perennial Lewis Springs site used only groundwater and did not take up surface water even following a large rain and flood event in August. Mesquite at this site likewise used mostly groundwater over the growing season, but shifted to predominantly surface water usage at an intermittent section of the stream (Boquillas site, Figure 1) following the monsoon rains (Williams et al., this issue). Further from the river (100-300 m) at Lewis, initial findings indicate that mesquite and sacaton did not use groundwater (Scott et al., this issue). These data, taken together, illustrate the complex nature of plant controls on ecosystem water fluxes within desert riparian ecosystems.

Initial results related to *objective four* indicate that sensible heat flux estimates from the 3-D sonics and scintillometer compare well over uniform terrain in both the U.S. and Mexico. Good comparisons were also obtained between the scintillometer and a weighted average sensible heat fluxes derived from the 3-D sonics over the mesquite-grass transect at Lewis Springs. In this case, this implies that a simple aggregation rule can provide good results (Chehbouni et al., this issue).

Results related to *objective five* indicated a 35% decrease in the grasslands, an 11% increase in desert shrubs, and a 50% increase in woodlands over a large area of the USPB in the 13-year span between 1974 and 1987. The grasslands also became much more fragmented with the number of grassland patches increasing 61% while the average patch size of the grasslands decreased 60%. These changes will have different implications for habitat suitability for various species due to differences in life history and scale requirements (i.e. patch size). However, for large grazing mammals such as the Mexican pronghorn the changes appear to be quite severe with an estimated 70% reduction in suitable habitat (Kepner, et al., 1995).

In conclusion it should also be noted that the SALSA Program activities have broken new ground in the approach to large-scale multidisciplinary science where limited resources were available. Careful planning resulted in the identification of critical scientific

challenges, of such complexity, as to require the joining of disciplinary forces to address them. Careful attention to fostering interdisciplinary communication built the foundation for trustful collaborations. This enabled unselfish sharing of numerous small grants and in-kind resources to accomplish a goal which is much greater than the sum of the disciplinary parts. An additional driving force behind the SALSA Program's success is the knowledge that the results of this research will directly aid land managers and decision makers in the near term.

## 6. ACKNOWLEDGMENTS

Financial support from the USDA-ARS Global Change Research Program, NASA grant W-18,997, NASA Landsat Science Team, grant #S-41396-F, USDA National Research Initiative Grant Program, Electrical Power Research Institute, Arizona Department of Water Resources, U.S. Environmental Protection Agency-Office of Research and Development, CONACYT, ORSTOM, the French Remote Sensing Program (PNTS) via the VEGETATION Project and the ERS2/ATSR2 Project, and US Bureau of Land Management is gratefully acknowledged. Assistance was also provided in part by the NASA/EOS grant NAGW2425, EPA STAR Graduate Student Fellowship Program, National Science Foundation, US Geological Survey, US Department of Energy contract W-7405-ENG-36, California Institute of Technology-Jet Propulsion Laboratory (NASA, EOS/ASTER), WAU (Wag. Agricultural University, Netherlands), Cochise County Highway and Flood Control Dept., and Ft. Huachuca; this support is also gratefully acknowledged. Special thanks are extended to the ARS staff located in Tombstone, Arizona for their diligent efforts and to USDA-ARS Weslaco for pilot and aircraft support. We also wish to extend our sincere thanks to the many ARS and University of Arizona staff and students, and local volunteers who generously donated their time and expertise to make this project a success.

## 7. REFERENCES

- Goodrich, D. C., 1994: SALSA-MEX: A Large Scale Semi-Arid Land-Surface-Atmospheric Mountain Experiment. Proc. 1994 Intern. Geoscience and Remote Sensing Sym. (IGARSS'94), Pasadena, CA, Vol. 1, Aug. 8-12, p. 190-193.
- Kepner, W., K. Riitters, and J. Wickham. 1995: A landscape approach to monitoring and assessing ecological condition of rangelands-San Pedro River case study. Proc. Symp. EMAP Science, Chapel Hill, NC, March 7-9, 1995. U.S.-EPA, Environ. Monitoring and Assessment Prog., Research Triangle Park, NC.
- Renard, K. G., Lane, L. J., Simanton, J. R., Emmerich, W. E., Stone, J. J., Weltz, M. A., Goodrich, D. C., and Yakowitz, D. S., 1993: Agricultural impacts in an arid environment: Walnut Gulch case study. *Hydrological Science and Technology*, 9(1-4):145-190.
- Wallace, J., 1995: Multidisciplinary Program Studies Land-Atmosphere Interactions in Semi-Arid Regions, *EOS, Transactions, American Geophysical Union*, 76(46), 465,469.
- Woolhiser, D.A., Keefer, T.O., and Redmond, K.T., 1993: Southern oscillation effects on daily precipitation in

the southwestern United States. *Water Resources Research*, 29(4):1287-1295.

### 7.1 References in this Issue

- Arias, H.M., C. Watts, I. Parra, T. Maddock III, and R. MacNish, SALSA: An international integrated effort to aid in addressing socio-economic issues in the Upper San Pedro River Basin.
- Bégué, A., D. Luquet, J. Dauzat, Y. Nouvellon, and H. Rey, Vegetation hemispherical fluxes calculated from 3d computerized vegetation canopies and ray tracing technique: application to a semi-arid grassland.
- Chehbouni, A., O. Hartogensis, L. Hipps, J-P. Brunel, C. Watts, J. Rodriguez, G. Boulet, G. Dedieu, Y. Kerr, and H. De Bruin, (a) Sensible heat flux measurements using a large aperture scintillometer over heterogeneous surfaces.
- Cooper, D.I., W. Eichinger and L. Hipps, Spatial properties of water vapor scalar and flux over a riparian corridor.
- Harlow, C., J. Toth, and J. Shuttleworth Evaluation of the SIB2 surface scheme in a semi-arid region.
- Hipps, L.E., D. Cooper, W. Eichinger, D. Williams, S. Schaeffer, K. Snyder, R. Scott, A. Chehbouni, C. Watts, O. Hartogensis, J-P. Lhomme, B. Monteny, J-P. Brunel, G. Boulet, J. Schieldge, H. De Bruin, J. Shuttleworth, and Y. Kerr, A summary of processes which are connected to evaporation of riparian and heterogeneous upland vegetation in arid regions.
- MacNish, R., C. Peters, M. Schulte, D. Goodrich, D. Pool, T. Maddock, III, C. Unkrich, M. Whitaker, B. Goff, Quantification of groundwater - surface water interactions in a southwestern riparian system.
- Maddock, T., III, R. MacNish, D. Goodrich, D. Williams, J. Shuttleworth, B. Goff, R. Scott, S. Moran, D. Cooper, L. Hipps, A. Chehbouni, An overview of atmospheric and surface water coupling to regional groundwater models in semi-arid basins.
- Moran, S., D. Williams, D. Goodrich, A. Chehbouni, A. Bégué, G. Boulet, D. Cooper, R. Davis, G. Dedieu, W. Eichinger, J. Everitt, B. Goff, C. Harlow, M. Helfert, L. Hipps, C. Holifield, D. Hymer, A. Kahle, T. Keefer, Y. Kerr, R. Marsett, W. Ni, Y. Nouvellon, J. Qi, S. Schaeffer, J. Schieldge, R. Scott, K. Snyder, J. Toth, C. Watts, M. Whitaker, and I. Yucel, Overview of remote sensing of semi-arid ecosystem function in the upper San Pedro river basin, Arizona.
- Moran, S. (b), D. Hymer, J. Qi, E. Sano, R. Marsett, M. Helfert, and E. Sano, Soil moisture evaluation using Synthetic Aperture Radar (SAR) and optical remote sensing in semi-arid rangeland.
- Nouvellon, Y., S. Rambal, A. Bégué, S. Moran, J-P. Lhomme, J. Qi, A. Chehbouni, D. Lo Seen, An ecophysiological model for semi-arid grasslands.
- Scott, R., J. Shuttleworth and D. Goodrich, Water use of two dominant riparian vegetation communities in southeastern Arizona.
- Shao, X. M., R.C. Carlos, M.W. Kirkland, Observations of precipitable water vapor in the planetary boundary layer via microwave interferometry.
- Toth, J.J., D.C. Goodrich, M.S. Moran, R. Scott, D. Hymer, J. Shuttleworth, I. Yucel, J. Garatuza-Payan, C. Harlow, A. Chehbouni, G. Boulet, D. Cooper, M. Kirkland, C.J. Watts, R. Avissar, R. Pinker, and L. Hipps, An overview of spatially distributed surface fluxes of heat and moisture across the semi-arid San Pedro basin
- Whitaker, M.P.L., T. Maddock, B. MacNish, D. Goodrich and B. Goff, Monitoring bank storage in the San Pedro riparian national conservation area, Arizona.
- Williams, D.G., J-P. Brunel, G. Lin, S. Schaeffer, and K. Snyder, Biotic control over the functioning of desert riparian ecosystems.
- Yucel, I., J. Garatuza-Payan, J. Toth, J. Shuttleworth, and R. Pinker, Cloud screened surface radiation estimates with high resolution from GOES-9 data.




Figure 2: Lewis Springs Instrumentation and Measurement Locations





