

1020

MÉTHODE PROVISOIRE POUR DÉPISTER LA RÉSISTANCE À L'ENDOSULFAN CHEZ LE SCOLYTE DE LA CERISE DU CAFÉ, *HYPOTHENEMUS HAMPEI* (COLEOPTERA: SCOLYTIDAE)

Résumé

Cet article décrit et compare trois méthodes de mesure de la résistance à l'endosulfan du scolyte du café, *Hypothenemus hampei*: la pulvérisation directe des femelles avec la tour de Potter et l'exposition directe ou indirecte à un papier filtre préalablement traité. La similarité des résultats obtenus suggère que la tour de Potter sera utilisée pour toutes les études toxicologiques et que la méthode d'exposition indirecte aux insecticides conviendra aux programmes visant à détecter précocement de nouveaux cas de résistance ou à gérer une résistance déjà établie.

METODO PROVISIONAL PARA DETECTAR LA RESISTENCIA AL ENDOSULFAN EN LA BROCA DEL CAFE, *HYPOTHENEMUS HAMPEI* (COLEOPTERA: SCOLYTIDAE)

Resumen

En el presente artículo se comparan tres métodos para determinar la resistencia de *Hypothenemus hampei* al endosulfán: la técnica de pulverización directa, ya descrita en otra publicación, y los métodos de exposición directa e indirecta utilizando papel de filtro tratado. Los resultados indican que la eficacia es parecida con los tres métodos. El método indirecto, que utiliza la acción del vapor mediante la exposición de los insectos en un espacio cerrado por encima del papel de filtro tratado, fue el más fácil de aplicar. Se recomienda este método como sistema normal provisional para la detección de resistencia al endosulfán en esta especie. En un estudio comparativo entre la pulverización directa y la exposición indirecta sólo se observó una diferencia en el diagnóstico de la resistencia en 162 cepas estudiadas.

2

## Provisional method for detecting endosulfan resistance in coffee berry borer, *Hypothenemus hampei* (Coleoptera: Scolytidae)

L.O. BRUN, V. GAUDICHON  
and C. MARCILLAUD

**Summary.** This paper compares three methods for assessment of endosulfan resistance in *Hypothenemus hampei*: the direct spray technique reported previously, and direct and indirect exposure methods using treated filter paper. The results indicate comparable accuracy with all three methods. The indirect method, using vapour action by exposing insects in a confined space above the treated filter, was the easiest to use. This method is recommended as the provisional standard method for detection of endosulfan resistance in this species. A comparative survey of the direct spray and indirect exposure methods showed only one difference in diagnosis of resistance out of 162 strains tested.

Coffee berry borer (CBB), *Hypothenemus hampei* (Ferrari), was reported for the first time in 1948 from the northern part of New Caledonia (Bugnicourt, 1950). *H. hampei* spread rapidly and became the major pest of coffee, which is predominantly composed of *Coffea caniphora* var. *robusta*. CBB is controlled throughout the world predominantly with the use of a single cyclodiene insecticide, endosulfan (Huttenbach, 1982). Alternative insecticides are generally less effective and not widely used (Rhodes and Mansingh, 1981).


The discovery of high levels of resistance to endosulfan in New Caledonia may indicate future problems elsewhere, given similar pat-

The authors are with the Institut français de recherche scientifique pour le développement en coopération (ORSTOM), B.P. A5, Nouméa, New Caledonia.

They are grateful for the suggestions and comments of Dr D.M. Suckling (DSIR, New Zealand) and thank Dr D.F. Waterhouse (ACIAR, Australia) for useful comments and discussions on the development of the research programme. They are also indebted to Mr Moulis who constructed the first detection kit. This work was supported in part by the Coffee Board of New Caledonia (*l'Opération Café*).

FAO Plant Prot. Bull.

Fonds Documentaire IRD



010023816

Fonds Documentaire IRD

Cote : B\*23816 Ex : 1



Comparison of dosage-mortality responses of endosulfan-susceptible and resistant strains of *Hypothenemus hampei*

Strain	N <sup>3</sup>	Slope (SE)	LC <sub>50</sub> (95% CL)	LC <sub>90</sub> (95% CL)	LC <sub>99</sub>
Direct spray <sup>1</sup>					
S	1 380	2.87(0.14)	28(24-32)	78(66-97)	180
R	240	0.46(0.75)	*** <sup>2</sup>	*** <sup>2</sup>	*** <sup>2</sup>
Indirect exposure					
S	300	4.67(0.79)	64(38-82)	119(91-266)	200
R	480	0.59(0.14)	26 900 (5.592 - 1 × 10 <sup>6</sup> )	1 × 10 <sup>6</sup> (1.9 × 10 <sup>5</sup> - *** <sup>2</sup> )	*** <sup>2</sup> *** <sup>2</sup>
Residue					
S	90	4.29(0.88)	50(32-66)	99(74-148)	173
R	150	0.51(0.35)	*** <sup>2</sup>	*** <sup>2</sup>	*** <sup>2</sup>

<sup>1</sup> Reported in Brun *et al.*, 1989.

<sup>2</sup> Resistance level too high to calculate parameter by probit analysis.

<sup>3</sup> Number of females tested.

pipetted on filter paper with a rotational motion.

A thin sheet of transparent plastic (75 × 135 mm) with ten holes (15 mm diameter) was used to cover partly the testing arenas to prevent escape. Any test with more than 10 percent mortality was discarded, but control mortality was generally lower than 5 percent. Mortality was assessed after 6 h exposure at 25°C and 80-85 percent R.H.

**Indirect exposure.** The technique used was essentially the same as the residue exposure method, but a fine white nylon gauze was placed above the filter paper prior to adding the beetles to each hole. In this way the CBB were only exposed to the vapour of endosulfan. The test was repeated twice with each sample and a water-only treatment was included for each batch of tests. This procedure ensured that the beetles did not eat the filter paper, which occasionally occurred with the residue exposure method. Exposure time and mortality assessment were the same as above. The diagnostic dosage used was 400 ppm of endosulfan.

**Statistical analysis.** Probit analysis was done with POLO (Robertson, Russell and Savin, 1980) to determine the LC<sub>99</sub> of susceptible insects with each technique. The diagnostic dosages were set at approximately twice the LC<sub>99</sub> of susceptibles. Sites with any beetles surviving the diagnostic dosage were considered to include resistant individuals.

## RESULTS AND DISCUSSION

**Comparison of methods.** A very close similarity of endosulfan toxicity to susceptible insects was evident with the three methods which overlap in confidence limits between "residue" and "indirect exposure" methods (see Table). This similarity in responses of susceptible CBB between techniques (for slopes and LC<sub>50</sub> S, see Table) led us to use the same endosulfan concentration (400 ppm) as a diagnostic dosage for the survey using "direct" spray and indirect exposure methods.

The high level of resistance to endosulfan determined from the Potter tower direct spray technique has been reported previously (Brun

*et al.*, 1989). The resistance factor ( $LC_{50} R / LC_{50} S$ ) was estimated at over one thousand fold by graphical analysis for direct spray and residue tests, while the resistance factor was 572 fold (see Table).

The authors have assessed mortality of resistant strains at 400 ppm for longer periods of time (up to 96 h). No substantial increase in mortality of resistant females was observed between 6 and 48 h, so a period of 6 h was chosen as recommended by FAO (FAO, 1980). The increase in mortality of treated insects after 48 h was at the same rate as in untreated control insects. The advantage of this short time period is that results are rapid.

The main limitation of the Potter tower direct spray technique is the need for specialized spray application equipment. The residue exposure technique was developed with the aim of reproducibility, low cost, and ease of use. However, an occasional problem occurring with the residue exposure technique was beetles eating or shredding the filter paper. To eliminate this possibility, the beetles were caged above the treated surface using fine gauze. We found that it was easier to get clear results with this indirect exposure method. Mortality assessment was also made easier, since the beetles could not become entangled in the paper because of the behaviour of the scolytid, which is inclined to bore holes in berries. Beetle mortality in this type of bioassay relies entirely on the high vapour action of endosulfan (Knauf, 1982).

The close similarity of endosulfan toxicity evident with the three methods, suggests that toxicity from exposure to the "gas phase" may be of primary importance in all three testing methods.


**Survey results.** The survey included 162 collection sites throughout New Caledonia covering the West Coast, East Coast, Northern Region, Middle Region and Southern Region, but resistance was detected only on the East Coast. The two detection methods showed excellent agreement, with a difference in detection of resistance at only one site. Resistance was present at 43.2 percent of the sites according to

the direct spray method, and at 42.6 percent according to the indirect exposure method.

Further details of the distribution of resistance in New Caledonia are reported by Brun *et al.* (in preparation).

## CONCLUSIONS

The indirect exposure technique offers reproducibility, low cost, and ease of use. It is recommended for future surveys of endosulfan resistance in the coffee berry borer. The efficiency of the technique permits extensive screening of strains and should allow early detection of resistance problems. Kits for the indirect exposure method (see Figure) are available from the authors, ORSTOM, Nouméa, New Caledonia.



*Components of the test method, showing the different layers of Perspex, filter paper and gauze. The layers are assembled as shown*

## REFERENCES

- ABBOTT, W.S. 1925. A method of computing the effectiveness of an insecticide. *J. Econ. Entomol.*, 18: 265-267.
- BRUN, L.O., MARCILLAUD, C., GAUDICHON, V. & SUCKLING, D.M. 1989. Endosulfan resistance in coffee berry borer, *Hypothenemus hampei* (Coleoptera: Scolytidae) in New Caledonia. *J. Econ. Entomol.*, 87 (5): 1311-1316.
- BRUN, L.O., MARCILLAUD, C., GAUDICHON, V. & SUCKLING, D.M. *Monitoring endosulfan and lindane resistance in the coffee berry borer: Hypothenemus hampei* (Ferrari) (Coleoptera: Scolytidae) in New Caledonia. (In preparation)
- BRUN, L.O., MARCILLAUD, C., GAUDICHON, V. & SUCKLING, D.M. *Influence of temperature on the indirect exposure method for detecting endosulfan resistance in coffee berry borer.* (In preparation)
- BUGNICOURT, F. 1950. Le scolyte du grain de café en Nouvelle-Calédonie. *Rev. Agric. Nouv.-Calédonie*, 1-2: 3-4.
- FAO. 1980. Recommended methods for measurements of pest resistance to pesticides. Method for adults of some major pests of stored cereals with malathion or lindane. FAO method no. 15. *FAO Plant Production and Protection Paper*, 21: 77-89.
- HUTTENBACH, H. 1982. Fields of application and survey of technical literature. *Residue Reviews*, 83: 112-137.
- KNAUF, W. 1982. Effect of endosulfan and its metabolites on arthropods. *Residue Reviews*, 83: 89-112.
- POTTER, C. 1952. An improved, laboratory apparatus for applied direct sprays and surface film, with data on the electrostatic charge on atomized spray fluids. *Ann. Appl. Biol.*, 39: 1-28.
- RHODES, L.F. & MANSINGH, A. 1981. Susceptibility of the coffee berry borer *Hypothenemus hampei* (Ferrari) to various insecticidal formulations. *Insect Sci. Appl.*, 2: 227-231.
- ROBERTSON, J.L., RUSSELL, R.M. & SAVIN, N.A. 1980. *POLO: A user guide to probit or logit analysis*. USDA Forest Service General Technical Report, PSW-38.

# OUTBREAKS AND NEW RECORDS ATTAQUES ET NOUVEAUX ENNEMIS SIGNALÉS NUEVOS FOCOS DE ENFERMEDADES Y PLAGAS

## BHUTAN

W.T.H. PEREGRINE, FAO Office (Project UGA/87/006), Kampala, Uganda, & THINLAY, EEC Plant Protection Project, Ministry of Agriculture, Box 119, Thimphu, Bhutan.

### **Black leaf streak of banana *Mycosphaerella fijiensis* Morelet**

Black leaf streak of banana, *Mycosphaerella fijiensis* Morelet, was recognized in 1963 as a distinct and more severe disease than yellow Sigatoka, *M. musicola* Leach ex Mulder.<sup>1</sup>

The authors now report the presence of black leaf streak in the Himalayan kingdom of Bhutan, with the fungal identity confirmed in 1985 by the CAB International Mycological Institute as *Paracercospora fijiensis* (Morelet) Deighton state, (*Mycosphaerella fijiensis* Morelet), IMI No. 290376.

Since bananas are only a backyard crop in Bhutan, the disease is not of economic significance there. Yellow Sigatoka is still more common. However, the route taken by black leaf streak into the country is of considerable interest.

As the disease is not known to occur in the Indian subcontinent, one possible route is from Malaysia or Thailand via Myanmar, with which country Bhutan shares an eastern border. The authors' record, however, was made in southwest Bhutan and in an area where cross-border trade and exchange of planting materials with India is normal. It is possible therefore that the disease may already be in India or another country.

Its advent in Bhutan reinforces the importance of an effective plant quarantine system, especially in countries with an agriculture-based economy. The situation should be reviewed in the light of the explosive spread of some other diseases, such as groundnut rust (*Puccinia arachidis*) throughout Asia and Africa, following its discovery in Brunei in 1970.<sup>2</sup>

<sup>1</sup> MEREDITH, D.S. & LAWRENCE, J.S. 1969. Black leaf streak disease of banana (*Mycosphaerella fijiensis*): symptoms of disease in Hawaii and notes on the conidial state of the casual fungus. *Transactions of the British Mycological Society*, 52: 459-476.

<sup>2</sup> PEREGRINE, W.T.H. 1971. Groundnut rust (*Puccinia arachidis*) in Brunei. *PANS*, 17: 3.

## CANADA

Reported to FAO by D. Coates-Milne, Director, Plant Protection Division, Canada.

### **Eradication of potato spindle tuber viroid (PSTV) in the Canadian provinces of Prince Edward Island and New Brunswick**

Scientific surveys conducted from 1984 to 1987 in New Brunswick and Prince Edward Island, Canada, have confirmed the absence of potato

spindle tuber viroid in seed and ware potato fields. Leaf samples taken during the survey were tested by means of polyacrylamide gel electrophoresis (PAGE), nucleic acid hybridization (dot-blot) and return-polyacrylamide gel electrophoresis (R-PAGE). These laboratory results corroborate field inspection and post-harvest testing data collected by Canada's Plant Protection Division over the past seven years, during which no PSTV was detected.

The eradication of this infectious pathogen was accomplished through rigorous screening