

NOTE

TOXICITY OF FOUR INSECTICIDES TO *PHYTOSEIULUS MACROPILIS* (BANKS) AND *P. PERSIMILIS* ATHIAS-HENRIOT (ACARINA: PHYTOSEIIDAE)L. O. BRUN¹, J. CHAZEAU¹, and V. E. EDGE²¹ ORSTOM Centre de Noumea, Boite Postale A5, Noumea-Cedex, New Caledonia² Biological and Chemical Research Institute, Department of Agriculture N.S.W., P.M.B. 10, Rydalmere, N.S.W. 2116.

Abstract

Commercial formulations of dimethoate, methomyl, fenvalerate and endosulfan were tested against a New Caledonian strain of *Phytoseiulus macropilis* and an Australian strain *P. persimilis* using a slide-dip technique. The responses of the 2 species to fenvalerate, methomyl and endosulfan were similar, with fenvalerate being the most toxic and endosulfan the least. In contrast, dimethoate was 50x more toxic to *P. macropilis* than to *P. persimilis* at the LC₅₀ level.

Phytoseiulus persimilis Athias-Henriot has been widely used for the control of tetranychid mites and its response to pesticides has been evaluated (Everson and Tonks 1981). Resistance to several organophosphorus compounds has been confirmed in *P. persimilis* (Schulten *et al.* 1976). *P. macropilis* Banks is an effective predator of tetranychid mites (Hamlen and Poole 1980), but there are no data on its response to pesticides.

In the present investigation commercial formulations of 4 insecticides from different chemical groups were tested against a New Caledonian strain of *P. macropilis* and a strain of *P. persimilis* from Sydney.

The test strain of *P. macropilis* was established with several hundred individuals collected from French beans adjacent to a commercial nursery near Noumea, New Caledonia. The culture was maintained at 25°C in continuous illumination on potted French bean plants infested with *Tetranychus urticae* Koch, twospotted mite. A slide-dip technique (Anon. 1974) was used to test the following materials: dimethoate (30% w/v, Rogor®), methomyl (20% w/v, Lannate®), fenvalerate (10% w/v, Sumicidin 10®), and endosulfan (35% w/v, Thiodan®). Twenty gravid females were attached to each slide using double sided adhesive tape and dipped in an aqueous suspension of the insecticide for 5 seconds. Four concentrations of each insecticide were used per test and the tests were replicated 5 times. A water only treatment was included in each replicate as a control. After dipping, excess liquid was removed from around each female with absorbent paper and slides were held at 27 ± 1°C and ca 95% RH. Mortality was assessed 48 h after treatment; the criterion for death was failure to exhibit leg movement when gently prodded with a fine bristle. The data were corrected for control mortality, which did not exceed 10%, and analysed by probit analysis (Finney 1971).

The same samples of the insecticide formulations were then tested in Australia against a Sydney strain of *P. persimilis* (Goodwin and Schicha 1979). The rearing conditions and the test method were the same as those described for *P. macropilis*, but only one concentration of each insecticide (that giving 90-99% mortality of *P. macropilis*) was tested against *P. persimilis* using 20 females per test and replicated 5 times. Dose response data were obtained for dimethoate against *P. persimilis* as described for *P. macropilis* after the initial dosage of 50 ppm gave <30% mortality.

The log-dose probability data obtained for *P. macropilis* are shown in Table 1. These data were characterised by low slope values and the χ^2 values for residual between doses (Table 1) indicate significant departure from linearity for dimethoate, methomyl and endosulfan. Low slope values were also reported by Schulten *et al.* 1976) after testing a range of insecticides against a susceptible strain of *P. persimilis*. Both sets of data may indicate that there is considerable inherent variation in the response of *Phytoseiulus* sp. to pesticides, particularly in unselected strains. This problem may be compounded by the test method, although it is widely used for predatory mites (Croft 1977).

The single dosage tests (Table 2) indicated that the relative toxicities of methomyl, fenvalerate and endosulfan were similar for *P. macropilis* and *P. persimilis*. The low toxicity of endosulfan is consistent with

TABLE 1
TOXICITY OF 4 INSECTICIDES TO *P. MACROPILIS* FEMALES

	LC50 ppm (a.i.) - (95% fiducial limits)	Slope value ± SE	χ^2
Dimethoate	7 (3-13)	1.9 ± 0.46	9.7*
Methomyl	39 (27-58)	1.5 ± 0.2	10.5*
Fenvalerate	1 (0.5-1.9)	1.1 ± 0.18	5.1
Endosulfan	1100 (500-2100)	1.8 ± 0.39	9.5*

* Significant departure from linearity (P < 0.01)



Fonds Documentaire IRD
Cote : B* 24 233 Ex : 1

TABLE 2
RESPONSES OF *P. MACROPILIS* AND *P. PERSIMILIS* FEMALES TO 4 INSECTICIDES

Concentration ppm (a.i.)	% mortality	
	<i>P. macronilis</i>	<i>P. persimilis</i>

- COCHEREAU, P. (1976).—Contrôle biologique, en Nouvelle Calédonie de *Tetranychus urticae* (Acarien: *Tetranychidae*) au moyen de *Phytoseiulus persimilis* (Acarien: *Phytoseiidae*), en cultures maraichères. *Entomophaga* 21: 151-156.
- COLLYER, E. (1980).—Phytoseiidae (Acari) from the Pacific Islands: note. *N.Z. Entomol.* 7: 138-139
- CROFT, B. A. (1977).—Resistance in arthropod predators and parasites. In Watson, D. L. and Brown, A. W. A. (Eds.). *Pesticide management and insecticide resistance*. Academic Press: New York.
- EVERSON, P. R. and TONKS, N. V. (1981).—The effect of temperature on the toxicity of several pesticides to *Phytoseiulus persimilis* (Acarina: *Phytoseiidae*) and *Tetranychus urticae* (Acarina: *Tetranychidae*). *Can. Ent.* 113: 333-336.
- FINNEY, D. J. (1971).—*Probit Analysis*. 3rd edition. Cambridge University Press: London.
- GOODWIN, S. and SCHICHA, E. (1979).—Discovery of the predatory mite *Phytoseiulus persimilis* Athias-Henriot (Acarina: *Phytoseiidae*) in Australia. *J. Aust. ent. Soc.* 18: 304.
- HAMLEN, R. A. and POOLE, R. T. (1980).—Effects of a predaceous mite on spider mite populations of *Dieffenbachia* under greenhouse and interior environments. *Hort. Science* 15: 611-612.
- HINKLEY, A. D. (1963).—Trophic records of some insects, mites and ticks in Fiji. *Bull. Dept. Agric. Fiji* 45, 116 pp.
- PRASAD, V. (1967).—Biology of the predatory mite *Phytoseiulus macropilis* in Hawaii (Acarina: *Phytoseiidae*). *J. econ. Ent.* 60: 905-908.
- SABA, F. (1974).—Life history and population dynamics of *Tetranychus tumidus* in Florida (Acarina: *Tetranychidae*). *Fla Ent.* 57: 47-63.
- SCHULTEN, G. G. M., VAN DE KLASHORST, G. and RUSSELL, V. M. (1976).—Resistance of *Phytoseiulus persimilis* A. H. (Acari: *Phytoseiidae*) to some insecticides. *Z. ang. Ent.* 80: 337-341.
- STRICKLER, K. and CROFT, B. A. (1982).—Selection for permethrin resistance in the predatory mite *Amblyseius fallacis*. *Ent. exp. & appl.* 31: 339-345.

[Manuscript received 31 August 1982, Revised 20 January 1983]