

Nickel Content of Deep-Sea Deposits

INVESTIGATIONS made in this Institute of the ferrides present in sediment cores from great depths, taken by the Swedish Deep-Sea Expedition, have shown their content of nickel to be specially interesting. The only analysis of nickel in deep-sea deposits known to us was published in 1914 by F. W. Clarke and G. Steiger, and gave 0.032 per cent as the content of nickel oxide in a composite sample of red clay from fifty-one different localities¹. By the aid of modern microanalytical methods (the colour reaction with dimethyl glyoxime) quantities of nickel as small as 5-6 γ can easily be measured.

In a core of red clay from the north-west Atlantic Ocean (lat. 30° 29' N., long. 56° 07' W., depth 5,620 metres, length of core 1,530 cm.) the content of nickel oxide was found to vary between 0.005 and 0.010 per cent, whereas in a red clay core from the central Pacific Ocean (lat. 07° 38' S., long. 152° 53' W., depth 5,000 metres, length of core 1,500 cm.) much higher values were found, ranging between 0.041 and 0.089 per cent of nickel oxide. Assuming the decrease in radium content from the surface layer downwards found in the latter core to be due to the disintegration of ionium, precipitated from the sea-water, namely, a decrease to 50 per cent in 83,000 years, the average rate of deposition in the uppermost parts of the same core would be of the order 0.5 mm. in one thousand years, whereas in the Atlantic core the rate of deposition appears to have been at least ten times higher. The content of nickel oxide in the two cores, therefore, appears to be roughly in inverse proportion to the rate of sedimentation. As is seen from the accompanying graph, where the content of iron-, manganese- and nickel-oxide in the Pacific core is set out as a function of the distance from the top of the core, there are five distinct maxima of nickel content, which do not seem to be related to those of the two other elements.

In another Pacific core taken about 400 nautical miles farther south and in a smaller depth, namely, 4,625 metres, there was also a pronounced maximum of nickel oxide in a 'depth' of 223 cm., namely, 0.098 per cent, which possibly corresponds to the uppermost maximum found in the more northerly core.

Fonds Documentaire IRD




010025824

Fonds Documentaire IRD

Cote : Bx25824 Ex : *un*

n° 481



Vertical axis shows percentage of Fe_2O_3 and MnO_2 (left), and NiO (right); horizontal axis, depth in metres.
 —, MnO_2 ; ---, NiO ; ·····, Fe_2O_3 .

Regarding the origin of the nickel found by us in the red clay, three different explanations appear possible :

(1) It may have originated from sea water. This, however, appears improbable, considering the extremely low nickel content in the water (about 10^{-8} per cent or about 500 times less than its content of iron).

(2) It may have originated from magma spread over the ocean floor by submarine eruptions. Thanks to the courtesy of Dr. S. Landermon of the Geo.

(3) That the nickel may have originated from meteors and from meteoritic dust, on the other hand, seems worthy of serious consideration. Murray and Renard, in their classical studies of the deposits taken by the *Challenger* Expedition, recognized a small number of "cosmic spherules" containing nickel-iron in the red clay and other similar deep-sea deposits. It is reasonable to expect that other fragments from shooting stars in a metallic or oxidized state, as well as minor fragments of interplanetary dust, drifting into the earth's atmosphere, must be present in the deposits to an extent inversely proportional to the rate of sedimentation. A highly approximate calculation, from what we know at present regarding the nickel content and the rate of sedimentation of the cores we have so far examined, leads to unexpectedly high values for the rate of accretion of meteoric material over the whole earth. Our estimates indicate several thousand tons per day (assuming an average nickel content of 2 per cent by weight), which is considerably more than astronomers