



Pergamon

Tetrahedron Letters, Vol. 36, No. 48, pp. 8853-8856, 1995
Elsevier Science Ltd
Printed in Great Britain
0040-4039/95 \$9.50+0.00

0040-4039(95)01837-9

Caledonin, a Natural Peptide Bolaphile with Zn^{II} and Cu^I Complexing Properties from the Tunicate *Didemnum rodriguezii*

María Jesús Vázquez¹, Emilio Quiñoa¹ and Ricardo Riguera^{1*}
Anuncia Ocampo² and Teresa Iglesias²
Cécile Debitus³


¹Departamento de Química Orgánica, Facultad de Química, Universidad de Santiago,
15706 Santiago de Compostela, Spain, Tel/FAX 34-81-391091

²Facultad de Ciencias, Universidad de Vigo, Vigo, Spain

³Centre ORSTOM, B. P. AS, Nouméa Cédex, New Caledonia

Abstract: Caledonin (**1**) is a modified peptide isolated from the marine tunicate *Didemnum rodriguezii*. Caledonin comprises a central L-phenylalanine residue connected via its amino group to (S)-3-aminopropylmercaptopentanoic acid (a new sulfur-containing β-amino acid), and by its carboxyl group to a six membered cycloguanidine moiety, bearing an n-octyl chain. Caledonin is a natural bolaphile which strongly binds Zn^{II} and Cu^I ions.

In the last decades, tunicates have been subjected to very close scrutiny and as a result, a large number of bioactive nitrogenous metabolites with significant biological activity have been described.¹ They include acyclic and cyclic peptides and depsipeptides with novel and complex structures and also compounds of physiological importance characterized by the presence of cyclic guanidine groups.^{2,3} As part of our work on pharmacologically active metabolites from marine organisms, we now report the structure of caledonin (**1**), a novel modified peptide, which was isolated from the tunicate *Didemnum rodriguezii*. Caledonin, among other relevant structural features, possesses a new β-amino acid residue, a cycloguanidine group and presents strong metal-complexing properties.



Didemnum rodriguezii (Tunicata, Didemniidae),⁴ an encrusting ascidian of a deep pink, red or yellow colour, was collected from the Baie des Citrons, Nouméa (New Caledonia). We isolated⁵ pure caledonin (**1**) [48 mg; mp= 170-172°C; [α]_D²⁰ = +24 (MeOH, c=1.45 mg/mL)] from the methylene chloride soluble material that showed *in vitro* cytotoxicity against K562 cells (85% inhibition at 10 μg/mL). The IR spectrum⁶ showed characteristic absorption bands due to amide groups [(KBr) 1656, 1630 cm⁻¹] and the UV indicated the presence of a phenyl chromophore [(MeOH) λ_{max} 234 (ε 2816), 280 (ε 666; sh) nm]. Its (+) FABMS (glycerol) spectrum showed a peak corresponding to the molecular ion at m/z 489 ([M]⁺), confirmed when NaCl was added to the

8853

Fonds Documentaire IRD



010026203

Fonds Documentaire IRD

Cote : Bx 26 203

Ex : Unique

matrix (m/z 511, $[\text{Na}(\text{M}-\text{H})]^+$). Its EIMS showed the $[\text{M}-\text{H}]^+$ ion at m/z 488 and so did the HREIMS (m/z 488.30650, $\Delta = +1.2$ ppm, corresponding to $\text{C}_{26}\text{H}_{42}\text{N}_5\text{O}_2\text{S}$, 8 unsaturations).

One- (^1H , ^{13}C , DEPT) and two-dimensional (DQF-COSY, TOCSY, HMQC) NMR analysis revealed the presence of four main substructures generating independent ^1H spin systems, and linked by one C-N and two peptide bonds. Those fragments were: 1) a new β -amino acid (3-amino-5-mercaptopentanoic acid; 2) phenylalanine; 3) a guanidine group with two nitrogens linked by a $-\text{CH}_2\text{CH}_2\text{CH}_2-$ chain and forming a six membered ring and 4) an *n*-octyl chain. These substructures satisfy the unsaturation requirements (6 double bonds and 2 rings) suggested by the molecular formula.

Diagnostic ^{13}C and ^1H shifts (Table 1) showed the nature of the different heteroatoms and functional groups present in the molecule: one primary amine bound to a methine (δ 61.8, δ 3.24, 3-CH) with γ heteroatom substituents, one guanidine group (δ 160.2, 11-C), two amide groups (δ 171.3, 5-C; δ 166.0, 7-C) and finally, the characteristic shifts of 1- CH_2 (δ 26.9, δ 1.10) indicate a thiol group attached to that position. Exchangeable ^1H NMR spectra ($\text{CDCl}_3/\text{D}_2\text{O}$) confirmed that the broad singlets at δ 4.8 (4H) and δ 6.0 (1H) were originated by hydrogens attached to heteroatoms; and ^1H - ^{15}N HMQC and COSY experiments identified the signal at δ 6.0 as the NH attached to the 10- CH_2 .




Figure 1. a. Selected HMBC ($^2J_{\text{CH}}$ and $^3J_{\text{CH}}$) correlations. b. Selected (+) FABMS fragmentations.

The precise connectivities between the substructures were established by HMBC and corroborated by ROESY experiments. Selected correlations are shown in Figure 1a. Among them, those between 5-CO and 6-CH; 7-CO and 8-CH₂; and 11-C and the CH₂ groups at positions 8, 10 and 12, were the most useful. Caledonin was thus shown to comprise a central phenylalanine residue flanked by a 3-amino-5-mercaptopentanoic acid and the cycloguanidine moiety, with the *n*-octyl chain bound to the exo-nitrogen. Furthermore, the observed (+) FABMS fragmentation patterns are in good agreement with that structure (Figure 1b).

Several chemical transformations were carried out to confirm the structure of caledonin. Acid hydrolysis of **1** (6N HCl; 12h at 110°C) yielded three products identified (^1H -NMR, MS) as phenylalanine, 3-amino-5-mercaptopentanoic acid and the substituted guanidine **2**. Derivatization (1. *n*-BuOH; 110°C. 2. TFAA; 150°C) followed by chiral GC-MS analysis (Chirasil Val III; He at 1 mL/min; 120 to 200°C) indicated that the phenylalanine belonged to the *L*-series (*S* absolute configuration). Derivatization of caledonin with (*R*)- and (*S*)-methoxyphenylacetic acid (MPA), afforded after HPLC separation, the MPA amides **3** and **4**. Analysis of their ^1H -NMR spectra⁷ indicated the absolute configuration at 3-C to be *S* (Figure 2a).⁸

Tunicates are known to produce metal complexing metabolites, such as the tunichroms.⁹ Some features of caldonin, and in particular the hydrophobic chain at one end and the penicillamine-like β -amino acid¹⁰ at the other, suggest that this metabolite is a natural biolipophile¹¹ which may be involved in ion transport across membranes.

To test this hypothesis, a solution of caldonin in CD_3OD was titrated with $ZnCl_2$ and monitored by 1H -NMR spectroscopy. The 2:1 caldonin/ $ZnCl_2$ adduct was isolated and characterized by (+) FABMS (m/z 552, $[Zn(M-H)]^+$) and NMR spectroscopy (1H , ^{13}C , COSY, TOCSY and HMQC experiments).

Table 1. 1H and ^{13}C NMR data for caldonin (1)^a

At. no.	^{13}C (δ)	1H (δ), <i>m</i> , J(Hz)	At. no.	^{13}C (δ)	1H (δ), <i>m</i> , J(Hz)
1	26.9	1.10, dd (7.1, 13.1)	12	40.6	2.68, <i>m</i>
2	31.7	1.55	13	31.0	1.75, dd (7.5, 15.1)
		1.42			
3	61.8	3.24, <i>m</i>	14-17	29.3-29.6	1.38-1.55
4	39.0	3.16, <i>t</i> (12.1)	18	22.7	1.38
		2.12, dd (8.1, 14.1)			
5	171.3		19	14.1	0.89, <i>t</i> (6.6)
6	66.3	4.50, <i>t</i> (1.0)	20	35.6	3.39, dd (4.4, 13.6)
					3.20, dd (2.0, 13.5)
7	166.0		21	134.8	
8	52.4	3.53, <i>d</i> (17.0)	22/26	130.5	7.16-7.26
		2.79, <i>m</i>			
9	27.4	1.55	23/25	128.6	7.16-7.26
10	38.9	3.75, <i>m</i>	24	127.7	7.16-7.26
		2.75, <i>d</i> (17.0)	11-NH		6.00, <i>s</i>
11	160.2		NH(4H)		4.80, <i>bs</i>

^a NMR spectra were recorded on a Bruker AMX-500 spectrometer in $CDCl_3$ at 298 K.

Comparison of the 1H -NMR data of 1 and its Zn^{II} complex (Table 2) showed that the thiol and amino groups are involved in complex formation with the Zn^{II} ion in a tetrahedral environment, as shown in Figure 2b.

Very similar 1H -NMR spectrum was obtained when 1 was titrated with $CuCl$, suggesting formation of a similarly bonded adduct, but the chelated Cu^I ion was rapidly oxidized to Cu^{II} , hampering isolation and precise structure determination.

Table 2. Selected NMR data for caldonin (1) and for caldonin- Zn^{II} complex in CD_3OD .

At. no.	1		1- Zn^{II} complex	
	1H (δ)	^{13}C (δ)	1H (δ)	^{13}C (δ)
1	1.30	27.55	1.18	29.72
2	1.61/1.52	35.75	1.63/1.33	35.51
3	3.24	49.00	4.37	51.85
4	3.16/2.12	40.48	2.41/2.22	41.88




Figure 2. a. Perspective and elongated Newman projections for 3 and 4 showing the magnetic shielding effects based on Ref. 7. b. Caledonin-Zn^{II} complex, showing its bolaphite characteristics.

Acknowledgements

This work was financially supported by the CICYT (SAF 92-1023) and the Xunta de Galicia (XUGA 20907B/93 and XUGA 20907B/94).

REFERENCES AND NOTES

- Ireland, C. M.; Molinski, T. F.; Roll, D. M.; Zabrickie, T. M.; Mokec, T. C.; Swersey, J. C.; Foster, M. P.; in *Bioorganic Marine Chemistry*, vol 3, Scheuer, P.J. Ed., Springer-Verlag, Berlin Heidelberg, 1989, pp. 1-46.
- Ohtani, I.; Knaumi, T.; Kakisawa, H.; Kaetman, Y.; and Hirsh, S. *J. Am. Chem. Soc.*, 1992, 114, 8472.
- Ohtani, I.; and Moore, R.; *J. Am. Chem. Soc.*, 1992, 114, 7941.
- Da Rocha, R.M.; and Monniot, F. *Ann. Inst. Océanogr., Paris*, 1993, 69, 261-265.
- Freshly collected tunicate was frozen on site, transferred over dry ice and then lyophilized. 1 Kg of lyophilizate was homogenized in MeOH and extracted at r.t. for four days (4 x 2 l). The MeOH was evaporated *in vacuo* and the residue was partitioned between water and CH₂Cl₂. 5 g of the CH₂Cl₂-soluble material were successively chromatographed on silicagel (CH₂Cl₂-MeOH) and Sephadex LH-20 (MeOH) columns, and then on reversed-phase HPLC (μ -Bondapak C₁₈ column; MeOH:H₂O 95:5; flow: 4 mL/min; retention time 12 min; RI detector), to afford 48 mg of pure caldonin (1).
- IR (KBr) 2922, 2852, 1656, 1630, 1548, 1457 cm⁻¹.
- Latypov, S. K.; Soco, J. M.; Quiñod, E.; and Riguera, R. *J. Org. Chem.*, 1995, 60, 1538.
- For instance, characteristic shifts for the pair of derivatives 3(4) were δ 4.93(4.65) and 2.21(2.18) (vs 6-H and 4-H respectively).
- a) Sraith, M. J.; Kim, D.; Horenstein, B.; and Nakanishi, K. *Acc. Chem. Res.*, 1991, 24, 117.
b) Michael, J. P.; and Pattenden, G. *Angew. Chem. Int. Ed. Engl.*, 1993, 32, 1.
- Penicillamine is a well known antirheumatic and chelating agent for copper in Wilson's disease: Merck Index, 11th Edition, Merck & Co., Inc. Rahway, N.J., U.S.A., 1989, p. 7029 and references therein. Caledonin also resembles the antihypertensive captopril, Merck Index, 11th Edition, Merck & Co., Inc. Rahway, N.J., U.S.A., 1989, p. 1773 and references therein.
- a) Fuhrhop, J.-H.; and Fritsch, D. *Acc. Chem. Res.*, 1986, 19, 130, and references therein.
b) Escamilla, G. H.; and Newkome, G. R. *Angew. Chem. Int. Ed. Engl.*, 1994, 33, 1937, and references therein.

(Received in UK 13 July 1995; revised 27 September 1995; accepted 29 September 1995)