
**Callipeltin A, an Anti-HIV Cyclic
Depsipeptide from the New Caledonian
Lithistida Sponge *Callipelta* sp.**

**Angela Zampella, M. Valeria D'Auria, Luigi Gomez Paloma,
Agostino Casapullo, Luigi Minale, Cécile Debitus, and Yvette Henin**

Contribution from the Dipartimento di Chimica delle Sostanze Naturali,
Universita' degli Studi di Napoli Federico II, via D. Montesano 49,
80131 Napoli, Italy, ORSTOM, Centre de Nouméa, B.P. A5 Nouméa
Cedex, New Caledonia, and Rhone-Poulenc Rorer Recherche
Developpement, Centre de Recherche de Vitry-Alfortville 13,
Quai Jules Guesde -BP 14, 94403 Vitry Sur Seine Cedex, France

**JOURNAL
OF THE
AMERICAN
CHEMICAL
SOCIETY®**

Reprinted from
Volume 118, Number 26, Pages 6202-6209



Fonds Documentaire IRD

Cote : B * 26 208 Ex : unique


Callipeltin A, an Anti-HIV Cyclic Depsipeptide from the New Caledonian Lithistida Sponge *Callipelta* sp.

Angela Zampella,[§] M. Valeria D'Auria,[§] Luigi Gomez Paloma,[§]
Agostino Casapullo,[§] Luigi Minale,^{*,§} Cécile Debitus,[‡] and Yvette Henin[#]

*Contribution from the Dipartimento di Chimica delle Sostanze Naturali,
Universita' degli Studi di Napoli Federico II, via D. Montesano 49, 80131 Napoli, Italy,
ORSTOM, Centre de Nouméa, B.P. A5 Nouméa Cedex, New Caledonia, and Rhone-Poulenc
Rorer Recherche Developpement, Centre de Recherche de Vitry-Alfortville 13,
Quai Jules Guesde -BP 14, 94403 Vitry Sur Seine Cedex, France*

Received December 28, 1995[®]

Chart 1



(700 g) and extracted sequentially with *n*-hexane, dichloromethane, and a mixture of 8:2 dichloromethane–methanol. Fractionation of the 8:2 dichloromethane–methanol extract by droplet counter current chromatography (DCCC) followed by reversed-phase HPLC yielded callipeltin A (1) as colorless solid (500 mg) ($[\alpha]_D$ in methanol = +3.56).

The FABMS of callipeltin A (1) displayed a protonated molecular ion at m/z 1505 $[M + H]^+$, whereas ESIMS showed a major peak (100%) at m/z 753.5 for $[M + 2H]^{2+}$ accompanied by a minor one (10%) at m/z 1506, corresponding to a molecular weight of 1504 daltons. An intense $[M + 2H]^{2+}$ peak in the ESIMS spectrum indicated the presence of two strong basic groups in the molecule, such as two guanidinium units. The peptidal nature of the compound was indicated by NMR (Table 1) and FTIR spectra (ν_{\max} 1660 and 1740 cm^{-1} , with the latter absorption supporting also the presence of an ester or lactone function). The ^1H NMR spectrum in pyridine- d_5 contained eight amide NH signals between δ 10.1 and 7.27 as implied by COSY and TOCSY (HOHAHA) experiments revealing NH-CH(α) connectivities for eight amino acid residues and two *N*-methyl signals at δ 2.90 and 3.07, totally accounting for the presence of ten amino acids. The signal at δ 7.40 could be assigned to a phenolic OH on the basis of its correlation with the

The aromatic region of the ^1H NMR spectrum (methanol- d_4) contained signals for a *para*-substituted benzene ring at δ 6.81 (d, 2H, $J = 8.5$ Hz) and 7.24 (d, 2H, $J = 8.5$ Hz) and a -CHOMe-CH(α) spin system at δ 3.15 (s, 3H), 4.55 (d, 1H, $J = 9.5$) and 5.01 (d, 1H, $J = 9.5$ Hz). On the basis of ^1H and ^{13}C NMR chemical shifts and COLOC data, the *para*-substituted benzene ring was part of a β -methoxytyrosine residue, which appears to be undescribed. It should be noted that the related β -methoxyphenylalanine was detected in discokialides, peptides isolated from *Discodermia kiiensis*¹⁹ and more recently in cyclomarins, potent antiinflammatory cyclopeptides from marine bacteria.²⁰

The 3,4-dimethylglutamine (or glutamic acid) was difficult to establish by ^1H NMR spectroscopy in methanol- d_4 , because the signals at δ 2.39 ($\text{H}\beta$) and 2.84 ($\text{H}\gamma$) did not give any correlation peak in the COSY spectrum, implying that the dihedral angle between the two hydrogens is close to 90 degrees. However, the COLOC spectrum, which showed correlations between the Me carbon signal at δ 14.6 (Me-C β) and both $\text{H}\beta$ and $\text{H}\gamma$ via $^2J_{\text{CH}}$ and $^3J_{\text{CH}}$, respectively, indicated the proximity of two methines substituted by methyl groups. A correlation peak between the $\text{H}\alpha$ at δ 4.28 and $\text{H}\beta$ at δ 2.39 was observed in the COSY spectrum, whereas correlations between $\text{H}\alpha$ and

Table 1. ^1H and ^{13}C NMR Data^a of Callipeltin A (1) in pyr-*d*₅ and Methanol-*d*₄ at 500 MHz


	pyr- <i>d</i> ₅ ^b		methanol- <i>d</i> ₄		COLOC
	δ_{H}	δ_{C}	δ_{H}	δ_{C}	
			MeAla		
α	5.68 q (6.8) ^c	50.1	5.37 q (7.5)	52.3	NCH ₃ , C β , CO
β	1.13 s	13.7	1.35 d (7.5)	16.3	CO
NCH ₃	2.90 s	30.1	2.86 s	30.1	C α , CO, CO- β MeOTyr
CO				171.5	
			β MeOTyr		
α	5.42 t (9.6)	53.4	5.01 d (9.5)	53.5	CONH
β	4.96 d (9.6)	84.1	4.55 d (9.5)	84.0	CO, C α , C5/C9, OCH ₃
C4		129.2		129.2	
C5/C9	7.64 d (8.0)	131.1	7.24 d (8.5)	130.8	C7
C6/C8	7.25 d (8.0)	116.2	6.81 d (8.5)	115.9	C4
C7		158.5		158.5	
OH	7.40 br s				
OCH ₃	3.15 s	56.9	3.15 s	56.7	C β
CONH	8.72 d (9.6)			171.6	
			MeGln		
α	5.27 br t (7.2)	56.7	4.82 m	56.7	
β	1.82, 1.78 m	26.4	1.65, 1.50 m	26.5	
γ	2.08, 1.93 m	32.4	1.67 m	32.4	
NCH ₃	3.07 s	30.6	3.03 s	30.6	C α , COLeu
CO				172.9 ^e	
CONH ₂ ^d	7.30 overlapped			180.4 ^e	
			Leu		
α	4.92 m	49.7	4.67 dd (10.5, 3.0)	50.1	CONH
β	1.74, 1.50 m	39.6	1.73, 1.33 m	39.8	
γ	1.67 m	25.4	1.70 m	26.3	
δ	0.77 d (6.4)	21.6	0.98 d (6.4)	23.4	C β , C γ
ϵ	0.83 d (6.4)	23.5	0.93 d (6.4)	21.5	C β , C γ , C δ
CONH	7.27 d (8.4)			174.7	
			Arg		
α	4.73 m	52.8	4.36	53.1	
β	2.15, 1.90 m	27.9	2.00, 1.67 m	27.9	
γ	1.90, 1.70 m	26.1	1.67, 1.53 m	26.1	
δ	3.32 m	42.2	3.19 m	41.7	guan
δ NH	8.20 br s				
CONH	8.29 br s			176.1 ^e	
guan	7.92 overlapped			158.3 ^e	
			Thr-1		
α	4.40 d overlapped	63.4	3.96 d (3.7)	63.4	CONH
β	4.56 m	66.8	4.36 overlapped	66.8	
γ	1.45 d (6.0)	21.0	1.33 d (7.4)	20.1	C α , C β
OH	8.65 br d				
CONH	8.54 br s			172.0	
			Thr-2		
α	5.88 (9.2)	56.5	5.48 d (2.7)	56.3	CONH, C β
β	6.07 m	72.2	5.60 m	72.4	
γ	1.52 (6.4)	15.1	1.26 d (6.8)	14.6	C α
CONH	10.1 br s			172.9 ^e	
			DiMeGln		
α	5.03 d overlapped	59.3	4.28 d (9.5)	58.7	CONH, C β
β	2.81 m	38.5	2.39 m	37.7	β Me
β Me	1.60 d (7.2)	13.3	1.26 d (6.8)	14.6	
γ	3.35 m	40.8	2.84 m	42.8	β Me
γ Me	1.40 d (6.8)	16.2	1.33 (6.5)	14.7	CONH ₂
CONH	9.78 br s			173.5	
CONH ₂ ^d	7.50			180.4 ^e	
			AGDHE		
α	4.72 d (8.4)	72.0	3.99 d (9.1)	72.0	
β	4.10 br d (8.4)	75.2	3.65 br d (9.1, 2.2)	75.2	
γ	4.60 overlapped	50.6	4.21 m	50.8	
δ	1.96, 1.78 m	29.6	1.70, 1.60 m	26.1	
ϵ	1.78 m	26.1	1.58 m	25.4	
ζ	3.39 m	42.2	3.19 m	42.1	guan
γ NH	8.30 br s				
ζ NH	8.50 br s				
CO				177.2 ^e	
guan	7.92 overlapped			158.4	

Table 1 (Continued)

	pyr- <i>d</i> ₅ ^b		methanol- <i>d</i> ₄		
	δ_H	δ_C	δ_H	δ_C	COLOC
			Ala		
α	4.80 m	50.4	4.40 overlapped	51.0	CONH, C β CONH, C α , CO-TMHEA
β	1.51 d (7.1)	18.3	1.48 d (7.1)	17.8	
CONH	8.90 d (6.8)			176.2	
			TMHEA		
1				178.7	
2	2.90	44.5	2.68 m	44.7	
3	3.71 dd (8.8, 1.6)	79.5	3.52 dd (8.8, 3.0)	79.5	
4	1.71 m	33.0	1.78 m	33.5	
5	1.31, 1.21 m	38.8	1.26 m	39.3	C9
6	1.50 m	25.7	1.75 m	25.8	
7	0.81 d (6.4)	24.7	0.98 d (6.4)	24.7	C9
8	1.11 d (6.4)	14.6	1.11 d (6.8)	14.4	C2, C4, C1
9	1.01 d (6.8)	17.7	1.01 d (6.8)	17.0	
10	0.74 d (6.4)	21.6	0.91 d (6.8)	21.5	C5

^a Assignments based on 2D-COSY, 2D-HOHAHA, HETCOR, and COLOC experiments. ^b Quaternary carbons in pyr-*d*₅ were not assigned due to the failure of obtaining ¹H-¹³C long range connectivities under various experimental conditions in this solvent. ^c Coupling constants are enclosed in parentheses and given in Hz. ^d Tentative assignments based on NOESY experiments: MeGln, cross peak δ 7.30/1.93; DiMeGln cross peak δ 7.50/3.55. ^e These assignments are interchangeable. ^f AGDHE: 4-amino-7-guanidino-2,3-dihydroxyheptanoyl residue. ^g TMHEA: 3-hydroxy-2,4,6-trimethylheptanoyl residue.

to those of Arg, confirmed by the cross peak between the low field resonating methylene protons at δ 3.19 and the guanidino



(2R,3R,4S)-AGDHE (2)

carbon at δ 158.4 ppm in the COLOC spectrum, whereas COSY and HOHAHA²¹ spectra revealed three continuous methines

implied by the presence of the phenolic group of the β -methoxytyrosine residue. All the chemical and spectroscopic data so far collected indicate a C₆₈H₁₁₆N₁₈O₂₀ molecular formula in agreement with the mass spectral data.

The stereochemistry of standard amino acids was determined by HPLC analysis of the acid hydrolysate derivatized with Marfey's reagent,²⁴ which allowed us to assign the L configuration for the Ala, *N*-MeAla, Leu, and Thr residues and the D configuration for the Arg residue. In our hands the Marfey's derivatives of D-Arg and L-Thr had identical retention times. The configuration of both residues were solved by applying the Marfey's method on the amino acids isolated after acid hydrolysis (see below).

The *S* configuration at C-4 of the new 4-amino-7-guanidino-2,3-dihydroxyheptanoic acid residue (AGDHE 2) was deter-





Chart 2



L-configuration of Thr and the D-configuration of Arg. The β -methoxytyrosine, labile under acid conditions, was the only piece lost during the acid hydrolysis. The ^1H NMR spectrum of the total hydrolysate mixture revealed only dispersed aromatic signals confused in the background of the spectrum.

Amino acid sequence of callipeltin A (1) was deduced by interpretation of the NOESY spectrum in pyridine- d_5 , which gave sequential information from correlations between amide protons and α -protons of the adjacent residue *via* amide bond, as shown in 1. The attachment of the 3-hydroxy-2,4,6-trimethylheptanoyl residue to the α -amino group of the N-terminus Ala, as first indicated by the COLOC correlation between the Ala α -proton and the carbonyl of the heptanoyl residue at 178.7 ppm and supported by a NOESY cross peak between the Ala amide proton and the proton attached to C-2 of the heptanoyl residue, was definitively confirmed by isolation, after mild acid hydrolysis in 6 N HCl at 140 °C for 1 h, of the *N*-(3-hydroxy-2,4,6-trimethylheptanoyl)-L-alanine. The amino acid was characterized by EIMS, m/z 259, M^+ , and ^1H NMR spectroscopy. The remaining connection to be clarified was that in the macrocyclic lactone between the carbonyl group of the C-terminal MeAla, whose α -proton did not give rise to cross peaks with any amide protons in the NOESY spectrum, and one of the hydroxyl groups of the two Thr residues. The β -proton of Thr-2 resonated at low-field, δ 5.60 in methanol- d_4 and 6.07 in pyridine- d_5 , thereby revealing participation of

In order to evaluate the antiviral activity of callipeltin A (1), we studied the inhibition of cytopathic effects (CPE) induced by HIV-1, using the MTT cell viability to determine the CD_{50} (50% cytotoxic dose) and ED_{50} (50% effective dose) as described previously.³⁰ At day six post-infection, callipeltin A (1) exhibited a CD_{50} of 0.29 $\mu\text{g}/\text{mL}$ and a ED_{50} of 0.01 $\mu\text{g}/\text{mL}$ giving a selectivity index (SI ratio $\text{CD}_{50}/\text{ED}_{50}$) of 29. AZT reference has a CD_{50} of 50 μM and a ED_{50} of 30 nM. The antifungal activity of callipeltin A (1) was measured against *Candida albicans*, whose growth was inhibited at 100 $\mu\text{g}/\text{disc}$ (6 mm) with 30 mm of inhibition.

Conclusion

Callipeltin A (1) is a novel antiviral and antifungal cyclodepsipeptide which contains three unusual amino acid residues: (2*R*,3*R*,4*S*)-4-amino-7-guanidino-2,3-dihydroxyheptanoic acid (AGDHE), which is apparently derived by a two carbon atoms homologation of arginine; β -methoxytyrosine, whose stereochemistry waits to be elucidated; and (3*S*,4*R*)-3,4-dimethyl-L-glutamine. All three, to the best of our knowledge, have not been isolated from natural sources. The general structure of callipeltin A (1) with the N-terminus blocked and the C-terminus lactonized with a Thr residue, giving rise to a macrocyclic lactone, is similar in these aspects to other marine cyclodepsipeptides such as, for example, didemnins, a family of potent antitumoral and antiviral compounds from *Trididemnum tuni-*

employing the conventional pulse sequence. The COSY spectra were obtained using a data set ($t_1 \times t_2$) of 1024 \times 512 points for a spectral width of 4201.7 Hz (relaxation delay 1 s) in methanol- d_4 and 5154.6 Hz in pyr- d_5 , respectively. The data matrix was processed using an unshifted sine bell window function, following transformation to give a magnitude spectrum with symmetrization (digital resolution in F2 dimension: 4.10 Hz/pt in methanol- d_4 and 5.03 Hz/pt in pyr- d_5).

The 2D-HOHAHA²¹ (pyr- d_5) experiment was performed in the phase sensitive mode (TPPI) using a MLEV 17 sequence for mixing. The spectral width (t_2) was 5434.8 Hz; 512 experiments of 48 scans each (relaxation delay 1.5 s, mixing time 100 ms) were acquired in 1K data points. For processing, an unshifted sine bell window function was applied in both dimension before transformation. The resulting digital resolution in F2 was 5.30 Hz/pt.

The NOESY (pyr- d_5) experiment was performed in the phase-sensitive mode (TPPI). The spectral width (F₂) was 6172.8 Hz; 512 experiments of 64 scans each (relaxation delay 1.0 s, mixing time 400 ms) were acquired in 2K data points. For processing, a sine bell window function was applied in both dimension before transformation. The resulting digital resolution in F2 was 3.01 Hz/pt.

¹³C, ¹H shift correlation experiment (HETCOR) was performed in methanol- d_4 (125 MHz). The spectral width in ¹³C dimension was 17 241.3 (1024 points) and 3703.7 Hz (128 time increments) along the ¹H domain; for each FID 256 scans were recorded. The digital resolution in F2 was 37.5 Hz/pt.

The HMQC experiment (pyr- d_5) was performed according to Bax et al.²² The spectral width in ¹H dimension was 3703.7 Hz; 512 experiments of 32 scans each were acquired in 1K data points (relaxation delay 1.5 s). A sine square window function was applied in both dimension before transformation (digital resolution in F2 was 3.62 Hz/pt).

¹³C, ¹H long range shift correlation experiment (COLOC²³) was performed in methanol- d_4 (125 MHz). Spectral width in F2 was 22 727.23; 256 experiments of 128 scans were acquired in 1K data points (relaxation delay 1.5 s). The digital resolution in F2 was 22.2 Hz/pt.

Optical rotations were measured on a Perkin-Elmer 141 polarimeter using a sodium lamp operating at 589 nm. Fast atom bombardment mass spectra (FAB MS) were recorded in a glycerol-matrix in the positive ion mode on a VG ZAB instrument (Argon atoms of energy of 2–6 KV). IR spectrum was performed on a IFS 48 Bruker instrument. UV spectra were recorded on a Beckman DU70 spectrophotometer, and CD spectra were recorded on a JASCO 710 spectrometer.

Molecular Mechanics Calculations. All the calculations were carried out on a SGI Personal Iris 35/G using the force field CHARMM (QUANTA 4.0 software package). The two diastereomeric molecular models for the AGDHE residue (2R,3R,4S) and (2S,3S,4S) were generated by imposing a hydrogen bond between the amide carbonyl and the hydroxyl group attached to C-3. Then the lowest energy conformation for each diastereomer was searched by energy minimizing a number of different initial conformations. Energy minimization was performed using a conjugate gradient algorithm and a distance dependent dielectric (to partially compensate for the absence of the solvent) until the energy gradient was less than 0.001 Kcal/mol. The two conformations (one for each diastereomer) displaying the lowest potential energy were selected and the ϕ dihedral angle (H β –C β –C γ –H γ) measured, so that the coupling constants for the two stereoisomers could be calculated (by means of a modified Karplus equation) and then compared to the corresponding experimental value. The same procedure was applied for searching the lowest energy minimum conformations of the diastereomeric β -hydroxy acid (3).

Isolation of Callipeltin A (1). *Callipelta* sp. (Demospongiae, Lithistida, Corallistidae) was collected in 1992 and 1993 in the shallow waters of East coast of New Caledonia. Taxonomic identification was performed by Professor Claude Lévi, Muséum National d'Histoire Naturelle, Paris, France, and reference specimens are on file (reference 1572) at the ORSTOM Centre of Nouméa. Preliminary tests of bioactivity on polar extracts showed antifungal activity against *Fusarium oxysporum*, *Helminthosporium sativum*, and *Phytophthora hevea*, cytotoxic activity against KB and P388 cells (10 μ g/mL, ca. 80% inhibition, in both cases), and anti-HIV activity.

The organism were freeze-dried, and the lyophilized material (700 g) was extracted with *n*-hexane and CH₂Cl₂ in a Soxhlet apparatus and then with 8:2 CH₂Cl₂:MeOH (3 \times 2 L) at room temperature. The dichloromethane–methanol extract was filtered and concentrated under reduced pressure to give 10 g of a brown amorphous solid. The crude dichloromethane–methanol extract was applied in five runs to a DCCC apparatus (7:13:8 CHCl₃:MeOH:H₂O, ascending mode; fractions of 4 mL were collected). Fractions 13–17, mainly containing callipeltin A (1), were further purified by HPLC on a Waters C-18 μ -Bondapak column (7.8 mm i.d. \times 30 cm) with MeOH:H₂O (50:50) as eluent to give 500 mg of pure 1 (t_r = 10.8 min).

Callipeltin A (1): [α]_D = +3.56° (c 0.012 M, MeOH); UV (MeOH) λ_{max} 232 (ϵ = 6919), 274 (ϵ = 1272); IR (KBr) 3330, 1740, 1660, 1520 cm⁻¹; ¹H and ¹³C NMR in Table 1; FABMS and ESMS in the text.

Amino Acid Analysis of Callipeltin A (1). For standard amino acid analysis, 100 μ g of callipeltin A (1) was dissolved in 0.5 mL of 6 N HCl in a evacuated glass tube and heated at 110 °C for 16 h. After evaporation, the residue was dissolved in 0.5 mL of HCl and subjected to amino acid analysis on a Beckman 118BL system. Retention times in the amino acid analysis (min) were as follows: Thr (18.9), Ala (33.3), Leu (44.2), Arg (102.4), NH₃ (160.0).

For a large scale hydrolysis, a 50 mg sample of callipeltin A (1) dissolved in 5 mL of 6 N HCl and heated at 130 °C for 4 h. The crude hydrolysate was extracted (2 \times 1 mL) with CH₂Cl₂. The CH₂Cl₂ layer afforded 4 mg of 3-hydroxy-2,4,6-trimethylheptanoic acid (3); FABMS *m/z* 189 [M + H]⁺; ¹H NMR (CDCl₃) δ 0.84 (3H, d, J = 6.8 Hz, H-9), 0.92 (3H, d, J = 6.8 Hz, H-10), 0.96 (3H, d, J = 5.8 Hz, H-7), 1.12–1.18 (2H, m, H-5 and H-5'), 1.26 (3H, d, J = 7.7 Hz, H-8), 1.64 (1H, m, H-6), 1.70 (1H, m, H-4), 2.73 (1H, quintet, J = 7.7 Hz, H-2), 3.48 (1H, dd, J = 4.1, 7.7 Hz, H-3). The aqueous layer was fractionated by DCCC [10:10:1:6 CHCl₃:MeOH: *n*BuOH:H₂O (0.1% TFA), descending mode], and 4 mL fractions were collected. Fractions 18–43 (A) contained a mixture of 3,4-dimethylpyroglutamic acid and *N*-methylpyroglutamic acid; fractions 59–70 (B) contained Leu, fractions 143–158 (C) contained a mixture of Ala and Me-Ala, fractions 197–217 (D) contained Thr and, fractions 231–241 (E) contained a mixture of Arg and 4-amino-7-guanidino-2,3-dihydroxyheptanoic acid (2); FABMS *m/z* 252 (4-amino-7-guanidino-2,3-dihydroxyheptanoic acid, [M + NH₄]⁺); ¹H NMR (pyr- d_5) δ 1.90 (4H, m, H-6 AGDHE and H-4 Arg), 2.15 (2H, m, H-5 AGDHE and H-3 Arg), 2.25 (2H, m, H-5' AGDHE and H-3' Arg), 3.30 (4H, m, H-7 AGDHE and H-5 Arg), 4.13 (2H, m, H-4 AGDHE and H-2 Arg), 4.59 (1H, dd, J = 7.5, 3.0 Hz, H-3 AGDHE), 4.98 (1H, d, J = 7.5 Hz, H-2 AGDHE). Fraction A was further purified by HPLC on a Vydac 218 TP protein-peptide RP column (flow rate 4 mL/min, eluent: H₂O (0.1% TFA) for 10 min, CH₃CN–H₂O (0.1% TFA) 10–80% in 30 min) to give 0.5 mg of pure *N*-methylpyroglutamic acid (t_r = 23.2 min) and 0.4 mg of pure 3,4-dimethylpyroglutamic acid (t_r = 23.4 min).

N-Methylpyroglutamic acid: CD [θ]_{216nm} = +3151 (c 0.13 10⁻⁴ M, H₂O); ¹H NMR (pyr- d_5) δ 2.05 (1H, m, H-3), 2.12 (1H, m, H'–3), 2.20 (1H, m, H-4), 2.40 (1H, m, H-4'), 2.88 (3H, s, N-Me), 4.12 (1H, dd, J = 9.3, 3.3 Hz, H-2); ¹³C NMR: δ 22.6, 28.4, 29.2, 61.7, 174.5 and 175.0. The compound shows identical spectral properties with a synthetic sample obtained by heating in water under reflux overnight the commercial *L*-*N*-methylglutamic acid.²⁶

3,4-Dimethylpyroglutamic acid: CD [θ]_{208nm} = +1534 (c 0.001 M, H₂O); ¹H NMR (pyr- d_5) δ 1.01 (3H, d, J = 6 Hz, Me on C-3), 1.04 (3H, d, J = 6.8 Hz, Me on C-4), 2.65 (1H, m, H-4) 2.71 (1H, m, H-3), 4.02 (1H, d, J = 3.8 Hz, H-2). A 0.1 mg aliquot of compound was treated with 50 μ L of MTBSTFA [*N*-methyl-*N*-(*tert*-butyldimethylsilyl)-trifluoroacetamide] in 0.5 μ L of CH₃CN at 150 °C for 150 min. After cooling to room temperature the solution was directly applied to GC-MS under conditions reported by Des Neves et Vasconcelos;²⁸ *m/z* 370, 328, 229, 385, 300.

Determination of Absolute Configuration. (a) HPLC Analysis of the Marfey's Derivatives.²⁴ To a 10% acetone solution (50 μ L) of 1-fluoro-2,4-dinitrophenyl-5-alanine amide (*L*-FDAA) and 1 N NaHCO₃ (40 μ L) was added 100 μ g of the peptide acid hydrolysate, and the mixture was kept at 50 °C for 1 h. After cooling to room temperature 2 N HCl (40 μ L) was added, and the resulting solution was taken to dryness and then dissolved in 500 μ L of DMSO. A 5 μ L aliquot of

the FDAA derivatives was analyzed by reversed-phase HPLC. A linear gradient from triethylammonium phosphate (50 nM, pH 3.0)/MeCN 90% to triethylammonium phosphate (50 nM, pH 3.0)/MeCN 50% over 45 min (flow rate 2 mL/min) was used to separate the FDAA derivatives which were detected by UV at 340 nm²⁴. Amino acids were derivatized as described above. The peaks were identified by co-injection with a DL-mixture of standard amino acids. Due to the overlapping of L-Thr and D-Arg peaks, we used the DCCC enriched fractions D and E for the identification of these amino acids. Retention times (min) are given in parentheses. L-Thr and D-Arg (18.1), L-Ala (21.4), L-MeAla (28.3), L-Leu (37.6).

(b) **Periodate Oxidation of Callipeltin A (1).** A solution of callipeltin A (1) (1.5 mg) and sodium periodate (200 mg) in water (0.5 mL), adjusted to pH 4.0 with acetic acid, was stirred overnight at 25 °C. The solvent was lyophilized, and the residue was dissolved in hydrogen peroxide (2 mL) and formic acid (1 mL). The solution was heated at 70 °C for 20 min and cooled, and the solvent was evaporated under reduced pressure. The reaction product was subjected to acid hydrolysis and Marfey's derivatization as described above. An additional peak, corresponding to L-Arg (retention time: 16.4 min), was observed in the HPLC trace of Marfey's derivatives.

(c) **R (+) and S (-) Mosher Esters of 3.** Freshly distilled (+)- α -methoxy- α -(trifluoromethyl)phenylacetic (MTPA) chloride (6 μ L) was added to a solution of **3** (2 mg) with a catalytic amount of 4-(dimethylamino)pyridine and 5 μ L of freshly distilled triethylamine, in 1 mL of freshly distilled CH₂Cl₂ and allowed to stand at room temperature for 12 h under argon atmosphere. The residue, obtained after evaporation of the solvent, was subjected to reverse-phase HPLC using a linear gradient from water to CH₃CN (100%), UV detector: λ = 260 nm, to obtain 0.5 mg of (R)-(+)-MTPA ester of **3**. The same procedure was used to obtain 0.4 mg of (S)-(-)-MTPA ester of **3**. 3-(+)-(R)-MTPA ester of **3**: ¹H NMR (500 MHz, CDCl₃) δ 0.78 (3H, d, *J* = 5.8 Hz, H-7), 0.87 (3H, d, *J* = 6.8 Hz, H-10), 0.92 (3H, d, *J* = 6.8 Hz, H-9), 1.10 (2H, m, H-5 and H-5'), 1.36 (3H, d, *J* = 7.5 Hz, H-8), 1.62 (1H, m, H-6), *ca.* 1.70 (1H, overlapped with H₂O, H-4), 3.16 (1H, quintet, H-2), 3.70 (1H, dd, H-3). 3-(-)-(S)-MTPA ester of **3**: ¹H NMR (500 MHz, CDCl₃) δ 0.82 (3H, d, *J* = 5.8 Hz, H-7), 0.90 (3H, d, *J* = 6.8 Hz, H-10), 0.97 (3H, d, *J* = 6.8 Hz, H-9), 1.13 (2H, m, H-5 and H-5'), 1.32 (3H, d, *J* = 7.5 Hz, H-8), 1.64 (1H, m, H-6), *ca.* 1.70 (1H, overlapped with H₂O, H-4), 3.14 (1H, quintet, H-2), 3.72 (1H, dd, H-3).

Methylation of Callipeltin A (1). A solution of 1 mg of callipeltin A (1) in MeOH was treated with an excess of CH₃N₂ in ether at room temperature for 1 h. The solution was taken to dryness under a stream of N₂ and analyzed by FABMS, *m/z* 1519 [M + H]⁺, and ¹H NMR: the spectrum contained a new methyl signal at δ 3.80 (3H, s).

Methanolysis of Callipeltin A (1) Giving the Opened Methyl Ester (4). A solution of 1 mg of callipeltin A (1) was treated with 1.1 mg of NaOMe in dry methanol (0.2 mL) at room temperature for 2 h. The reaction mixture was neutralized with 0.1 N HCl, poured into ice-water, and then extracted with *n*-BuOH. The *n*-BuOH phase was evaporated under reduced pressure and the crude product (0.8 mg) was directly subjected to FABMS analysis. FAB-MS fragmentations are shown in 4.

Partial Hydrolysis of Callipeltin A (1) Giving the N-(3-Hydroxy-2,4,6-trimethylheptanoyl)-L-Alanine Residue. Callipeltin A (1) (30 mg) was treated with 2 N HCl (2 mL), and after degassing the reaction mixture was heated at 145 °C for 1 h in a sealed tube. The reaction mixture was extracted with CH₂Cl₂ (3 \times 2 mL) and the dichloromethane layer afforded 1.5 mg of N-(3-hydroxy-2,4,6-trimethylheptanoyl)-L-alanine: EIMS *m/z* 259 (M⁺); ¹H NMR (methanol-*d*₄) δ 0.87 (3H, d, *J* = 6.4 Hz, H-13), 0.94 (3H, d, *J* = 6.5 Hz, H-14), 0.95 (3H, d, *J* = 6.5 Hz, H-10), 1.11 (1H, ddd, *J* = 14.0, 9.0 and 4.5 Hz, H-8), 1.18 (3H, d, *J* = 7.1 Hz, H-12), 1.33 (1H, ddd, *J* = 14.0, 9.5 and 3.0 Hz, H-14), 1.41 (3H, d, *J* = 7.5 Hz, H-11), 1.67 (2H, m, H-7 and H-9), 2.60 (1H, quintet, *J* = 7.1 Hz, H-5), 3.67 (1H, t, *J* = 7.1 Hz, H-6), 4.38 (1H, q, *J* = 7.5 Hz, H-2).

Acknowledgment. This contribution is part of the EEC project "Marine Sciences and Technology MAST III" Contract MAS 3-CT95-0032. Mass spectra were provided by the "Servizio di Spettrometria del CNR e dell'Università di Napoli". The staff is acknowledged. The NMR spectra were recorded at CRIAS "Centro Interdipartimentale di Analisi Strumentale", Faculty of Pharmacy, University of Naples. We thank the diving team of the ORSTOM Centre de Noumèa for the collection of the sponge (ORSTOM-CNRS SMIB Program) and Professor Claude Lévi, Muséum National d'Histoire Naturelle, Paris, France, for the identification of the sponge. Dedicated to Professor Mario Piattelli, Università di Catania, on the occasion of his 70th birthday.

Supporting Information Available: ¹H NMR, ¹³C NMR, 2D-COSY, and 2D-NOESY spectra of callipeltin A (1) (12 pages). See any current masthead page for ordering and Internet access instructions.

JA954287P

