

# Birth Spacing and Child Survival in Rural Senegal

CARINE RONSMANS

*enfant espacé à l'époque*

Ronsmans C (Maternal and Child Epidemiology Unit, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT, UK). Birth spacing and child survival in rural Senegal. *International Journal of Epidemiology* 1996; 25: 989-997.

**Background.** Studies examining the associations between short birth spacing and child mortality have often concentrated on the strength of the associations whilst the public health importance of short spacing in specific communities has received less attention. This study re-examines the association between short birth intervals and child mortality in rural Senegal and discusses the potential direct effects of efforts to delay births on child mortality in this community.

**Methods.** The study uses longitudinal data in a cohort of 4852 children born between 1983 and 1989. The associations between birth spacing and child mortality are examined using logistic and Cox proportional hazards regression models.

**Results.** The probability of dying before age five is 224 per 1000 livebirths. The median interval between births is 33 months and only 12% of the birth intervals are less than 24 months in length. The odds of dying in the neonatal and post-neonatal period is 2.27 and 2.12 times higher respectively for children born after preceding birth intervals of one year or less compared to children born after longer intervals. Children born within two years of a subsequent birth are at 4.09 times higher risk of dying in the second year of life than children whose mother gave birth more than 2 years after the index birth.

**Conclusions.** In this community where prolonged breastfeeding causes women to space their births at long intervals, short birth intervals are a consequence rather than a cause of child mortality and the potential direct effects of birth spacing efforts on child mortality are limited. To reduce the high levels of child mortality, efforts will have to be made to ensure effective preventive and curative health services, and to maintain the traditional pattern of breastfeeding.

**Keywords:** birth intervals, birth spacing, child mortality, family planning, developing countries, Senegal

The associations between short intervals between births and impaired child survival are well established.<sup>1-4</sup> It is generally accepted that if closely spaced births were delayed, particularly in countries where mortality and fertility are still high, child mortality levels would fall.<sup>5</sup> Increasing the interval between births to improve the health of women and children has thus become one of the primary rationales for family planning programmes.<sup>5,6</sup>

Despite the existence of a large body of evidence on the association between birth spacing and child survival, many questions remain regarding the strength and nature of the association, and the magnitude of mortality reductions that can be expected from family planning programmes are still the subject of debate.<sup>7-9</sup> Most studies have been based on retrospective surveys, and recall biases may have led to an overestimation of the strength of the association between short birth intervals and mortality.<sup>10</sup> In addition, the potential

confounding effect of unmeasured variables, particularly prematurity,<sup>11</sup> but also socio-economic characteristics of the households, may call into question the existence of a causal association. Finally, empirical evidence on the causal pathways that explain the associations is poor.<sup>3,12,13</sup>

This study re-examines the association between short birth intervals and mortality in childhood using longitudinal data from rural Senegal. The objectives of the study are (1) to describe the distribution of birth intervals; (2) to explore the extent to which a child death shortens the length of time between subsequent births; and (3) to examine the association between a short preceding birth interval and neonatal and post-neonatal mortality and between a short subsequent birth interval and mortality after age one. The findings are discussed in the context of the potential effects of child spacing efforts on child mortality.

## METHODS

### Population

The study population lives in Niakhar, a rural area located about 150 km east of Dakar, the capital of

At the time of this study the author was a doctoral student at the Harvard School of Public Health, Boston, USA.  
Reprint requests: Carine Ronsmans, Maternal and Child Epidemiology Unit, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT, UK.





Senegal. The population consists largely of a single ethnic group, the Sereer, who are subsistence farmers growing millet and peanuts. The farming activities are almost entirely concentrated in the rainy season from July to October. The population is mostly Muslim and the level of education is poor.

The study area has about 25 000 inhabitants and has been under demographic surveillance since March 1983.<sup>14</sup> From 1983 to 1987, yearly censuses were conducted to obtain information on births, deaths and migration among all residents of the study area, and from January 1987 onwards, demographic data were collected by weekly visits to all households.<sup>15,16</sup>

Prolonged breastfeeding is widespread and there was no deliberate control of family size during the period studied. The average age at weaning was 23 months.<sup>14</sup> From 1983 onwards, two physicians visited the area regularly and provided medical care and advice. From January 1987, a physician was living permanently in Niakhar. Vaccine coverage was low in 1984, but reached almost 100% in 1989.<sup>15</sup>

#### *Sample Selection and Analysis*

The analysis consists of four parts. For each part, separate samples were selected from the 4852 single births which occurred between March 1983 and December

pregnant again, children who died before the estimated date of conception of the subsequent birth were excluded. The date of conception was estimated by subtracting 280 days from the date of birth. Among those children whose mother had a subsequent birth, 2161 survived after the first year of life and 1963 after the second year of life. Of those, 103 children aged 12–23 months and 18 aged 24–35 months died before the estimated date of the subsequent conception, and were excluded from the analysis of the effect of subsequent birth intervals.

Mortality was expressed per 1000 livebirths for neonatal mortality and per 1000 child-years at risk for all other age groups. In addition to the duration of birth intervals, the following variables were studied. For all age groups, mortality was analysed by birth order and sex of the child, by maternal age and literacy (whether the mother could read or write), and by season. The seasonal risks were expressed as binary variables, referring to the month of birth for the neonatal period and to the month of exposure for other age groups. For neonatal mortality, the high risk season was defined from June through October, reflecting the months of seasonal weight losses in women in this area.<sup>15</sup> For mortality in the other age groups, the high risk season was taken to extend from August through November




FIGURE 1 Distribution of birth interval length by age at death of the index child

eliminating the least significant variables and retaining those in which the regression coefficients yielded a *P*-value of  $\leq 0.10$ .

RESULTS

*Levels of Mortality*

The probability of dying in the first year of life is 103.5 per 1000 livebirths and for children surviving after one year of age, the probability of dying before age five is 134.3 per 1000. The probability of a newborn child dying before its fifth birthday is 223.8 per 1000.

*Distribution of the Length of Birth Intervals*

child is weaned off the breast and thus placed at greater risk of infection and undernutrition when its mother becomes pregnant again. It may also be related to the effect of a child death on the return of ovulation.<sup>13,19</sup> One way to separate these two mechanisms is to calculate the proportion of children who died *before* the estimated date of conception of the next child, since only the latter can influence the timing to the next birth. The majority (413/501) of the children who died before 2 years of age and were followed by another birth, died *before* the estimated date of conception of the next child, hence their death will have resulted in the next pregnancy rather than having been caused by it. Although the large majority of neonatal (96%) and post-

TABLE 1 *Crude differentials in neonatal and post-neonatal mortality by selected variables*

Variable	Neonatal mortality <sup>a</sup> (number of deaths) [number of observations]	Post-neonatal mortality <sup>a</sup> (number of deaths) [number of observations]
Duration of preceding birth interval (months)		
≤12	79.54* (7)[88]	105.34 (7)[81]
13-24	38.08 (19)[499]	53.97 (21)[477]
25-36	35.02 (52)[1485]	50.06 (56)[1420]
37-48	42.94 (14)[326]	54.29 (13)[310]
≥49	119.40** (8)[67]	55.37 (2)[58]
Immediately preceding child <sup>b</sup>		
survived	38.94 (90)[2311]	52.33 (83)[2202]
died	64.94 (10)[154]	60.84 (16)[144]
Season of birth/exposure <sup>b</sup>		
low risk	33.53 (45)[1342]	34.88 (41)
high risk	48.98 (55)[1123]	84.84** (58)
Birth order		
2-5	39.19 (52)[1327]	48.58 (49)[1261]
6-9	32.77 (31)[946]	59.29 (42)[912]
≥10	88.54** (17)[192]	60.15 (8)[173]
Maternal age		
15-19	40.00 (2)[50]	97.56 (4)[2081]
20-29	35.66 (44)[1234]	55.33 (50)[1176]
30-39	34.99 (33)[943]	43.63 (31)[906]
≥40	88.23** (21)[238]	86.42* (14)[216]
Sex		
boy	45.60 (56)[1228]	55.07 (50)[1159]
girl	35.57 (44)[1237]	52.02 (49)[1187]
Maternal literacy		
yes	32.26 (2)[64]	42.09 (2)[60]
no	40.70 (98)[2401]	53.82 (97)[2286]
Year of birth		
1984-1985	21.74 (4)[184]	49.70 (8)[180]
1986-1987	44.50 (49)[1101]	61.48 (57)[1041]
1988-1989	39.83 (47)[1180]	44.53 (34)[1125]
All	40.57 (100)[2465]	53.52 (99)[2346]

<sup>a</sup> Neonatal mortality is expressed in deaths per 1000 livebirths, post-neonatal mortality in deaths per 1000 child-years at risk.

<sup>b</sup> See text.

\*  $P < 0.05$ , \*\*  $P < 0.01$  when compared to category of lowest risk.

mother was 40 years or older when she gave birth, and in children born at birth order 10 or more. Similar patterns are observed for post-neonatal mortality, although the associations only reach significance at the 5% level for seasonal exposure and old maternal age.

The multivariate analysis does not modify the findings, except for birth order which does not retain

statistical significance (Table 2). Children born within one year after their older sibling was born are twice as likely to die in the neonatal and post-neonatal period than children born after longer intervals. The most striking feature of this analysis is the strong seasonality of post-neonatal mortality: in the rainy season, children are at 2.58 times higher risk of dying than in the dry season.

TABLE 2 *Multivariate analysis of neonatal and post-neonatal mortality*

Variable	Neonatal mortality Odds ratio (95% CI)	Post-neonatal mortality Hazard ratio (95% CI)
Duration of preceding birth interval (months)		
≤12	2.27 (1.01-5.10)*	2.12 (0.98-4.54)
13-48	1.00	1.00 <sup>a</sup>
≥49	3.05 (1.39-6.68)**	-
Season of birth/exposure		
low risk	1.00	1.00
high risk	1.44 (0.96-2.16)	2.58 (1.73-3.85)**
Maternal age		
15-19	1.12 (0.27-4.72)	1.94 (0.71-5.29)
20-39	1.00	1.00
≥40	2.52 (1.51-4.20)**	1.79 (1.01-3.16)*

<sup>a</sup> Reference is ≥13 months.

\*  $P < 0.05$ , \*\*  $P < 0.01$ .

#### *Effects of the Length of the Subsequent Birth Interval*

The duration of the subsequent birth interval clearly affects mortality in the second year of life and to a lesser extent, but not significantly so, in the third year of life (Tables 3 and 4). If the child's mother delivers another child within 2 years of the birth of the index child, the risk of mortality is four times higher in the second year of life than if the next birth takes place more than 2 years after the index child's birth. In addition, the rainy season raises the risk twofold in the second year of life and almost threefold in the third year of life. Trends in mortality over time were impressive: mortality in both age groups almost halved over the study period. Birth order, age of the mother, sex of the child or literacy, on the other hand, do not seem to affect childhood mortality to a large extent.

#### DISCUSSION

The findings of this study add evidence to the widely observed association between the length of birth intervals and child survival. In this population, a short preceding interval increases the risk of neonatal and to a lesser extent post-neonatal mortality, and a short subsequent interval increases the risk of mortality in childhood.

There are, however, two observations in this study which differ from what has been found in other studies, and which have implications for the potential effect of spacing programmes on mortality. Firstly, preceding birth intervals of 1-2 years did not affect neonatal or post-neonatal mortality. The latter finding merits

attention, since it is generally believed that children throughout the developing world are much more likely to die if they were born less than 2 years after their mother's previous birth.<sup>2,4</sup> Secondly, short succeeding birth intervals, in contrast, have major adverse consequences for child survival, and the size of the effect is larger than those found by other authors.<sup>2,4,20</sup>

The lack of an association between preceding intervals of between one and two years and neonatal and post-neonatal mortality may in part be due to the small sample in this risk group and in part to the fact that short birth intervals almost invariably follow an infant death. If there is no surviving young child to compete for the mother's resources at the time of birth of the next child, the adverse effect of short spacing may be reduced. However, similar trends have been found in an earlier study in the same population using a much larger sample and incorporating a control for the survival status of the immediately preceding child.<sup>21</sup> In addition, many studies have found a stronger association between a short preceding birth interval and neonatal or post-neonatal mortality when the preceding sibling died than when it survived.<sup>3,20,22</sup> In fact, there is also evidence from Bangladesh that only very short preceding birth intervals affect neonatal and post-neonatal mortality.<sup>3,11</sup>

The association between preceding intervals of one year or less and neonatal and post-neonatal mortality raises concerns about the confounding effect of prematurity. Very short birth intervals are likely to include a large proportion of infants born after a short gestation, which affects child survival, and removing the confounding effect of prematurity may reduce the estimates

TABLE 3 Crude differentials in mortality between 12 and 23 months and between 24 and 35 months by selected variables

Variable	Mortality between 12 and 23 months <sup>a</sup> (number of deaths) [number of observations]	Mortality between 24 and 35 months <sup>a</sup> (number of deaths) [number of observations]
Duration of subsequent birth interval (months) <sup>b</sup>		
≤24	180.59** (40)[243]	54.32 (9)[197]
25-30	46.70 (30)[655]	39.09 (22)[619]
31-36	-	40.09 (29)[751]
37-42	-	8.09** (2)[240]
Season of exposure <sup>c</sup>		
low risk	64.38 (85)	25.98 (31)
high risk	120.27** (90)	74.97** (49)
Birth order		
1	71.61 (15)[220]	48.09 (9)[202]
2-5	85.79 (102)[1244]	38.61 (41)[1131]
6-9	82.95 (49)[624]	49.53 (26)[566]
≥10	135.56 (9)[73]	70.68 (4)[64]
Maternal age		
15-19	81.61 (15)[194]	55.00 (9)[177]
20-39	83.45 (149)[1870]	40.29 (64)[1700]
≥40	124.19 (11)[97]	89.43 (7)[86]
Sex		
boy	90.54 (98)[1140]	42.28 (41)[1032]
girl	97.13 (77)[1021]	45.30 (39)[931]
Maternal literacy		
illiterate	155.16 (11)[103]	80.00 (6)[53]
literate	85.14 (175)[2161]	43.70 (80)[1963]
Year of birth		
1983-1984	115.31** (103)[951]	57.21** (46)[844]
1985-1986	61.17 (59)[1001]	32.15 (29)[927]
1987-1988	65.57 (13)[209]	39.95 (5)[192]
All	85.14 (175)[2161]	43.70 (80)[1963]

<sup>a</sup> Mortality is expressed as deaths per 1000 child-years at risk.

<sup>b</sup> Excluding children who died before the estimated date of conception of the next birth (excluding 103 children for mortality age 12-23 months and 18 children for mortality age 24-35 months). Rates are not shown for cells in which the subsequent birth could not affect mortality in the age group of interest.

<sup>c</sup> See text.

\*  $P < 0.05$ . \*\*  $P < 0.01$  when compared to category of lowest risk.

TABLE 4 *Multivariate analysis of mortality between 12 and 23 months and between 24 and 35 months*

Variable	Mortality between 12 and 23 months Hazard ratio (95% CI)	Mortality between 24 and 35 months Hazard ratio (95% CI)
Duration of subsequent birth interval (months)		
≤24	4.09 (2.54–6.58)**	1.38 (0.65–2.96)



abstinence.<sup>24</sup> Although it is commonly believed that traditional birth spacing practices may be abandoned as a society develops, there is very little empirical evidence to support this hypothesis.<sup>25</sup> In this population, as is the case in Senegal as a whole, there is no indication that the duration of breastfeeding has been declining over time.<sup>14,26</sup>

When the direct effects of child spacing on child survival are viewed in the context of other factors associated with child survival, such as seasonal variations in mortality, their relative importance diminishes even more. The high child mortality rates in the rainy season in West Africa have been documented many decades ago, and are probably associated with a higher incidence of infectious diseases such as malaria, acute respiratory and diarrhoeal diseases, compounded by the nutritional stresses while awaiting the upcoming harvest.<sup>18</sup> The fact that the large reductions in infant and child mortality in this population have not brought about perceptible reductions in seasonally high mortality rates indicates that much remains to be done. The focus on preventable diseases such as measles and pertussis through immunization programmes may have had a more pronounced effect on child mortality in the dry than in the rainy season. Prevention of infectious diseases associated with the rainy season, on the other hand, may be difficult and the reduction of mortality from these diseases may require the provision of effective and accessible curative services.<sup>27,28</sup>

Family planning programmes can of course do much more than increase the intervals between births.<sup>9</sup> Although difficult to measure, the most important public health consequences of reduced fertility rates are probably improved economic well-being and reduced obstetric mortality and morbidity. However, family planning services are sometimes presented as a more or less exclusive answer to situations that in fact can only change if other conditions are fulfilled. The low levels of education in this population, and the high prevailing levels of child mortality—one out of four newborn children does not survive to its fifth birthday—indicate the need for an increased effort to educate women and men, to provide an adequate health infrastructure, and to guarantee effective preventive and curative health services.

#### ACKNOWLEDGEMENT

This paper is a modified version of a chapter of the author's doctoral dissertation. Comments and suggestions from Michel Garenne, John Cleland and Kate Stewart have been very helpful. The data used in this study were collected by the 'Unité de Recherche

Population et Santé' at ORSTOM. The author's doctoral studies were supported by the Nutricia Research Foundation, The Netherlands, and the Alumni of the Harvard School of Public Health, USA.

#### REFERENCES

- <sup>1</sup> Yerusalmly J. On the interval between successive births and its effect on survival of infant. I. An indirect method of study. *Hum Biol* 1945; **17**: 65–106.
- <sup>2</sup> Hobcraft J N, McDonald J W, Rutstein S O. Demographic determinants of infant and early child mortality: a comparative analysis. *Popul Stud* 1985; **39**: 363–85.
- <sup>3</sup> Koenig M A, Phillips J F, Campbell O M, D'Souza S. Birth intervals and childhood mortality in rural Bangladesh. *Demography* 1990; **27**: 251–65.
- <sup>4</sup> Boerma J T, Bicego T G. Preceding birth intervals and child survival: searching for pathways of influence. *Stud Fam Plann* 1992; **23**: 243–56.
- <sup>5</sup> World Development Report. *Investing in Health*. Oxford: Oxford University Press, 1993.
- <sup>6</sup> Locoh T. *Vingt ans de planification familiale en Afrique Sub-Saharienne*. Les Dossiers du CEPED No.19. Paris: CEPED, 1992.
- <sup>7</sup> Bongaarts J. Does family planning reduce infant mortality rates? *Popul Dev Rev* 1987; **13**: 323–34.
- <sup>8</sup> Trussell J. Does family planning reduce infant mortality? An exchange. *Popul Dev Rev* 1988; **14**: 171–78.
- <sup>9</sup> Potter J E. Does family planning reduce infant mortality? *Popul Dev Rev* 1988; **14**: 179–87.
- <sup>10</sup> Potter J E. Birth spacing and child survival: a cautionary note regarding the evidence from the WFS. *Popul Stud* 1988; **42**: 443–50.
- <sup>11</sup> Miller J E, Trussell J, Pebley A R, Vaughan B. Birth spacing and child mortality in Bangladesh and the Philippines. *Demography* 1992; **29**: 305–18.
- <sup>12</sup> Winikoff B. The effects of birth spacing on child and maternal health. *Stud Fam Plann* 1983; **14**: 231–45.
- <sup>13</sup> Zimmer B G. Consequences of the number and spacing of pregnancies on outcome, and of pregnancy outcome on spacing. *Soc Biol* 1979; **26**: 161–78.
- <sup>14</sup> Garenne M, Maire B, Fontaine O, Dieng K, Briend A. *Risques de décès associés à différents états nutritionnels chez l'enfant d'âge préscolaire*. Dakar, Senegal: ORSTOM, 1987.
- <sup>15</sup> Becker C, Chanazarian A, Delaunay V, Preziosi M, Samb B, Simondon F, Simondon K B. *Population et santé à Niakhar: Niveaux et tendances des principaux indicateurs démographiques et épidémiologiques de la zone d'étude 1984–1991*. Dakar, Senegal: ORSTOM, 1992.
- <sup>16</sup> Garenne M, Leroy O, Beau J P, Whittle H, Sene I, Sow A R. *Efficacy, safety, and immunogenicity of two high titer measles vaccines*. Dakar, Senegal: ORSTOM, 1991.
- <sup>17</sup> Newell C. *Methods and Models in Demography*. New York: The Guilford Press, 1988.
- <sup>18</sup> Cantrelle P, Diop I L, Garenne M, Gueye M, Sadio A. IV. The profile of mortality and its determinants in Senegal, 1960–1980. In: *Determinants of Mortality Change and Differentials in Developing Countries*. New York: United Nations, 1985 (UN Population Studies-E.85.XIII.4).
- <sup>19</sup> Santow G, Bracher M D. Child death and time to the next birth in Central Java. *Popul Stud* 1984; **38**: 241–53.

- <sup>20</sup> De Sweemer C. The influence of child spacing on child survival. *Popul Stud* 1984; **38**: 47-72.
- <sup>21</sup> Cantrelle P, Leridon H. Breast feeding, mortality in childhood and fertility in a rural zone in Senegal. *Popul Stud* 1971; **25**: 505-33.
- <sup>22</sup> Cleland J G, Sathar Z A. The effect of birth spacing on childhood mortality in Pakistan. *Popul Stud* 1984; **38**: 401-18.
- <sup>23</sup> Cleland J, Rutstein S. Contraception and birthspacing. *Int Fam Plann Perspect* 1986; **12**: 83-90.
- <sup>24</sup> Rosetta L. Intervalles intergénésiques et indicateurs de santé dans une population Sérère au Sénégal. *Bulletins et Mémoires de la Société d'Anthropologie de Paris* 1984; **XIV**, pp. 265-70.
- <sup>25</sup> Millman S. Trends in breastfeeding in a dozen developing countries. *Int Fam Plann Perspect* 1986; **12**: 91-95.
- <sup>26</sup> Onuoha N. Contributions of the proximate determinants to fertility change in Senegal. *Soc Sci Med* 1992; **35**: 1317-20.
- <sup>27</sup> Lamb W H, Foord F A, Lamb C M B, Whitehead R G. Changes in maternal and child mortality rates in three isolated Gambian villages over ten years. *The Lancet* 1984; **ii**: 912-14.
- <sup>28</sup> Pison G, Trape J F, Lefebvre M, Enel C. Rapid decline in child mortality in a rural area of Senegal. *Int J Epidemiol* 1993; **22**: 72-80.

(Revised version received March 1996)