

LEAD AUTHOR

Sam Geerts
(KU Leuven University, Leuven,
Belgium), Magali Garcia
(Universidad Mayor de San
Andrés, La Paz, Bolivia)

CONTRIBUTING AUTHORS & INSTITUTIONS

Sven-Erik Jacobsen
(University of Copenhagen,
Taastrup, Denmark),

Jean Vacher
(Ambassade de France,
Lima, Peru),

Thierry Winkel
(IRD, CEFÉ-CNRS,
Montpellier, France),

Daniel Bertero
(Universidad de Buenos
Aires and CONICET, Buenos
Aires Argentina)

ACKNOWLEDGMENTS

We thank the Faculty of
Agronomy of the University
of San Andres, La Paz, Bolivia
(UMSA) for the support to the
work leading to the results
presented here.

We are also thankful for the
financial support for field work
from the VLIR-UOS,
Brussels, Belgium.

Dirk Raes
(KU Leuven University,
Leuven, Belgium)

Pasquale Steduto
(FAO, Land and Water Division,
Rome, Italy)

SCIENTISTS CONTRIBUTING WITH EXPERIMENTAL DATA AND TESTS FOR THE CALIBRATION OF AQUACROP

Edwin Yucra,
Cristal Taboada,
Roberto Miranda,
Jorge Cusicanqui,
Felix Mamani
(Universidad Mayor de San
Andrés, La Paz, Bolivia)

Quinoa

GENERAL DESCRIPTION


Quinoa (*Chenopodium quinoa* Willd.), a species of the goosefoot (*Chenopodium*) genus of the family of sugar beet, beetroot, mangold and spinach, is a grain-like crop grown primarily for its edible seeds. It is a seed crop, rather than a true cereal, or a grain, as it is not a member of the grass family. It is native to the Andean mountains where this traditional seed crop has been cultivated in the Peruvian and Bolivian Andes for more than 7 000 years. Although the production declined significantly during the Spanish conquest, popularity of quinoa rose again in the last century. Production is now widespread in the Andes, covering Bolivia, Peru, Ecuador, Colombia and the north of Argentina and Chile. As a crop with a large food utilization potential, it is rapidly gaining interest globally, being already fairly known in North and Central America, Brazil, Europe and Asia (Schulte auf'm Erley *et al.*, 2005). First results also indicate the potential of quinoa in Africa. In 2009, 83 thousand hectares were sown to quinoa, producing 69 000 tonne of grain at an average grain yield of 0.8 tonne/ha (FAO, 2011). The worldwide trend for cropping area and production over the last 50 years is shown in Figure 1.

Quinoa is characterized by an enormous intra-species variety and plasticity that allows the crop to grow under highly diverse climatic and agronomic conditions. It is well adapted to arid and semi-arid locations and grows from sea-level to high altitudes, up to Andean Altiplano at around 4 000 m above sea level, where its cultivation is of special importance. Quinoa is cultivated as a mono-culture (e.g. Southern Bolivian Altiplano) or in rotation with potato and barley, and sometimes with wheat and maize at the low altitudes. When cultivated as mono-culture, fields are left fallow for 1 to 3 years and sometimes even longer (up to 10-12 years) for pest control, soil fertility regeneration and build up of the soil water reserve. Traditionally, a range of quinoa landraces is cultivated in the same vicinity, though for export purposes a few local cultivars sown in monoculture are generally preferred (e.g. quinoa var. Real Blanca). Daylength neutral cultivars of quinoa can be grown under the long day conditions of northern Europe (Christiansen *et al.*, 2010).

GROWTH AND DEVELOPMENT

Quinoa is a C₃ annual dicot of 0.5 to 2 m height, terminating in a panicle consisting of small flowers, and with only one seed of around 2 mm

FIGURE 1 World quinoa harvested area and average yield over the period 1961-2009 (FAO, 2011).



produced per flower. In the Andean highlands, quinoa is grown from September to May, and normally without any fertilizer or pesticide. In the Bolivian Altiplano, the sowing date varies between the beginning of September and the end of November, according to the crop cycle length of the different cultivars and local climate, particularly when the soil is moist enough for germination. Sowing practice is the key for the success of quinoa. Superficial sowing runs the risk of seed dehydration or sunburn whereas deep sowing can prevent emergence; in both cases, a poor stand and uneven canopy cover occurs with detrimental effects on final yield. Common practice is to sow between 8 to 15 kg of seeds per hectare in rows 0.4 to 0.8 m apart, either on top of the bed or in the furrow, with plants spaced about 10 cm apart within the row after thinning. Another practice, adapted to arid environments where commercial production is widespread, is to group several plants each in pits spaced about 1 m apart. Less common are transplanting (Inter-Andean valleys) or broadcasting of the seeds. As nutrients are often scarce in the Altiplano, early weeding (\pm about 30 days after sowing) and thinning of excessive plants are important activities in the areas where rain is sufficient to allow for rapid plant growth. In arid areas, instead, weeding and thinning are not practised, even in commercial production.

Phenological development is highly variable among varieties. Additionally, phenology is highly flexible in response to water stress, with differences in time to maturity of as much as 30 days for the same cultivar (Geerts *et al.*, 2008c). Under no stress conditions, time from sowing to emergence is 3 to 10 days. The time to maximum canopy cover (CC_x) depends on plant density and temperature regime. Phenology is further complicated by a response to photoperiod — a short day response for duration of emergence to flowering, and for the duration of all developmental phases in some cultivars (Bertero, 2003). The time from emergence to

physiological maturity varies between 100 and 230 days, again because of wide variation among cultivars. Flowering starts between 60 to 120 days, and lasts around 20 days. Canopy senescence starts generally about one month before physiological maturity, and progresses relatively fast. It is important to note that these indicative values are given for cultivars cultivated at high elevations, and could be biased because growing conditions (temperature, fertility, water supply) are not optimal. Roots, often with numerous ramifications, can deepen to 1.80 m depth in cases of drought stress in light soils.

In the southern Andes, quinoa is harvested in April-May, mainly by pulling out or cutting the plants and leaving them in stacks on the field to dry. Harvest can take up to 1.5 months because of asynchronous flowering and ripening. In principle, realistic simulation of such production practice with *AquaCrop* would entail simulations runs for each maturity group, and then summing up the yields of all groups in proportion to their population density or land area occupied (Figure 2. Example of quinoa).

WATER USE & PRODUCTIVITY

In midseason, quinoa reaches maximum canopy cover (CC_x), shortly after first anthesis. CC_x is largely dependent on the management conditions and, to a large extent, determines quinoa transpiration. For a complete canopy cover of the ground and under non-limiting nutrient conditions, quinoa transpires at a rate similar to the reference evapotranspiration (ET_0). Seasonal ET values for quinoa with a normal season length of 150-170 days are around 500 mm under non-stressed conditions. As a C_3 crop, normalized crop water productivity (WP^*) is low,

FIGURE 2 Flower and grain colour presented in the figure are only an example. Depending on the quantity of anthocyanins, this colour varies from green-yellowish to deep purple and even black throughout quinoa cultivars.



with typical values around 10.5 g/m² under the low, natural fertility in the Bolivian Altiplano (Geerts *et al.*, 2009). Still-under poor fertility conditions, a decrease of the reference biomass water productivity (WP*) value only occurs at higher total transpiration sums, and only by 10 percent. The C₃ pathway is well adapted to the prevailing low average temperatures in the Altiplano. Reported values of seed yield per unit of water consumed (WP_{Y/ET}) are rather low and lie between 0.3 and 0.6 kg/m³ because of the generally prevailing low fertility conditions (Geerts *et al.*, 2009). On the other hand, it is a crop with a large nitrogen-sink thus causing an increased metabolic cost or higher glucose-equivalent per unit dry matter produced. To our knowledge, no research has been conducted on the response of quinoa to increases in atmospheric CO₂.

RESPONSE TO STRESSES

Quinoa is highly resistant to a number of abiotic stresses (Jacobsen *et al.*, 2003). Several drought resistant mechanisms are present in quinoa. Drought in early vegetative stages may prolong its life cycle, allowing the plant to make up for growth lost during the early drought if water is available later. Also, the availability of cultivars with different season length makes it possible to match the water requirement of the quinoa crop to the available rainfall or the stored soil water at a given location. Quinoa tissue is relatively high in osmotic solutes and undergoes substantial osmotic adjustment under drought, which enable stomata to remain somewhat open down to a leaf water potential range of -1.5 MPa. During soil drying the plants are able to maintain leaf water potential and photosynthesis due to the complex stomatal response, resulting in an increase of leaf water use efficiency. Root originated ABA plays a role in stomata performance during soil drying. ABA regulation seems to be one of the mechanisms utilized by quinoa when facing drought inducing decrease of turgor of stomata guard cells (Jacobsen *et al.*, 2009). The plant also avoids negative effects of drought through fast and deep rooting particularly in dry soils. Quinoa also reduces its leaf area by controlled leaf senescence under drought.

Quinoa is a facultative halophyte (Bosque-Sanchez *et al.*, 2003) and can grow in non-saline to extremely saline conditions, depending on the cultivar. Seed production is enhanced by moderate salinity (EC in the 5-15 dS/m range) and may not be drastically reduced even at EC of 40 to 50 dS/m in some cultivars (Jacobsen *et al.*, 2003). Osmotic adjustment by the accumulation of salt ions in tissues enables the plant to maintain cell turgor and transpiration under saline conditions.

Apart from drought, frost and cold are the other major growth limiting factors in the Altiplano. Quinoa is tolerant to frost, partly because of the protection provided by its heterogeneous canopy (Winkel *et al.*, 2009), although the tolerance varies with cultivar and appears to diminish at the late phenological stages (Jacobsen *et al.*, 2005). Leaf freezing of quinoa occurred only between -5 and -6 °C, and is delayed in case of mild water stress (Bois *et al.*, 2006). The resistance to frost is associated with super cooling of tissue water and tolerance of extracellular ice formation (in the cell wall), as is common for most winter crops.

Linked to frost resistance is a low base temperature (T_{base}) for plant processes. In a study of the leaf appearance rate of different quinoa cultivars originating from various altitudes and latitudes, T_{base} averaged 2 °C and the temperature at which maximum rate was reached averaged 22 °C. Other studies found a T_{base} of 3 °C, T_{opt} of 30-35 °C and T_{max} estimated to 50 °C

(Bertero *et al.*, 2000; Jacobsen and Bach, 1998). Temperature sensitivity of quinoa was highest in cultivars originating in cold and dry climates and lower in cultivars from warmer and humid climates (Bertero *et al.*, 2000).

Because soil fertility is generally poor in its centre of origin, cases of quinoa cultivation under non-limiting soil fertility are very rare. Research into nitrogen and phosphorus requirements conducted in Colorado, United States, found that maximum yields over 4.5 tonne per ha are possible when 170 to 200 kg N/ha were applied (Oelke *et al.*, 1992). If these results are confirmed in other studies, the WP* of quinoa under non-limiting fertility would be much higher than the value of 10.5 g/m² reported earlier for the low fertility natural conditions of the Andean Altiplano. No effect on yield was observed when 34 kg of phosphorus (as phosphoric acid) per ha was applied, in comparison to an untreated field plot (Oelke *et al.*, 1992). In areas of traditional cultivation, some sheep or lama dung is applied when available; but mostly, quinoa is sown in unfertilized fields. If sown after potato, nutrient supply is generally better because of the nutrients left over from the potato fertilization.

IRRIGATION PRACTICE

As quinoa is drought resistant, it is traditionally cultivated under rainfed conditions, even in semi-arid locations. Researchers, though, started to study the impact of additional water on quinoa production and found that deficit irrigation (DI) was highly beneficial in various experimental locations. DI is already practised for the reintroduction of quinoa in arid regions of Chile. On the other hand, currently quinoa is rarely cultivated under full irrigation, as research of quinoa under full irrigation gave only slightly better results than quinoa cultivated under deficit irrigation (Geerts *et al.*, 2008a and b), besides the issue that sufficient water for full irrigation is mostly unavailable.

YIELD

Quinoa produces nutritious seeds (Alvarez-Jubete *et al.*, 2009) with high protein content (from 12 up to 20 percent), as compared to maize, rice or even wheat. The balanced amino acids composition makes the protein quality comparable to that of milk, making it an effective meat and milk substitute. Additionally, quinoa is gluten free, which is advantageous for commercial food manufacturing for celiac consumption. On the other hand, the seeds also contain the anti-nutritional component saponin (Mastebroek *et al.*, 2000), that the plant produces as an inherent protection against pests. Saponin is removed by washing or dry polishing before consumption.

Although high yields (up to 4.5 tonne/ha and more) have been occasionally reported for some quinoa cultivars under non-limiting fertility and water conditions, rainfed yields in the Peruvian and Bolivian Altiplano do not exceed 0.85 tonne/ha as an average. Pests, including birds and rodents, and diseases are major causes of yield loss, in addition to low fertility and water deficits. Harvest index (HI) of quinoa in the field ranges between 0.3 and 0.5. Building up of HI takes a short time for the short season cultivars, and up from 80 to 100 days for the long season cultivars. Individual grain size is quite variable among cultivars, with 1 000 grain weight ranging from 1.2 to 6.0 g in non-stress conditions (Rojas, 2003).

REFERENCES

- Alvarez-Jubete, L., Arendt, E.K., Gallagher, E. 2009. Nutritive value of pseudocereals and their increasing use as functional gluten-free ingredients. *Trends in Food Science & Technology* 21: 106-113.
- Bertero, H.D. 2003. Response of Developmental Processes to Temperature and Photoperiod in Quinoa (*Chenopodium quinoa* Willd.). *Food Reviews International* 19, 87-97.
- Bertero, H.D., King, R.W. & Hall, A.J. 2000. Photoperiod and temperature effects on the rate of leaf appearance in quinoa (*Chenopodium quinoa* Willd.). *Australian Journal of Plant Physiology* 27, 349-356.
- Bois, J.F., Winkel, T., Lhomme, J.P., Raffailac, J.P. & Rocheteau, A. 2006. Response of some Andean cultivars of quinoa (*Chenopodium quinoa* Willd.) to temperature: Effects on germination, phenology, growth and freezing. *European Journal of Agronomy* 25, 299-308.
- Bosque-Sanchez, H., Lemeur, R., Van Damme, P. & Jacobsen, S.E. 2003. Ecophysiological analysis of drought and salinity stress of quinoa (*Chenopodium quinoa* Willd.). *Food Reviews International* 19: 111-119.
- Christiansen, J.L., Jacobsen, S.-E. & Jørgensen, S. T. 2010. Photoperiodic effect on flowering and seed development in quinoa (*Chenopodium quinoa* Willd.). *Acta Agriculturae Scandinavica, Section B - Plant Soil Science*, 1-6.
- FAO. 2011. FAOSTAT online database, available at link <http://faostat.fao.org/>. Accessed on December 2011.
- Geerts, S., Raes, D., Garcia, M., Vacher, J., Mamani, R., Mendoza, J., Huanca, R., Morales, B., Miranda, R., Cusicanqui, J. & Taboada, C. 2008a. Introducing deficit irrigation to stabilize yields of quinoa (*Chenopodium quinoa* Willd.). *European Journal of Agronomy* 28, 427-436.
- Geerts, S., Raes, D., Garcia, M., Condori, O., Mamani, J., Miranda, R., Cusicanqui, J., Taboada, C. & Vacher, J. 2008b. Could deficit irrigation be a sustainable practice for quinoa (*Chenopodium quinoa* Willd.) in the Southern Bolivian Altiplano? *Agricultural Water Management* 95, 909-917.
- Geerts, S., Raes, D., Garcia, M., Mendoza, J. & Huanca, R. 2008c. Indicators to quantify the flexible phenology of quinoa (*Chenopodium quinoa* Willd.) in response to drought stress. *Field Crops Research* 108, 150-156.
- Geerts, S., Raes, D., Garcia, M., Taboada, C., Miranda, R., Cusicanqui, J., Mhizha, T. & Vacher, J. 2009. Modelling the potential for closing quinoa yield gaps under varying water availability in the Bolivian Altiplano. *Agricultural Water Management* 96, 1652-1658.
- Jacobsen, S.-E. & A.P. Bach. 1998. The influence of temperature on seed germination rate in quinoa (*Chenopodium quinoa* Willd.). *Seed Science and Technology* 26, 515-523.
- Jacobsen, S.E., Mujica, A. & Jensen, C.R. 2003. The Resistance of Quinoa (*Chenopodium quinoa* Willd.) to Adverse Abiotic Factors. *Food Reviews International* 19, 99-109.
- Jacobsen, S.-E., Monteros, C., Christiansen, J.L., Bravo, L.A., Corcuera, L.J. & Mujica, A. 2005. Plant responses of quinoa (*Chenopodium quinoa* Willd.) to frost at various phenological stages. *European Journal of Agronomy* 22, 131-139.
- Jacobsen, S.-E., Liu, F. & Jensen, C.R. 2009. Does root-sourced ABA play a role for regulation of stomata under drought in quinoa (*Chenopodium quinoa* Willd.). *Scientia Horticulturae* 122, 281-287.
- Mastebroek, H.D., Limburg, H., Gilles, T. & Marvin, H.J.P. 2000. Occurrence of saponins in leaves and seeds of quinoa (*Chenopodium quinoa* Willd.). *Journal of the Science of Food and Agriculture* 80(1): 152-156.
- Oelke, E.A., Putnam, D.H., Teynor, D.M. & Oplinger, E.S. 1992. *Alternative field crops manual—quinoa*. USA, University of Purdue.
- Rojas, W. 2003. Multivariate Analysis of Genetic Diversity of Bolivian Quinoa Germplasm. *Food Reviews International* 19, 9-23.
- Schulte auf'm Erley, G., Kaul, H.-P., Kruse, M. & Aufhammer, W. 2005. Yield and nitrogen utilization efficiency of the pseudocereals amaranth, quinoa, and buckwheat under differing nitrogen fertilization. *European Journal of Agronomy* 22, 95-100.
- Winkel, T., Lhomme, J. P., Nina Laura, J.P., Mamani Alcon, C., del Castillo, C., Rocheteau, A. 2009. Assessing the protective effect of vertically heterogeneous canopies against radiative frost: The case of quinoa on the Andean Altiplano. *Agricultural and Forest Meteorology*, 149, 1759-1768.





Quinoa*

**Flower and grain colour presented in the figure are only an example. Depending on the quantity of anthocyanins, this colour varies from green-yellowish to deep purple and even black throughout quinoa varieties.*

Crop yield response to water



Crop yield response to water

FAO
IRRIGATION
AND
DRAINAGE
PAPER
66

by

Pasquale Steduto

(FAO, Land and Water Division, Rome, Italy)

Theodore C. Hsiao

(University of California, Davis, USA)

Elias Fereres

(University of Cordoba and IAS-CSIC, Cordoba, Spain)

Dirk Raes

(KU Leuven University, Leuven, Belgium)

The designations employed and the presentation of material in this information product do not imply the expression of any opinion whatsoever on the part of the Food and Agriculture Organization of the United Nations (FAO) concerning the legal or development status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. The mention of specific companies or products of manufacturers, whether or not these have been patented, does not imply that these have been endorsed or recommended by FAO in preference to others of a similar nature that are not mentioned.

The views expressed in this information product are those of the author(s) and do not necessarily reflect the views of FAO.

ISBN 978-92-5-107274-5

All rights reserved. FAO encourages reproduction and dissemination of material in this information product. Non-commercial uses will be authorized free of charge, upon request. Reproduction for resale or other commercial purposes, including educational purposes, may incur fees. Applications for permission to reproduce or disseminate FAO copyright materials, and all queries concerning rights and licences, should be addressed by e-mail to copyright@fao.org or to the Chief, Publishing Policy and Support Branch, Office of Knowledge Exchange, Research and Extension, FAO, Viale delle Terme di Caracalla, 00153 Rome, Italy.

© FAO 2012