

3D model related to the publication: A fossil terrestrial fauna from Tobène (Senegal) provides a unique early Pliocene window in western Africa

Fabrice Lihoreau^{1*}, Raphaël Sarr², Dominique Chardon³, Jean-Renaud Boisserie^{4,5}, Renaud Lebrun¹, Sylvain Adnet¹, Jérémy E. Martin⁶, Bernard Sambou², Rodolphe Tabuce¹, Moustapha Thiam⁷, Lionel Hautier¹

¹Institut des Sciences de l'Evolution, UMR5554, CNRS, IRD, EPHE, Université de Montpellier, Montpellier, France

²Département de Géologie, Faculté des Sciences et Techniques, Université Cheikh-Anta-Diop de Dakar, B. P. 5005 Dakar-Fann, Sénégal

³GET, Université de Toulouse, IRD, CNRS, UPS, CNES, Toulouse F-31400, France

⁴Paleovprim UMR 7262, CNRS, Université de Poitiers, France

⁵Centre Français des Études Éthiopiennes, USR 3137, CNRS, Ministère de l'Europe et des Affaires Étrangères, Addis Abeba, Éthiopie

⁶Univ. Lyon, ENS de Lyon, Université Claude Bernard Lyon 1, CNRS, UMR 5276 Laboratoire de Géologie de Lyon: Terre, Planètes, Environnement, F-69342 46 Allée d'Italie, Lyon, France

⁷Laboratoire de Biostratigraphie et Sédimentologie, Département de Géologie, Faculté des Sciences et Techniques, Université Cheikh-Anta-Diop de Dakar, B. P. 5005 Dakar-Fann, Sénégal.

*Corresponding author: fabrice.lihoreau@umontpellier.fr

Abstract

The present contribution contains the 3D virtual restoration of a Pliocene right femur belonging to a Lutrinae indet. (Carnivora) from Tobène, Senegal, described and figured in Lihoreau et al. (2021) : "A fossil terrestrial fauna from Tobène (Senegal) provides a unique early Pliocene window in Western Africa ". <https://doi.org/10.1016/j.gr.2021.06.013>

Keywords: Carnivora, Lutrinae, Femur, Pliocene, Senegal

Submitted:2020-06-03, published online:2021-06-24. <https://doi.org/10.18563/journal.m3.102>

Inv nr.

SN-Tob-12-02

Description

3D surface virtual restoration

Table 1. Related model of a right femur belonging to a Carnivora (Lutrinae indet). Collection: Université Cheikh-Anta-Diop, Dakar, Senegal.

INTRODUCTION

SN-Tob-12-02 is an uncomplete right femur of Lutrinae indet, preserving a fragmentary portion of the proximal part of the diaphysis, as well as a fragmentary distal extremity, including the condyles. This specimen belongs to the fauna retrieved from the Pliocene locality of Tobène, Senegal. As most specimens from this locality, SN-Tob-12-02 is fragmentary, and several parts, such as the patellar facet have been shifted post-mortem. Therefore, a virtual restoration of this specimen was performed and is presented here (see Table 1 and Fig. 1).

METHODS

SN-Tob-12-02 was digitized using X-ray microtomography (μ CT) using an EasyTom 150 microtomograph (MRI platform). The distal extremity of this femur contains four main fragments (the patellar facet, a part of the lateral condyle, the medial condyle, and a part of proximal shaft), which had been slightly broken and displaced post-mortem. 3D surfaces representing these four parts were extracted with Avizo 9.2 (Thermo Fisher Scientific). The surfaces belonging to the distal extremity and

to the proximal part of the diaphysis were positioned in space with MorphoDig software (Lebrun, 2018) in order to retrieve anatomical connection.

ACKNOWLEDGEMENTS

3D data acquisitions were performed using the μ CT facilities of the MRI platform member of the national infrastructure France-BioImaging supported by the French National Research Agency (ANR-10-INBS-04, «Investments for the future»), and of the Labex CEMEB (ANR-10-LABX-0004) and NUMEV (ANR-10-LABX-0020).

BIBLIOGRAPHY

Fabrice Lihoreau, Raphaël Sarr, Dominique Chardon, Jean-Renaud Boisserie, Renaud Lebrun, Sylvain Adnet, Jeremy Martin, Bernard Sambou, Rodolphe Tabuce, Mohamadou M. Thiam, Lionel Hautier (2021). A fossil terrestrial fauna from Tobène (Senegal) provides a unique early Pliocene window in western Africa. Gondwana Research. <https://doi.org/10.1016/j.gr.2021.06.013>




Figure 1. Distal extremity of SN-Tob-12-02 in cranial (A) caudal (B) and distal (C) orientation. Brown: proximal shaft fragment. Green: patellar facet. Yellow: medial condyle. Cyan: lateral condyle fragment.