

# Lipid Composition and State of Oxidation of Fortified Infant Flours in Low-Income Countries Are Not Optimal and Strongly Affected by the Time of Storage

Célia Moustiés, Claire Bourlieu, Bruno Barea, Adrien Servent, Pascaline Alter, Marc Lebrun, Youna Morgane Hemery, Arnaud Laillou, and Sylvie Avallone\*

Food fortification is widely used to address the public health problem of nutrient deficiencies. This study's purpose is to assess the lipid profile and nutritional quality of 13 fortified infant flours (FI) collected "in the field" in Africa and Asia after different periods of storage. The lipid content, fatty acid profiles, lipophilic vitamin content, and lipid oxidation state (peroxide values, secondary oxidation products) are determined. Mycotoxins and packaging materials are also characterized. The lipid content averages  $9.1 \pm 3.5$  g/100 g. Fatty acid profiles are dominated by linoleic ( $43.3 \pm 8.8\%$ ), oleic ( $29.5 \pm 7.4\%$ ), and palmitic acid ( $17.8 \pm 6.7\%$ ) and result in an average  $\omega 6/\omega 3$  ratio of  $12.2 \pm 5.9$  but with high values for some FI. Very high overages in vitamins A, D, and E are observed in products stored for short periods (1–6 months), whereas FI stored for more than 12 months has insufficient vitamin content. Lipid oxidation is acceptable but for six products presenting excessive peroxide values. Most products are contaminated by low amounts of mycotoxins but only two FI do not abide by the regulation. A strong correlation between peroxide values, hexanal content, and time of storage is observed.

**Practical Applications:** The expiration dates for FI commercialized in low-income countries should be shortened from 36 to 12 months so as to guarantee their nutritional quality of these functional foods and to abide by the fortified infant flour legislation. Indeed, FI quality significantly decreases over time of storage. The use of high barrier packaging materials must be generalized, although it is a necessary criterion but not sufficient to ensure the long-term stability of FI. Special attention should be given to reduce lipophilic vitamin overages and to improve their lipid profile, especially the  $\omega 6/\omega 3$  ratio, which has to be lower than 15.

## 1. Introduction

Nutrient deficiencies still affect two billion people worldwide, due to health issues, inappropriate diet, poverty, poor hygiene, and sanitation.<sup>[1]</sup> In 2016, 155 million preschool-age children were still affected by stunting.<sup>[1]</sup> In Africa and South-East Asia, the diet of low-income populations is mainly based on starchy products and do not include enough nutrient-rich foods (i.e., fruits, vegetables, beans, or animal products). These diets are lacking in essential micronutrients and in certain macronutrients, such as essential lipids, and children are particularly concerned by this nutritional issue. Lipids play different important roles, in particular energetic, structural, and transport.<sup>[2]</sup> Moreover, adequate intake of essential lipids is very important for visual acuity, brain development, and lipid metabolism.<sup>[2–5]</sup>

During the 1000 days between a woman's pregnancy and her child's 2nd birthday, adequate micronutrient intakes are essential for the child's cognitive and physical development. Indeed, during fetal and infant development, inadequate food intake strongly impacts the nutritional status, resulting in stunting with cognitive consequences and increased mortality.<sup>[6]</sup> As micronutrient deficiencies are public health

C. Moustiés, A. Servent, P. Alter, M. Lebrun, Prof. S. Avallone  
QualiSud, Univ Montpellier, CIRAD, Montpellier SupAgro, Université  
d'Avignon  
Université de la Réunion  
1101 Avenue Agropolis, Montpellier, France  
E-mail: Sylvie.avallone@supagro.fr

Dr. C. Bourlieu, B. Barea  
IATE, INRA, CIRAD, Univ Montpellier  
Montpellier SupAgro  
1101 Avenue Agropolis, Montpellier, France  
Dr. Y. M. Hemery  
Nutripass, Univ Montpellier, IRD  
Montpellier SupAgro  
911 Avenue Agropolis, Montpellier, France

Dr. A. Laillou  
UNICEF Ethiopia  
Zambezi-Building #020 2nd, 3rd & 4th floors Addis Ababa  
Addis Ababa 1169, Ethiopia

DOI: 10.1002/ejlt.201900173

issues, governments and international organizations have, for decades, implemented three main strategies to solve the problems: diet diversification, supplementation, and food fortification.<sup>[7]</sup> Diet diversification has the advantage to stimulate the local economy by generating incomes for small farmers and to promote the consumption of diversified food with complementary nutritional profiles, but in some contexts, the cost of a diversified diet seems too high and not reachable for people with low incomes.<sup>[8]</sup> Supplementation has the advantage of being very specific in the intervention and in terms of dose-adjustment, but expensive and with risk of toxicity, due to low compliance and self-prescription by users.<sup>[9]</sup> The use of food fortification is an alternative strategy which is considered as the most cost-effective to eliminate micronutrient malnutrition.<sup>[7]</sup> The biological efficacy of fortified food products has been demonstrated in clinical trials. Under controlled environments, the conditions are perfect and the results are encouraging, but they do not reflect the “real food system” in terms of conditions of storage in the supply chain of southern countries and practices at home.

Fortified infant flours (FI), mostly cereal-based, have been designed to fulfill the needs of infants and preschool children. FI are used in complement to breastfeeding from the age of 6 months, in the form of gruel or porridge, consumed on a daily basis. Most FI have adjusted lipid profiles and can be enriched in polyunsaturated fatty acids (PUFA); enrichment in essential fatty acids (FA), that is, linoleic C<sub>18:2</sub> ( $\omega$ 6 family precursor) and linolenic C<sub>18:3</sub> ( $\omega$ 3 family precursor) FA, and frequently in long-chain PUFA that are important for brain and vision development, such as docosahexaenoic acid (DHA, C<sub>22:6</sub>,  $\omega$ 3 family).<sup>[4,10]</sup> Indeed, lipids play several important functions for young children, as they supply most of the calories necessary for growth and contribute to the development of cerebral functions and to lipophilic vitamin transport.<sup>[2]</sup> More specifically, DHA contributes to visual acuity and brain membrane accumulation during the child's first 2 years of life. Unbalanced  $\omega$ 6/ $\omega$ 3 lipid input during infant growth do not support infant development as satisfactorily as balanced ones.<sup>[4]</sup>

A previous study conducted in low-income countries has reported that an adequate consumption of well-formulated fortified cereal-based complementary foods has a positive impact on the nutritional status of young children.<sup>[11]</sup> The nutritional quality of many complementary foods sold in African and South-east Asian countries have been assessed,<sup>[12–14]</sup> but these studies focused either on the macronutrients, minerals, and trace elements, or on the anti-nutrients, and none focused on lipid profile or reported on lipophilic vitamin contents. Moreover, unsaturated FA and lipophilic vitamins present poor chemical stability during food production, transportation, and storage.<sup>[15]</sup> Vitamin A can be rapidly degraded, especially in food products stored in non-optimal conditions. High losses in vitamin A have been reported in products stored at moderate temperatures ( $\approx$ 40 °C) and when the packaging does not display good oxygen and water vapor barrier properties.<sup>[16,17]</sup> Very recently, more than half of the vitamin A fortified flours used in 20 national fortification programs did not meet the fortification standards.<sup>[18]</sup> Performance data on fortification interventions are often missing and regular assessment of fortification programs in “field conditions” are needed.<sup>[19,20]</sup>

The aim of the present work was to characterize 13 FI, sold and consumed on a daily basis, in four countries in Africa and Asia. FI were collected “in the field” after different storage periods. Lipid composition (i.e., total and essential fatty acids, and lipophilic vitamins), lipid oxidation (peroxide and acid values, and volatile compounds), mycotoxin contents, and barrier properties of packaging materials were evaluated.

## 2. Experimental Section

### 2.1. Collection and Storage of FI

Thirteen FI consumed on a daily basis were collected in pharmacies, groceries, and markets in four countries in Africa and South-east Asia (Table 1). The products were sampled after being stored in “field conditions” in the real supply chain from 1 to 19 months. Storage time is thus the period of time elapsing from the manufacture to the collection (purchase). Products were then shipped to the laboratory (Montpellier, France). The products were produced either locally or imported.

In order to obtain a representative sample, several unit-doses were mixed (achieved average mass  $\approx$ 105–150 g). At opening, products were immediately analyzed for vitamin profiles, dry matter, and water activity. The rest of the samples was vacuum-packed in distinct bags and frozen at  $-80$  °C for further analyses (mycotoxins, total and neutral lipid extraction and characterization, and volatile compounds).

Each product was labeled with a code FI\_X, where X corresponds to a specific brand. The characteristics of the products in terms of formulation, packaging, and duration of storage since production date are reported in Table 1.

For two FI brands (FI\_2 and FI\_3), two distinct batches were characterized. These four products were tested rapidly at reception (i.e., after 1 or 2 months “in field” storage) or stored for two more months in the laboratory at ambient temperature ( $\approx$ 20 °C) and again characterized. For these two brands, the products were labeled as follows: FI\_X\_LY\_Z to indicate that the FI was produced by a brand (X) with a number of batch (LY) and a specific duration of storage in month (Z). Products that were stored for two more months under controlled conditions in the laboratory were noted with an asterisk (\*): FI\_2\_L2\_3\*, FI\_2\_L1\_4\*, FI\_3\_L2\_3\*, FI\_3\_L1\_4\*.

In total 13 brands of FI were purchased in local groceries and markets in Vietnam, Burkina Faso, Madagascar, and Benin. Their production place, time of storage, packaging, and composition are detailed in Table 1. The collected products can be considered as representative of the ratio (imported/local) of local products in the supply chain. This ratio, imported/local, was of 3/13. The three imported products had all been produced in the European Union.

FI were formulated with five to 13 ingredients and limited amounts of additives (except in FI\_6). FI were based on cereals, either local ones such as millet (*Pennisetum glaucum*), sorghum (*Sorghum bicolor*) in African FI, but also wheat (*Triticum*) in three products. In most FI, lipids were mainly brought by peanuts (*Arachis hypogaea*), some oleoproteaginous such as soy (*Glycine max*) (present in 8 FI) or sesame (*Sesamum indicum*) and milk. 38% of FI included added “fats” with no other precision. 85%

**Table 1.** Characteristics of the FI.

Code	Purchase place (production place) <sup>a)</sup>	Storage duration (months)/expiration date (months)	Packaging material	Ingredients	Producer typology
FI_1	Burkina Faso	5/6	PE <sup>b)</sup>	Mil, soy, peanut, sugar, iodized salt, industrial amylase, CMV <sup>c)</sup>	Local (national producer)
FI_2	Madagascar	1-2-3-4/12	Alu/PE <sup>d)</sup>	Corn, rice, soy, peanut, sugar, salt, vitamins, minerals	Local (national producer)
FI_3	Madagascar	1-2-3-4/12	Alu/PE <sup>d)</sup>	Corn, rice, soya, peanut, sugar, salt, vitamins, minerals	Local (national producer)
FI_4	Benin	6/24	PE <sup>b)</sup>	Cereal sprouts, soy germs, fats, sugar, CMV <sup>c)</sup>	Local (national producer)
FI_5	Benin	9/24	PE <sup>b)</sup>	Corn, mil, tubers, soy, sugar, fats, full milk, carrots, vanilla, CMV <sup>c)</sup>	Local (national producer)
FI_6	Benin (Ghana)	12/12	Alu/PE <sup>d)</sup>	Wheat flour, skimmed milk, sugar, palm olein, calcium carbonate, maltodextrin, vitamins, ferrous fumarate, vanillin, zinc sulfate, bifidus bacteria, sodium chloride, potassium iodide	Local (multi-national producer)
FI_7	Benin	2/6	PE/PP <sup>e)</sup>	Rice, corn, milk, sugar, CMV <sup>c)</sup>	Local (national producer)
FI_8	Benin	3/6	PE <sup>b)</sup>	Corn, sorghum, soy, sugar, CMV <sup>c)</sup>	Local (national producer)
FI_9	Benin	4/24	PE <sup>b)</sup>	Cereals, soy, tubers, fats, sugar, CMV <sup>c)</sup>	Local (national producer)
FI_10	Benin (France)	13/18	C/Epox <sup>f)</sup>	Flours (wheat, rice, corn), sugar, CMV <sup>c)</sup>	Imported (multi-national producer)
FI_11	Benin (France)	17/36	Alu/PE <sup>d)</sup>	Corn starch, rice flour, maltodextrins, tapioca, sunflower lecithin, CMV <sup>c)</sup>	Imported (multi-national producer)
FI_12	Benin (France)	19/36	Alu/PE <sup>d)</sup>	Flours (wheat, corn, rice), maltodextrins, sugar, caramel, vanillin, sunflower lecithin, CMV <sup>c)</sup>	Imported (multi-national producer)
FI_13	Vietnam	7/12	Alu/PE <sup>d)</sup>	Sticky rice, soy, green bean, sesame, whole milk powder, vegetal fat, CMV <sup>c)</sup> , amylase, yeast, vanillin, salt	Local (national producer)

<sup>a)</sup> The place of production is indicated when different from the place of purchase; <sup>b)</sup> PE, polyethylene; <sup>c)</sup> CMV, mineral and vitamin complex; <sup>d)</sup> Alu/PE, aluminum/polyethylene; <sup>e)</sup> PE/PP, polyethylene/polypropylene; <sup>f)</sup> C/Epox, cellophane/epoxy resin.

of the formulas included sugar and 23% added salt. Lipophilic vitamins were brought under the form of mineral and vitamin complex (CMV).

## 2.2. Materials and Chemicals

Solvents (ethanol, diethyl ether, hexane, chloroform, methanol, acetone, and *tert*-butyl ethyl ether) and reagents (potassium hydroxide, phenolphthalein, methylate sodium, hydrochloric methanol, potassium iodide, thiourea, and sodium thiosulfate) were obtained from Sigma-Aldrich (Saint Quentin Fallavier, France). Fatty acid methyl esters (FAMES) external calibration standards (Mixture ME 100) were obtained from Larodan (Sweden). HPLC standards ( $\alpha$ -tocopherol,  $\delta$ -tocopherol,  $\gamma$ -tocopherol,  $\beta$ -tocopherol, retinol palmitate, retinol acetate, and vitamin D<sub>2</sub> and D<sub>3</sub>) were obtained from Sigma-Aldrich (Saint Quentin Fallavier, France) and Biopharm Rhône Ltd (Glasgow, UK). The polytetrafluoroethylene (PTFE) membranes were obtained from Sartorius (Palaiseau, France). The water used in all the experiments was mQ water (HPLC grade).

## 2.3. Dry Matter and Water Activity Determination

The dry matter (DM) contents were determined in triplicate by drying 3 g of the samples at 105 °C in Memmert heat chamber (model 100 Schwabach, Germany) until constant weight follow-

ing method 44-15A.<sup>[21]</sup> Samples and their dehydrated residues were precisely weighted on a Mettler AE166 balance (Viroflay, France). Water activity ( $a_w$ ) was determined at 25 °C on approximately 1 g with a water activity meter 4TE AQUALAB (Hopkins, USA).

## 2.4. Determination of Vitamin A and D

The vitamin profiles were determined by liquid chromatography and specific detection by DAD (vitamin A and D), as previously described.<sup>[22]</sup> The UV detection allowed to quantify two forms of vitamin A (acetate and palmitate) and two forms of vitamin D (cholecalciferol and ergocalciferol).

Flour, 1 g, was mixed with 20 mL water and gently shaken for 10 min.<sup>[23]</sup> Samples were homogenized in 16 mL of ethanol/hexane (4:3, v/v) for 30 s at 30 000 rpm (Ultra-Turrax, Prolabo, France) and then centrifuged (30 min, 4 °C, 13 000 × g) (Heraeus Multifuge X1R, Thermo Fisher Scientific, Villebon sur Yvette, France). Afterward, the upper layer was transferred into amber glass tubes and dried with a Genevac for 25 min at 30 °C (EZ-2 series, Genevac LTD, Sp Scientific, England). The dried residues were dissolved in 2 mL of acetone, filtered with 0.45 µm PTFE minisart SRP4 membrane (Sartorius) and analyzed by liquid chromatography (Agilent System 1200 series, Massy, France). The column was a polymeric C<sub>30</sub> (4.6 mm i.d. × 250 mm, 5 µm particle size; YMC, Inc., Wilmington, NC) and the injection volume was 20 µL. The mobile phase was based on the blend of

two mixes at a flow rate of 1 mL min<sup>-1</sup> and the temperature was 25 °C. Mix A was constituted of methanol and water, 60:40 v/v, and mix B was constituted of methanol, methyl *tert*-butyl-ether and water, 28.5:67.5:4.0 v/v/v. The gradient applied progressed from 100% to 0% (A/B) over a period of 43 min and the column was washed for 5 min with ethyl acetate at the end of each run. Chromatograms were recorded with a UV-visible photodiode array detector (Agilent Technologies 1200 series) at the wavelength of maximum absorption of the vitamin A and D in the mobile phase (i.e., 325 and 265 nm, respectively). External calibration was realized weekly with standard solutions of the pure chemical in acetone in the range of 0.5 to 25 mg L<sup>-1</sup>. Each sample was analyzed with five repetitions. Vitamin A activity was expressed in retinol equivalents (RE).<sup>[24]</sup>

$$\begin{aligned} 1 \mu\text{g retinol equivalent} &= 1.146 \mu\text{g retinol acetate} \\ &= 1.832 \mu\text{g retinol palmitate} \end{aligned}$$

## 2.5. Determination of Vitamin E

Vitamin E was determined on the same extract that had been obtained for vitamin A and D quantification. Fluorescence was used to detect and quantify four distinct tocopherols ( $\alpha$ ,  $\beta$ ,  $\delta$ , and  $\gamma$ ). HPLC analysis was performed with an Agilent System 1290 series (Massy, France) equipped with a C<sub>18</sub> column (250 mm × 4.6 mm i.d., 5  $\mu$ m, HALO-5 column; AMT, Wilmington, Delaware, USA) and a fluorescence detector. The mobile phase consisted of ethanol/methanol (40:60, v/v) in isocratic conditions. The temperature of the column was maintained at 25 °C and flow rate was 0.8 mL min<sup>-1</sup>. Fluorescence detection was set at 296 nm for excitation and 330 nm for emission. The injection volume was 20  $\mu$ L and the calibration curves were realized with standard solutions from 0.1 to 1.5 mg L<sup>-1</sup>. Each sample was analyzed with five repetitions.

Vitamin E activity is expressed in  $\alpha$ -tocopherol equivalent ( $\alpha$ -TE) in mg for 100 g<sup>[25]</sup> and calculated as follows:

$$\alpha - \text{TE} = (1 \times \alpha - \text{T}) + (0.7 \times \alpha - \text{TA}) + (0.4 \times \beta - \text{T}) + (0.2 \times \gamma - \text{T}) + (0.1 \times \delta - \text{T}) \quad (1)$$

where  $\alpha$ -TE: tocopherol equivalent, T: tocopherol, TA: tocopherol acetate,  $\alpha$ :  $\alpha$ -tocopherol,  $\beta$ :  $\beta$ -tocopherol,  $\gamma$ :  $\gamma$ -tocopherol, and  $\delta$ :  $\delta$ -tocopherol.

## 2.6. Determination of Lipid Content and Fatty Acid Composition

The lipid content was determined gravimetrically in triplicate<sup>[26]</sup> and each extract was then analyzed in duplicate. Briefly, lipids were extracted from thoroughly homogenized food samples using chloroform/methanol (2:1, v/v). Fifteen grams of sample was rehydrated for 10 min in distilled water and, thereafter, dispersed in a sample/solvent ratio of 1:37 w/v for 1 min at 10 000 rpm (Ultra-Turrax T8, IKA Labortechnik, Staufen, Germany). NaCl 0.73% w/v was added to the organic phase with thorough shaking. After dephasing, the lower phase was filtered and retrieved.

Then, the sample was washed again with NaCl 0.58% w/v. Once clear, the organic layer was filtered and collected. Solvent was evaporated with a rotavapor at 40 °C until constant weight and residual lipids were precisely weighted (Laborota 400-efficient, Heidolph, Schwabach, Germany).

FAMES were obtained by methylation of lipids ( $\approx$ 200 mg) with 3 mL of sodium methylate for 10 min at 65 °C in heated mantle with saponification sticks under reflux. Then, 3 mL of chloride methanol was added and samples were heated again for 10 min at 64.7 °C. FAMES were extracted with 8 mL of hexane and washed with 10 mL of distilled water. 5  $\mu$ L of samples were analyzed by gas chromatography (Focus GC, Thermo Electron Corporation, Massachusetts, USA) equipped with a split injector (ratio of 1/20), a CP-Cil 88 Varian capillary column (50 m × 0.25 mm with 0.2  $\mu$ m film thickness; Chrompack, Middelburg, The Netherlands), and helium 1 mL min<sup>-1</sup> as the carrier gas. FAMES were analyzed by flame ionization detector and ChromCard software data system (version 2005; Thermo Fisher Scientific, Massachusetts, USA).

The column temperature started from 150 °C, then reached 150 to 225 °C with a rise of 5 °C min<sup>-1</sup> and was kept at 225 °C for 10 min. These chromatographic conditions allowed a correct and rapid separation of all FAMES from C<sub>10:0</sub> to C<sub>24:1</sub>. FAMES were identified using as external standards a mixture of methyl esters (Mixture ME 100, Larodan, Sweden). The injector and detector temperatures were 250 and 270 °C, respectively. FA contents were expressed in g per 100 g of fresh product, taking into account the total lipid content determined by Folch extraction.

## 2.7. Peroxide Value Determination

The peroxide values (PV) were determined by iodometric titration<sup>[27]</sup> with slight modifications. For each FI, average values were calculated from six determinations (three extractions analyzed in duplicate). Approximately 200 mg of oil was dissolved in 1 mL of chloroform. After addition of 1.5 mL of acetic acid and 0.1 mL of saturated potassium iodide, the solution was shaken for 1 min and placed for 5 min in the dark. After addition of 7.5 mL of distilled water and iodine indicator, the solution was titrated with Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> 0.002 N until the color disappeared. A blank determination was made, and the PV was calculated using the following equation:

$$\text{PV} = (V_s - V_b) \times N \times \frac{100}{W} \quad (2)$$

where PV: mEqO<sub>2</sub>/kg of lipid, V<sub>s</sub>: volume of Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> used in sample (mL), V<sub>b</sub>: volume of Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> used in the blank sample (mL), N: normality of Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> (mEq mL<sup>-1</sup>), and W: weight of lipids (g).

## 2.8. Acid Value Determination

The acid values (AV) were determined by titrimetry<sup>[28]</sup> and average values were obtained from six determinations (three extractions analyzed in duplicate). A precise quantity of lipid ( $\approx$ 200 mg) was dissolved in 0.4 mL ethanol/diethyl oxide (50:50, v/v). After



addition of phenolphthalein, the solution was titrated with KOH 0.01 N until the pink color persisted for at least 10 s. A blank determination was made, and the AV was expressed in oleic acid equivalent and calculated using the following equation:

$$AV = (V_s - V_b) \times N \times \frac{282.46}{10 \times W} \quad (3)$$

where AV: % lipid (acid oleic eq%),  $V_s$ : volume of KOH used in sample (mL),  $V_b$ : volume of KOH used in the blank sample (mL), N: normality of KOH, 282.46: molecular weight of oleic acid (g/mol), and W: weight of lipids (g).

## 2.9. Volatile Compound Determination

For each sample, the volatile compounds were extracted in triplicate using headspace solid-phase microextraction gas chromatography.<sup>[29]</sup> Extraction and analysis were carried out using divinylbenzene/carboxen/polydimethylsiloxane fiber (Supelco, Bellefonte, Pennsylvania, USA), and a tandem gas chromatograph 6890/MSD 5973 (Agilent Technologies, Palo Alto, USA)/Gerstel Multipurpose Sample MPS-2. Sample (1 g) was placed in headspace vials (10 mL) with the addition of benzyl butyrate (5  $\mu$ L/100 mL) as internal standard for semi quantification. Samples were left to equilibrate for 15 min and trapped for 60 min at 60 °C. DB-Wax UI (J&W Scientific Folsom, CA, USA) fused silica capillary column (30 m  $\times$  0.25 mm i.d.  $\times$  0.25  $\mu$ m film) was used with a 1.2 mL min<sup>-1</sup> constant flow of hydrogen. The temperature program for elution was: 3 °C min<sup>-1</sup> from 40 to 170 °C, then 10 °C min<sup>-1</sup> up to 240 °C and held for 10 min. The injection was performed in splitless mode injector at 250 °C. Mass spectrum was recorded in EI+ mode at 70 eV within 40 to 350 Da. Analyzer and source temperatures were 150 and 250 °C, respectively.

Data were analyzed with MassHunter version B. 06.00 (Agilent Technologies, Palo Alto, USA). Peak identification was realized comparing mass spectra with those of the National Institute of Standards and Technology (NIST) 14 database. Co-injection of alkanes from C<sub>8</sub> to C<sub>20</sub> (Sigma-Aldrich) was used for Kovats retention indice calculations and their comparison with the literature (flavornet and pherobase websites and NIST). The contents of the volatile compounds were expressed in mg equivalent of internal standard (i.e., benzyl butyrate), as previously described.<sup>[30]</sup>

## 2.10. Determination of Mycotoxins

Five grams of sample were homogenized with 20 mL of acetonitrile/water/acetic acid (50:50:1, v/v/v). After 10 min of gentle stirring, salts were added as follows: MgSO<sub>4</sub> (4 g), NaCl (1 g), sodium citrate tribasic dehydrate (1 g), and sodium citrate dibasic sesquihydrate (0.5 g). Samples were again stirred for 10 min and centrifuged (3000 rpm, 10 min). A mix of internal standard was added to 200  $\mu$ L of supernatant, and the overall extract was evaporated under nitrogen at 40 °C. Residue was diluted with 1 mL water/acetic acid 0.5%. The extract was filtered with 0.45  $\mu$ m PTFE minisart SRP4 membrane (Sartorius). The extracts were analyzed by ultra-high performance liquid chromatography<sup>[31]</sup>

and injection volume was 20  $\mu$ L. The pump model was a LC 20 AD Shimadzu (Kyoto, Japan) with an autosampler model: SIL 20 A XR Shimadzu and a column Kinetex 2.6  $\mu$ m XB-C18 100A (50 mm  $\times$  2.1 mm, Phenomenex Inc., California, USA). The oven temperature was 50 °C. The mobile phase A was water/acetic acid 0.5%, and mix B was isopropanol/acetic acid 0.5% at a flow rate of 0.4 mL min<sup>-1</sup>. The elution gradient started from 10% of eluent B, to 55% in 1.5 min, then 85% at 2 min, and decreased to 80% in 0.5 min and a stabilization step of 2% for 7 min. The tandem mass spectrometry conditions with system LCMS 8040 Shimadzu (Kyoto, Japan) were ionization ESI (positive and negative), MRM mode, ion source temperature (for desolvation line) set at 250 and 400 °C for heater block, gas nebulization at 3 L min<sup>-1</sup>, and gases drying at 15 L min<sup>-1</sup>.

Standard calibration curves were established with several mycotoxins (aflatoxins, ochratoxins, fumonisins, trichotecenes, zearalenon, and beauvericins) (Biopharm Rhône Ltd, Glasgow, UK). The contents were expressed in  $\mu$ g per 100 g of FI and analyses were performed in triplicate.

## 2.11. Packaging Characterization

Packaging was characterized by Fourier transform infrared spectroscopy (Table 1). Spectra were recorded using a Nexus 5700 spectrometer (Thermo Electron Corp., Massachusetts, USA) equipped with the diamond ATR Smart Orbit Accessory inserted. Spectral data were accumulated from 128 scans with a resolution of 4 cm<sup>-1</sup> in the range 800–4000 cm<sup>-1</sup>. All spectra pretreatments were performed using Omnic v7.3 and TQ Analyst v7.3 softwares (Thermo Electron Corp., Massachusetts, USA). Oxygen and water permeabilities were obtained from the literature for standard polymers.<sup>[32]</sup> Oxygen transmission rate (OTR) represents the quantity of O<sub>2</sub> which goes through the packaging per day and water vapor transmission rate (WVTR) represents the quantity of H<sub>2</sub>O which diffuses through the packaging per day. OTR and WVTR were calculated according to the packaging surface, the permeability coefficient of the material, and the time of storage of the product.<sup>[32]</sup>

## 2.12. Statistical Analysis

If not stated otherwise, all characterizations were conducted at least in triplicate. Analysis of variance (one-way ANOVA), followed by LSD post hoc tests were performed using Statgraphics plus 5.1 (Virginia, USA) to identify significant differences between samples. Differences were considered statistically significant for  $p$  value < 0.05. Still using the same software, correlations between the biochemical composition, time of storages, oxidation markers, and oxygen and water transfer rates were evaluated using principal component analysis (PCA).

## 2.13. Permission Statements

This work does not contain experiments using animals and does not contain human studies.

**Table 2.** Water content,  $a_w$ , lipid contents, and fatty acid profiles of the FI.

Sample	$a_w^*$	Water [g/100 g] <sup>*</sup>	Lipids [g/100 g] <sup>*</sup>	SFA [%] <sup>§</sup>	MUFA [%] <sup>§</sup>	PUFA [%] <sup>§</sup>	$\omega 6/\omega 3^§$
FI_1	0.22 ± 0.00	3.91 ± 0.16	13.6 ± 0.4 <sup>f</sup>	19.8 ± 0.1 <sup>bc</sup>	34.1 ± 0.1 <sup>ef</sup>	46.2 ± 0.2 <sup>d</sup>	8.5 ± 1.5 <sup>a</sup>
FI_2_L2_1	0.35 ± 0.00	5.72 ± 0.27	11.7 ± 3.5 <sup>def</sup>	16.7 ± 0.2 <sup>a</sup>	32.6 ± 0.5 <sup>def</sup>	50.7 ± 0.2 <sup>g</sup>	8.2 ± 0.5 <sup>a</sup>
FI_2_L1_2	0.33 ± 0.01	4.98 ± 0.09	8.7 ± 0.2 <sup>def</sup>	17.7 ± 0.2 <sup>a</sup>	33.1 ± 0.3 <sup>def</sup>	49.3 ± 0.6 <sup>g</sup>	8.2 ± 0.6 <sup>a</sup>
FI_2_L2_3*	0.38 ± 0.00	5.95 ± 0.10	9.9 ± 0.9 <sup>def</sup>	17.5 ± 0.1 <sup>a</sup>	32.4 ± 0.4 <sup>def</sup>	50.3 ± 0.3 <sup>g</sup>	8.8 ± 4.8 <sup>a</sup>
FI_2_L1_4*	0.33 ± 0.00	5.92 ± 0.25	10.0 ± 1.2 <sup>def</sup>	17.4 ± 0.1 <sup>a</sup>	33.8 ± 0.6 <sup>def</sup>	49.6 ± 0.1 <sup>g</sup>	9.1 ± 1.0 <sup>a</sup>
FI_3_L2_1	0.32 ± 0.00	5.09 ± 0.19	15.1 ± 3.2 <sup>ef</sup>	17.5 ± 0.2 <sup>a</sup>	33.8 ± 1.1 <sup>f</sup>	48.8 ± 0.7 <sup>ef</sup>	8.3 ± 1.0 <sup>a</sup>
FI_3_L1_2	0.33 ± 0.00	5.44 ± 0.67	10.6 ± 1.3 <sup>ef</sup>	15.7 ± 0.1 <sup>a</sup>	40.3 ± 0.5 <sup>f</sup>	48.5 ± 0.2 <sup>ef</sup>	8.2 ± 0.6 <sup>a</sup>
FI_3_L2_3*	0.31 ± 0.00	5.66 ± 0.29	10.3 ± 0.4 <sup>ef</sup>	17.5 ± 0.1 <sup>a</sup>	33.7 ± 0.1 <sup>f</sup>	48.8 ± 0.2 <sup>ef</sup>	9.2 ± 1.5 <sup>a</sup>
FI_3_L1_4*	0.35 ± 0.00	4.89 ± 0.16	9.8 ± 0.4 <sup>ef</sup>	17.8 ± 0.1 <sup>a</sup>	34.6 ± 0.7 <sup>f</sup>	48.2 ± 0.1 <sup>ef</sup>	9.1 ± 1.3 <sup>a</sup>
FI_4	0.18 ± 0.01	3.20 ± 0.12	8.8 ± 1.5 <sup>cdef</sup>	18.3 ± 0.3 <sup>abc</sup>	30.1 ± 0.7 <sup>cd</sup>	51.7 ± 0.3 <sup>gh</sup>	10.0 ± 1.1 <sup>bc</sup>
FI_5	0.15 ± 0.01	1.99 ± 0.13	9.0 ± 0.3 <sup>cdef</sup>	20.4 ± 2.5 <sup>c</sup>	29.6 ± 2.1 <sup>c</sup>	50.2 ± 1.0 <sup>g</sup>	11.6 ± 4.0 <sup>d</sup>
FI_6	0.20 ± 0.00	2.96 ± 0.13	10.2 ± 0.4 <sup>cdef</sup>	40.9 ± 0.5 <sup>h</sup>	42.8 ± 0.9 <sup>g</sup>	18.7 ± 0.8 <sup>a</sup>	21.2 ± 3.8 <sup>h</sup>
FI_7	0.40 ± 0.01	6.37 ± 0.08	8.8 ± 3.5 <sup>cdef</sup>	23.6 ± 0.3 <sup>d</sup>	34.7 ± 0.3 <sup>f</sup>	41.7 ± 0.4 <sup>c</sup>	31.7 ± 1.0 <sup>i</sup>
FI_8	0.40 ± 0.00	6.45 ± 0.30	6.6 ± 0.7 <sup>abcd</sup>	17.7 ± 0.1 <sup>ab</sup>	29.5 ± 0.3 <sup>c</sup>	52.8 ± 0.2 <sup>h</sup>	13.5 ± 5.9 <sup>e</sup>
FI_9	0.16 ± 0.01	3.07 ± 0.16	7.7 ± 0.2 <sup>bcd</sup>	18.1 ± 0.1 <sup>ab</sup>	28.8 ± 0.0 <sup>c</sup>	53.2 ± 0.1 <sup>h</sup>	10.7 ± 1.8 <sup>cd</sup>
FI_10	0.40 ± 0.01	5.49 ± 0.02	5.0 ± 1.4 <sup>abc</sup>	25.7 ± 0.4 <sup>de</sup>	14.1 ± 0.2 <sup>a</sup>	60.6 ± 0.6 <sup>i</sup>	14.8 ± 5.4 <sup>f</sup>
FI_11	0.28 ± 0.00	4.46 ± 0.52	1.5 ± 0.4 <sup>a</sup>	32.3 ± 0.4 <sup>f</sup>	20.4 ± 0.2 <sup>b</sup>	47.4 ± 0.7 <sup>de</sup>	16.4 ± 5.9 <sup>g</sup>
FI_12	0.41 ± 0.01	5.75 ± 0.14	2.5 ± 1.1 <sup>ab</sup>	27.1 ± 2.9 <sup>e</sup>	13.6 ± 9.3 <sup>a</sup>	59.3 ± 6.5 <sup>i</sup>	14.9 ± 0.4 <sup>f</sup>
FI_13	0.06 ± 0.01	1.78 ± 0.02	13.5 ± 0.7 <sup>f</sup>	36.3 ± 0.3 <sup>g</sup>	30.6 ± 0.6 <sup>cde</sup>	33.0 ± 0.3 <sup>b</sup>	8.9 ± 3.5 <sup>ab</sup>
Average	0.29 ± 0.1	4.69 ± 1.44	9.1 ± 3.6	22.0 ± 7.3	30.6 ± 7.5	47.8 ± 9.1	12.2 ± 5.9
Min	0.06	1.78	1.5	15.7	13.6	18.7	8.2
Max	0.41	6.45	15.1	40.9	42.8	60.6	31.7

Number of repetitions ( $n$ ),  $^*n = 3$ ,  $^§n = 6$ .  $a_w$ , water activity; SFA, saturated fatty acid; MUFA, monounsaturated fatty acid; PUFA, polyunsaturated fatty acid. Data are presented as mean ± standard deviation for 100 g of product. Values with distinct lowercase superscript letters in the same column are significantly different ( $p \leq 0.05$ ).

### 3. Results

#### 3.1. Dry Matter and Water Activity

The water contents of the FI ranged from  $1.78 \pm 0.02\%$  to  $6.45 \pm 0.30\%$  and  $a_w$  from  $0.06 \pm 0.01$  to  $0.41 \pm 0.01$  (Table 2).

#### 3.2. Lipid Content and Fatty Acid Profiles

A high heterogeneity was observed among FI in terms of lipid contents which averaged  $9.1 \pm 3.5$  g/100 g but ranged from 1.5 to 15.1 g/100 g (Table 2). The FI produced in Europe had a lower lipid content in comparison to the locally produced FI (Average  $2.9 \pm 1.8$  vs  $10.3 \pm 2.6$  g/100 g, respectively). European Union directive<sup>[33]</sup> requires a maximal lipid content of 3.3 g per 100 Kcal. For the imported products, the highest lipid content was  $5.0 \pm 1.4$  g/100 g corresponding to  $1.3 \pm 0.4$  g/100 Kcal (caloric equivalent: 1 g of lipid = 9 Kcal). Despite this heterogeneity, all the FI were in agreement with the European legislation for the maximal lipid contents.


The FA profiles in the FI were characterized by GC (Figure 1 and Table 2). The main FA were, by decreasing order, linoleic acid  $C_{18:2}$   $n-6$ , oleic acid  $C_{18:1}$   $n-9$ , and palmitic acid  $C_{16:0}$ . Their average percentages were, respectively,  $43.3 \pm 8.8\%$ ,  $29.5 \pm 7.4\%$ , and  $17.8 \pm 6.7\%$  of total lipids. FI contained on average  $22.0 \pm 7.3\%$  saturated FA (SFA) with important discrepancies between FI

since the lowest value for SFA was 15.7% (FI\_3) and highest reached 40.9% (FI\_6). Most FI contained a high amount of PUFA ( $47.8 \pm 9.1\%$ ), again with discrepancies (min = 18.7%, max = 60.6%). The  $\omega 6/\omega 3$  ratios of FI averaged  $12.2 \pm 5.9$  and were globally in agreement with the international recommendations, which indicates an optimal ratio range between 5 and 15.<sup>[3]</sup> However, it is important to underline that for two FI (FI\_6 and FI\_7), the ratios exceeded these recommendations largely ( $21.2 \pm 3.8$  and  $31.7 \pm 1.0$ , respectively).

Two FI (FI\_2 and FI\_3) were analyzed after being stored over 2 months in the laboratory. The main fatty acids (i.e.,  $C_{16:0}$ ,  $C_{18:1}$   $n-9$ ,  $C_{18:2}$   $n-6$ ) did not evolve significantly with the time of storage over this limited period of time (Table S1, Supporting Information).

#### 3.3. Lipophilic Vitamin Profiles

The lipophilic vitamin contents in FI are displayed according to their time of storage (Figure 2). Most of the FI were not consistent with the European directive.<sup>[33]</sup> For all lipophilic vitamins, a significant “time of storage effect” was observed whatever the FI brand or formulation: vitamin contents decreased with time of storage ( $p < 0.001$ ). Globally, the use of vitamin overages, that led to contents above the upper recommended limits, was far more frequent than insufficient contents. Whatever the vitamin, 13 out of 19 analyzed FI were above the upper limits and the excess reached up to 15, 35, and 10 times the



**Figure 1.** Main fatty acids of the FI. The products are classified according to increasing time of storage from left to right ( $n = 6$ ).

recommended upper limits for vitamin A, D, and E, respectively. Average amount of vitamin A for FI stored from 1 to 6 months reached  $378.8 \pm 88.0 \mu\text{g}/100 \text{ Kcal}$  corresponding to an overage of 210% compared to the authorized upper limit and dropped down to an average value of  $100.9 \pm 116.6 \mu\text{g}/100 \text{ Kcal}$  for FI stored for more than 6 months (six products). Similarly, values were quite homogeneous with important overages up to 5-month-storage, for vitamin D (average value =  $13.5 \pm 5.2 \mu\text{g}/100 \text{ Kcal}$  corresponding to an overage of 450% compared to the authorized upper limit) and for vitamin E (average value =  $4.3 \pm 2.7 \mu\text{g}/100 \text{ Kcal}$  corresponding to an overage of 143% compared to the authorized upper limit). After 5 months, averaged values dropped down to  $3.5 \pm 6.3 \mu\text{g}/100 \text{ Kcal}$  and  $1.4 \pm 1.5 \mu\text{g}/100 \text{ Kcal}$ , for vitamin D and E, respectively.

Seven out of 19 analyzed products were below the lower limits and most of them were close to the expiry date (time of storage >12 months): three FI for vitamin A, six for vitamin D, and four for vitamin E. All the imported products, i.e., FI produced in Northern countries and shipped to Africa (FI\_10, FI\_11, FI\_12) were included in this set of seven products and had an insufficient vitamin content.

When assessing compliance with the nutritional composition indicated on product label, we observed that vitamin contents were not always detailed. Moreover, the products were most of the time not compliant with the contents indicated on the label. To be more precise, for vitamin A, six products out of 13 had a content indicated on the label and two products only were compliant with indications. Regarding vitamin D, four products out of 13 had a content indicated on the label and none was compliant with indications. Regarding vitamin E, three products out of 13 had a content indicated on the label and one product only was compliant with indications. When comparing the two products stored over 2 months in the laboratory at ambient temperature ( $\approx 20^\circ\text{C}$ ), no significant difference was observed for any of the vitamins. A short time of storage at moderate temperature in the first months after production did not alter the vitamin profile of these fortified foods. However, it also is important to note that these products were packed with multilayer aluminum.<sup>[34]</sup>

### 3.4. Acid Values and Peroxide Values

The acidity values of all the products were acceptable (average value =  $0.06 \pm 0.04\%$ , range =  $[0.01; 0.16]\%$ ). Indeed, the values were below the maximum limit of 0.6% of total lipids as defined officially<sup>[35]</sup> (Figure 3). Regarding peroxide contents, five products did not meet the international recommendations, that is, were above the limit of  $10 \text{ meqO}_2 \text{ kg}^{-1}$  of lipids.<sup>[35,36]</sup> These products were those which had been stored for the longest period (i.e., 9 months and more). Peroxide contents increased significantly over the time of storage ( $p < 0.001$ ).


When comparing the two products stored over 2 months in the laboratory (FI\_2 and FI\_3), the time of storage significantly increased the acid and peroxide values ( $p < 0.001$ ), despite multilayer aluminum packaging.

### 3.5. Volatile Compounds

The main volatile oxidation compounds identified were saturated aldehydes and, by decreasing order, hexanal, pentanal, and nonanal (Table 3). These three volatile compounds significantly increased over the time of storage of FI ( $p < 0.001$ ). The highest content was reached with hexanal in the product stored the longest time ( $1.35\text{e}^{-08} \pm 2.19\text{e}^{-09} \text{ mg eq benzyl butyrate}/100 \text{ g}$  in FI\_12 after a 19-month-storage). When comparing the two products stored over 2 months in the laboratory (FI\_2, FI\_3), an increase in hexanal was observed.


### 3.6. Determination of Mycotoxins

In our experiments, trichotecene, zearalenone, ochratoxin A, and beauvericin were not detected. Low levels of fumonisins (FB1 and FB2) and aflatoxin were detected in 14 FI and only 4 FI were free of mycotoxins (Table 3). Samples were considered contaminated if the mycotoxin levels exceeded the EU limit,<sup>[37]</sup> which



**Figure 2.** Content in vitamins of the FI according to the time of storage: different letters above each bars indicate significant difference ( $p < 0.05$ ) ( $\mu\text{g RE}/100 \text{ Kcal}$ ;  $\mu\text{g}/100 \text{ Kcal}$ ;  $\text{mg } \alpha \text{ TE}/100 \text{ Kcal}$ ). The products are classified according to increasing time of storage from left to right ( $n = 5$ ). The limits are defined by straight lines. Lower and upper limits of vitamin A: 60–180  $\mu\text{g RE}/100 \text{ Kcal}$ ; lower and upper limits of vitamin D: 1–3  $\mu\text{g}/100 \text{ Kcal}$ ; lower and upper limits of vitamin E: 0.5–3.0  $\text{mg } \alpha \text{ TE}/100 \text{ Kcal}$ .

corresponds to 0.1  $\mu\text{g kg}^{-1}$  for aflatoxin B1 and 200  $\mu\text{g kg}^{-1}$  for fumonisins. Among the 19 analyzed products, only one product reached the limit for aflatoxin AFB1 (FI\_6,  $4.2 \pm 0.2 \mu\text{g kg}^{-1}$ ) and another for fumonisins (FI\_8, FB1 =  $198.1 \pm 3.5 \mu\text{g kg}^{-1}$  and FB2 =  $40.6 \pm 1.1 \mu\text{g kg}^{-1}$ ).



**Figure 3.** Acid and peroxide values of the FI according to the time of storage. Different letters above each bars indicate significant difference ( $p < 0.05$ ). The limits are defined by straight lines. Maximal limit of peroxide value: 10  $\text{meqO}_2 \text{ kg}^{-1}$  of lipid; maximal limit of acid value: 0.6% lipid.

### 3.7. Packaging Characterization

The packaging materials used to commercialize the FI were by decreasing frequency of use: aluminum/polyethylene multilayer, polyethylene, polyethylene/polypropylene multilayer, and cellophane/epoxy resin (Table 1).

OTR and WVTR were sometimes equal to zero if packaging is structured by an aluminum thin layer which is a very good barrier to oxygen and water transfer (Table 4). Such very good packaging materials were used for 7 FI out of 13, particularly for FI stored over long periods of time (>12 months, i.e., FI\_10, FI\_11, and FI\_12).

### 3.8. Determinants of the Degradation of Lipids and Vitamins

The key determinants of the stability of FI were explored by a correlation approach using PCA with the biochemical composition, oxidation level, OTR, and WVTR. However, this first analysis was not satisfying because several variables were not well projected in two or three dimensions. It was not surprising because of the great discrepancy between the product formulations, the packaging types, and conditions of storage (duration, temperature, etc.). In order to obtain a better representation of the variability of the data, OTR and WVTR data were excluded and a new PCA was realized. As expected for moisture sorption isotherm of starchy



**Table 3.** Major volatile compounds detected and contents in fumonisins and aflatoxin of FI ( $n = 3$ ).

	Hexanal [mg eqBB/100 g] <sup>a)</sup>	Pentanal [mg eqBB/100 g] <sup>a)</sup>	Nonanal [mg eqBB/100 g] <sup>a)</sup>	Fumonisin FB1 [ $\mu\text{g kg}^{-1}$ ]	Fumonisin FB2 [ $\mu\text{g kg}^{-1}$ ]	Aflatoxin AFB1 [ $\mu\text{g kg}^{-1}$ ]
FI_1	$5.38\text{e}^{-09} \pm 1.77\text{e}^{-09}$	$6.83\text{e}^{-10} \pm 2.23\text{e}^{-10}$	$4.35\text{e}^{-09} \pm 6.47\text{e}^{-10}$	—	—	—
FI_2_L2_1	$2.73\text{e}^{-0} \pm 1.19\text{e}^{-09\text{ab}}$	$6.27\text{e}^{-10} \pm 3.13\text{e}^{-10\text{a}}$	$6.03\text{e}^{-10} \pm 2.16\text{e}^{-10\text{a}}$	$113.1 \pm 3.6^{\text{b}}$	$16.4 \pm 0.4^{\text{a}}$	—
FI_2_L1_2	$2.49\text{e}^{-09} \pm 4.10\text{e}^{-10\text{a}}$	$4.89\text{e}^{-10} \pm 9.95\text{e}^{-11\text{a}}$	$6.45\text{e}^{-10} \pm 9.04\text{e}^{-11\text{a}}$	$122.3 \pm 2.0^{\text{b}}$	$32.0 \pm 1.2^{\text{a}}$	—
FI_2_L2_3*	$2.86\text{e}^{-09} \pm 2.48\text{e}^{-10\text{ab}}$	$5.25\text{e}^{-10} \pm 1.41\text{e}^{-11\text{a}}$	$6.53\text{e}^{-10} \pm 1.33\text{e}^{-10\text{a}}$	$115.7 \pm 3.0^{\text{b}}$	—	—
FI_2_L1_4*	$3.54\text{e}^{-09} \pm 5.39\text{e}^{-10\text{b}}$	$6.15\text{e}^{-10} \pm 8.28\text{e}^{-11\text{a}}$	$6.05\text{e}^{-10} \pm 4.83\text{e}^{-11\text{a}}$	$118.5 \pm 2.1^{\text{b}}$	—	—
FI_3_L2_1	$9.69\text{e}^{-10} \pm 1.46\text{e}^{-10\text{c}}$	$1.89\text{e}^{-10} \pm 1.87\text{e}^{-11\text{b}}$	$3.80\text{e}^{-10} \pm 3.12\text{e}^{-11\text{b}}$	$121.9 \pm 3.6^{\text{bc}}$	—	—
FI_3_L1_2	$1.00\text{e}^{-09} \pm 3.90\text{e}^{-10\text{c}}$	$2.43\text{e}^{-10} \pm 1.07\text{e}^{-10\text{bc}}$	$1.63\text{e}^{-10} \pm 2.06\text{e}^{-10\text{b}}$	$117.5 \pm 3.0^{\text{bc}}$	—	—
FI_3_L2_3*	$1.97\text{e}^{-09} \pm 5.38\text{e}^{-10\text{d}}$	$4.55\text{e}^{-10} \pm 1.69\text{e}^{-10\text{c}}$	$3.54\text{e}^{-10} \pm 3.18\text{e}^{-10\text{b}}$	$114.5 \pm 3.2^{\text{bc}}$	—	—
FI_3_L1_4*	$2.95\text{e}^{-09} \pm 1.18\text{e}^{-09\text{d}}$	$3.78\text{e}^{-10} \pm 9.76\text{e}^{-11\text{c}}$	$3.22\text{e}^{-10} \pm 3.09\text{e}^{-10\text{b}}$	$126.7 \pm 4.4^{\text{bc}}$	—	—
FI_4	$7.74\text{e}^{-10} \pm 9.36\text{e}^{-11}$	$1.59\text{e}^{-10} \pm 2.41\text{e}^{-11}$	$7.06\text{e}^{-10} \pm 9.42\text{e}^{-11}$	$123.6 \pm 4.9^{\text{c}}$	$37.7 \pm 1.0^{\text{c}}$	—
FI_5	$2.69\text{e}^{-09} \pm 5.97\text{e}^{-10}$	$4.50\text{e}^{-10} \pm 1.49\text{e}^{-10}$	$1.83\text{e}^{-09} \pm 8.73\text{e}^{-11}$	$134.7 \pm 4.4^{\text{d}}$	$30.1 \pm 0.8^{\text{c}}$	—
FI_6	$2.29\text{e}^{-09} \pm 5.40\text{e}^{-10}$	$2.70\text{e}^{-10} \pm 4.98\text{e}^{-11}$	$1.24\text{e}^{-09} \pm 3.49\text{e}^{-10}$	—	—	$4.2 \pm 0.2^{\text{a}}$
FI_7	$2.71\text{e}^{-09} \pm 8.89\text{e}^{-10}$	$3.10\text{e}^{-10} \pm 1.07\text{e}^{-10}$	$2.88\text{e}^{-09} \pm 6.19\text{e}^{-10}$	$133.5 \pm 3.3^{\text{d}}$	—	—
FI_8	$9.63\text{e}^{-09} \pm 3.24\text{e}^{-09}$	$1.21\text{e}^{-09} \pm 4.14\text{e}^{-10}$	$3.81\text{e}^{-09} \pm 6.16\text{e}^{-10}$	$198.1 \pm 3.5^{\text{e}}$	$40.6 \pm 1.1^{\text{c}}$	—
FI_9	$9.91\text{e}^{-10} \pm 3.35\text{e}^{-10}$	$1.82\text{e}^{-10} \pm 4.51\text{e}^{-11}$	$6.72\text{e}^{-10} \pm 1.62\text{e}^{-10}$	—	—	—
FI_10	$5.64\text{e}^{-09} \pm 1.92\text{e}^{-09}$	$9.28\text{e}^{-10} \pm 3.99\text{e}^{-10}$	$3.97\text{e}^{-10} \pm 8.00\text{e}^{-12}$	$105.2 \pm 3.1^{\text{a}}$	—	—
FI_11	$2.63\text{e}^{-09} \pm 3.65\text{e}^{-10}$	$4.35\text{e}^{-10} \pm 6.75\text{e}^{-11}$	$1.33\text{e}^{-10} \pm 1.92\text{e}^{-11}$	$116.0 \pm 1.9^{\text{b}}$	$30.3 \pm 0.7^{\text{c}}$	—
FI_12	$1.35\text{e}^{-08} \pm 2.19\text{e}^{-09}$	$2.14\text{e}^{-09} \pm 4.17\text{e}^{-10}$	$1.22\text{e}^{-09} \pm 1.02\text{e}^{-10}$	—	—	—
FI_13	$1.28\text{e}^{-08} \pm 1.67\text{e}^{-10}$	$2.86\text{e}^{-09} \pm 3.00\text{e}^{-10}$	$9.90\text{e}^{-10} \pm 6.71\text{e}^{-11}$	—	—	—

<sup>a)</sup>mg equivalent benzyl butyrate/100 g fresh product. Data are presented as mean  $\pm$  standard deviation for 100 g or kg of fresh product. Values with distinct lowercase superscript letters in the same column are significantly different ( $p \leq 0.05$ ). —, not detected.


**Table 4.** Oxygen and water vapor transmission rates of fortified infant flour packaging.

Code	Packaging	OTR <sup>a)</sup> [ $\text{cm}^3 \text{mm}^{-2}$ ]	WVTR <sup>b)</sup> [ $\text{g mm}^{-1}$ ]
FI_1	PE <sup>c)</sup>	17.46	0.03
FI_2	Alu/PE <sup>d)</sup>	0.00 <sup>e)</sup>	0.00 <sup>e)</sup>
FI_3	Alu/PE <sup>d)</sup>	0.00 <sup>e)</sup>	0.00 <sup>e)</sup>
FI_4	PE <sup>c)</sup>	13.87	0.02
FI_5	PE <sup>c)</sup>	13.87	0.02
FI_6	Alu/PE <sup>d)</sup>	0.00 <sup>e)</sup>	0.00 <sup>e)</sup>
FI_7	PE/PP <sup>f)</sup>	12.14	0.03
FI_8	PE <sup>c)</sup>	20.96	0.03
FI_9	PE <sup>c)</sup>	13.87	0.02
FI_10	C/Epo <sup>g)</sup>	0.00 <sup>e)</sup>	0.02
FI_11	Alu/PE <sup>d)</sup>	0.00 <sup>e)</sup>	0.00 <sup>e)</sup>
FI_12	Alu/PE <sup>d)</sup>	0.00 <sup>e)</sup>	0.00 <sup>e)</sup>
FI_13	Alu/PE <sup>d)</sup>	0.00 <sup>e)</sup>	0.00 <sup>e)</sup>

<sup>a)</sup>OTR, oxygen transmission rate; <sup>b)</sup>WVTR, water vapor transmission rate; <sup>c)</sup>PE, polyethylene; <sup>d)</sup>Alu/PE, aluminum/polyethylene; <sup>e)</sup>Aluminum multilayer packaging are considered as total oxygen and water barriers; <sup>f)</sup>PE/PP, polyethylene/polypropylene; <sup>g)</sup>C/Epo, cellophane/epoxy resin.

products, a positive correlation (0.78) between  $a_w$  and water content was observed. As a consequence, only water content was integrated in the PCA (Figure 4).

The three components extracted explained 79% of the variability of the data set. The representation of the variables along



**Figure 4.** Principal component analysis of the determinants of the stability of the FI (VE: vitamin E; VA: vitamin A; VD: vitamin D; PV: peroxide value; AV: acid value).

the first component accounted for 51% of the variability of the data, the second for 18%, and the third one for 10%. Despite the design of the study triggering high variability in FI's formulation, packaging materials, and conditions of storage, the set of analyzed products presented variables very well correlated (negatively or positively) with time of storage. The contents in vitamins A and E were negatively correlated with the time of storage and the correlation was rather strong with  $R = 0.86$  for vitamin A and

$R = 0.85$  for vitamin E. The peroxide content and the time of storage were positively correlated ( $R = 0.81$ ). The content in volatile compounds was not correlated to the lipid content of the FI.

#### 4. Discussion

The composition of FI sold in low-income countries was not always adequate in terms of fatty acids composition, lipophilic vitamins content, peroxide value, and mycotoxins content. The nutritional profiles of the 13 FI greatly differed depending on the formulation and on the location of the producing unit. The FI produced in Africa and in Asia contained three times more lipids than the products from northern countries. Such difference can be explained by the fact that children in developing countries need to cover a higher level of macronutrients with these fortified foods than children from northern countries that have access to a large choice of food to cover their needs. Indeed, the daily diet of children from southern countries is poorly diversified and can generate nutrient and caloric deficiencies.

Even if the FI lipid content greatly differs, most of them had a similar lipid profiles with three predominant fatty acids (i.e.,  $C_{18:2\ n-6}$  followed by  $C_{18:1\ n-9}$ , and  $C_{16:0}$ ). Lipids are crucial macronutrients that support most of the growth during the first 6 months of life and represent 40% to 60% of caloric requirements. Then, the lipid need gradually decreases to  $\approx 35\%$  of the total energy intake at 24 months. Palmitic acid, the third major FA in the fortified FI, plays an important role in infant nutrition.<sup>[2]</sup> It is the main saturated FA in breast milk. Nevertheless, its bioavailability will depend on food matrix composition (content in  $Ca_2^+$ ) and on its position on the triacylglycerol backbone. In breast milk, palmitic acid is located in *sn*-2 position in the triacylglycerol, the latter position will favor its absorption and role in the bone mineralization.<sup>[2]</sup> In vegetable oils used in our fortified FI, palmitic acid is predominantly located in external positions.<sup>[2]</sup> We can thus suspect that the consumption of formulas with high contents of saturated fat from vegetable oils or palm olein will not favor  $C_{16:0}$  absorption, but can be counterbalanced by complementary breast feeding until 24 months.

Globally,  $\omega 6/\omega 3$  ratios were high in fortified FI and sometimes excessive compared to the recommended range (5–15). Furthermore, the amounts of essential FA (linoleic acid  $C_{18:2\ n-6}$  and  $\alpha$ -linolenic acid  $C_{18:3\ n-3}$ ) present in the FI were sometimes too high. These FA represent approximately  $43.3 \pm 8.8\%$  and  $2.8 \pm 0.9\%$  of the total FA in the fortified FI, while it represents only 10.5–12.7% and 0.72–0.95% of the total FA in human milk.<sup>[3]</sup> Linoleic acid (LA) and  $\alpha$ -linolenic acid (ALA) intake recommendations for 6-month-old infant to 5-year-old children are 3.0–4.5% and 0.5–0.6% of energy intake, respectively,<sup>[38]</sup> which corresponds to 8.1–25.2 and 1.4–3.4 Kcal/day. If the FI analyzed in our study were consumed on a daily basis, LA intakes would be higher than the nutritional recommendations.<sup>[38]</sup> For a daily consumption of 100 g flours, an average  $29.7 \pm 11.7$  Kcal would be attributed to LA, which is higher than the recommendation, while ALA intakes would be satisfying.

Unbalanced LA intake can decrease blood DHA and cerebral lipids of infants.<sup>[4]</sup> Excessive intakes of ALA and LA can limit DHA biosynthesis by a competitive inhibition of PUFA bioconversion.<sup>[3,5]</sup> Conversely, a decrease in ALA contents in

infant food complement while maintaining  $C_{18:2\ n-6}/C_{18:3\ n-3}$  ratio below 5 could promote DHA biosynthesis. Unbalanced  $\omega 6/\omega 3$  ratios during infancy and childhood have long-term consequences on adipogenesis and lipid metabolism.<sup>[3,5]</sup>

In most FI, over fortification in vitamins was observed to ensure that vitamins would still remain in the product until its consumption, whatever the storage conditions. However, consumption of large amounts of vitamin A over a long period of time can induce liver damage, bone abnormalities, and vomiting.<sup>[39]</sup> Furthermore, in most developing countries, children aged 6–59 months are supplemented twice a year with a dose of 100 000 UI. The daily intakes from diet should not exceed 900  $\mu\text{g}$ , which is well above the mean requirement of about 200  $\mu\text{g}/\text{day}$  for infants.<sup>[39]</sup> In our experiments, seven FI could be at risk of excessive contents. Risk of hypervitaminosis (D or E) were already documented.<sup>[39,40]</sup> In addition, one can wonder about the nature and safety of the lipophilic vitamin overages degradation products. This aspect should be further investigated.

Globally, a strong effect of time of storage was observed on the set of FI despite heterogeneous conditions of storage in the supply-chain. This strong effect of time of storage is an important outcome of our study. For the FI produced in Europe and purchased in Africa with time of storage >12 months, the vitamin contents were below the recommended amounts. Two hypothesis can be formulated; either the vitamins were not adequately added during the industrial production but this hypothesis is so unlikely that it can be rejected, or the temperature during shipping and storage have activated chemical degradation within the products. There was a clear trend of vitamin degradation and peroxide generation over the storage period (between 1 and 19 months) in the “real supply chain.” The use of high barrier packaging material (ALU/PE) did not modify this trend. An overall adequacy of the FI, with acceptable nutritional profiles and low levels of degraded lipids, was observed up to 12 months, which could be a reasonable expiration date. According to the climatic conditions in tropical countries, the average temperatures are high and the temperature control within supply chain is not always possible. Indeed, vitamin A is susceptible to light and oxygen exposure while vitamin E is more stable. For instance, 90% degradation of vitamin A degradation was observed in fortified wheat flour after a 3-month storage at 40 °C.<sup>[16]</sup> On the contrary, vitamin E has been reported to be stable over 70 days even after opening a unit dose of infant milk.<sup>[41]</sup>

According to their  $a_w$ , the studied FI can be altered during storage mainly by hydrolysis reactions ( $a_w > 0.25$ ) and enzymatic activity ( $a_w > 0.35$ ). On the contrary, very low  $a_w$  (<0.2) favors very high oxidation rate, so the ideal window in terms of  $a_w$  is narrow probably (0.2–0.3). In our set of FI, three products were in this window and six products had  $a_w$  that exceeded 0.35.

With regards to peroxide and acidity values, they were higher when the storage duration increases, but the products were still acceptable for human consumption up to 9 months of storage, whatever the packaging used. The degradation reactions are rapid, especially in food matrices in powder form and at low  $a_w$ .

FI have a powdery structure, with a high surface area between the product and atmosphere which makes them an ideal matrix for the propagation of lipid oxidation.<sup>[42]</sup> Moreover, reactive oxygen species can initiate the oxidation of PUFA and, as a consequence, induce the production of hydroperoxides as well as other

oxidation compounds such as aldehydes (hexanal, pentanal, and propanal),<sup>[43]</sup> breakdown of *n*-6 FA (hexanal, pentanal) and *n*-3 FA (propanal).<sup>[44]</sup> These molecules are good markers of infant formula oxidation and participate in the off-flavor development.<sup>[45]</sup>

## 5. Conclusions

The panorama of fortified infant foods collected “in the field” after 1 to 19 months of storage indicated that their lipid profile and nutritional quality could be improved. Indeed, in most FI stored less than 6 months, lipophilic vitamin overages were observed. Long time of storages (>12 months) have deleterious impact on the fortified food nutritional profile despite the use of protective packaging. The expiration dates should be shortened in order to guarantee an optimal quality. In addition,  $\omega 6/\omega 3$  ratios should be maintained below 15. Such recommendation could improve the biological efficacy of fortified FI and better help preventing deficiencies. In perspectives, it would be very interesting to develop a mathematical model that could predict the degradation of lipophilic vitamins in complex matrices stored under different environmental conditions. This model would help optimizing vitamin levels and limiting overages, so that the level of added vitamins is acceptable from production up to consumption of fortified food products. Such model would improve significantly the vitamin profile of fortified foods and would constitute a helpful tool to abide by the regulation and fulfill infant needs.

## Abbreviations

ALA, alpha-linolenic acid; Alu/PE, aluminium/polyethylene; AV, acid value;  $a_w$ , water activity; C/Epox, cellophane/epoxy resin; CMV, mineral and vitamin complex; DAD, diode array detector; DHA, docosahexaenoic acid; EU, European Union; FA, fatty acid; FAME, fatty acid methyl ester; FFA, free fatty acid; FI, fortified infant flour; GC, gas chromatography; HPLC, high-performance liquid chromatography; HPLC DAD FLUO, high pressure liquid chromatography with diode array or fluorometric detector; HS-SPME, head-space solid phase microextraction; LA, linoleic acid; MSMS, tandem mass spectrometry; MUFA, monounsaturated fatty acid; *n*, number of repetitions; OTR, oxygen transmission rate; PCA, principal component analysis; PE, polyethylene; PE/PP, polyethylene/polypropylene; PTFE, polytetrafluoroethylene; PUFA, polyunsaturated fatty acid; PV, peroxide value; RE, retinol equivalent; ROS, reactive oxygen species; SFA, saturated fatty acid; UHPLC-MS, ultra high pressure liquid chromatography coupled to mass spectrometry; UV, ultra violet; VA, vitamin A; VE, vitamin E; VD, vitamin D; WVTR, water vapor transmission rate

## Supporting Information

Supporting Information is available from the Wiley Online Library or from the author.

## Acknowledgements

This work was supported by the French Ministry of Higher Education, Research and Innovation (scholarship program for PhD grant) and by the French Research on Lipid & Nutrition (<https://lipides-et-nutrition.net/legln/>). The authors warmly acknowledge Pierre Villeneuve for welcoming them in his lipid chemistry team (LiFFHeS, Lipids Functions Functionalities in Heterogeneous Systems) and all members of the team for their support and advice in the undertaken work. Arnaud Laillou is a UNICEF

staff member. The opinions and statements in this article are those of the authors and may not reflect official UNICEF policies.

## Conflict of Interest

The authors declare no conflict of interest.

## Keywords

lipid oxidation, mycotoxins, packaging, vitamins, volatile compounds

Received: April 19, 2019

Revised: July 26, 2019

Published online: October 2, 2019

- [1] C. Hayashi, J. Krasevec, R. Kumapley, V. Mehra, M. de Onis, E. Borghi, M. Blössner, M. F. Urrutia, E. B. Prydz, U. Serajuddin, *Levels and Trends in Child Malnutrition: Key Findings of the 2018 Edition of the Joint Child Malnutrition Estimates Report*, UNICEF/WHO/International Bank for Reconstruction and Development, Washington, DC **2018**.
- [2] S. M. Innis, *Adv. Nutr.* **2011**, *2*, 275.
- [3] G. Ailhaud, F. Massiera, P. Weill, P. Legrand, J.-M. Alessandri, P. Guesnet, *Prog. Lipid Res.* **2006**, *45*, 203.
- [4] P. Guesnet, J.-M. Alessandri, *Biochimie* **2011**, *93*, 7.
- [5] F. Massiera, P. Barbry, P. Guesnet, A. Joly, S. Luquet, C. M. Brest, T. Mohsen-Kanson, E.-Z. Amri, G. P. Ailhaud, *J. Lipid Res.* **2010**, *51*, 2352.
- [6] T. Sanghvi, J. Ross, H. Heymann, *Food Nutr. Bull. Suppl.* **2007**, *28*, S167.
- [7] I. Chakravarty, *Food Nutr. Bull. Suppl.* **2000**, *21*, 135.
- [8] G. Baldi, E. Martini, M. Catharina, S. Muslimatun, U. Fahmida, A. B. Jahari, Hardinsyah, R. Frega, P. Geniez, N. Grede, *Food Nutr. Bull. Suppl.* **2013**, *34*, S35.
- [9] L. Serra-Majem, R. Ortega, J. Aranceta, A. Entrala, A. Gil, *Public Health Nutr.* **2001**, *4*, 1331.
- [10] C. Bourlieu, A. Deglaire, S. C. De Oliveira, O. Ménard, Y. Le Gouar, F. Carrière, D. Dupont, *OCL* **2017**, *24*, D206.
- [11] K. G. Dewey, *Food Nutr. Bull. Suppl.* **2009**, *30*, S239.
- [12] M. Gibbs, K. B. Bailey, R. D. Lander, U. Fahmida, L. Perlas, S. Y. Hess, C. U. Loechl, P. Winichagoon, R. S. Gibson, *J. Food Compos. Anal.* **2011**, *24*, 418.
- [13] R. A. Gibson, B. Muhlhausler, M. Makrides, *Matern. Child Nutr.* **2011**, *7*, 17.
- [14] N. Roos, J. C. Sørensen, H. Sørensen, S. K. Rasmussen, A. Briend, Z. Yang, S. L. Huffman, *Matern. Child Nutr.* **2013**, *9*, 47.
- [15] B. Öztürk, *Eur. J. Lipid Sci. Technol.* **2017**, *119*, 1500539.
- [16] Y. M. Hemery, A. Laillou, L. Fontan, V. Jallier, R. Moench-Pfanner, J. Berger, S. Avallone, *Food Chem.* **2018**, *240*, 43.
- [17] Y. Kim, E. Strand, R. Dickmann, J. Warthesen, *J. Food Sci.* **2000**, *65*, 1216.
- [18] J. Wirth, N. Petry, S. Tanumihardjo, L. Rogers, E. McLean, A. Greig, G. Garrett, R. Klemm, F. Rohner, *Nutrients* **2017**, *9*, 190.
- [19] K. Eichler, S. Hess, C. Twerenbold, M. Sabatier, F. Meier, S. Wieser, *Value Health* **2017**, *20*, A545.
- [20] C. L. Luthringer, L. A. Rowe, M. Vossenaar, G. S. Garrett, *Glob. Health: Sci. Pract.* **2015**, *3*, 446.
- [21] Anonymous, *AACC Method 44-15A*, MN American Association of Cereal Chemists, Saint-Paul, MN **2000**.
- [22] Y. M. Hemery, L. Fontan, R. Moench-Pfanner, A. Laillou, J. Berger, C. Renaud, S. Avallone, *Food Chem.* **2015**, *184*, 90.
- [23] A. Laillou, C. Renaud, J. Berger, R. Moench-Pfanner, L. Fontan, S. Avallone, *Food Nutr. Bull. Suppl.* **2014**, *35*, 449.

- [24] International Vitamin A Consultative Group [IVACG], *Conversion Factors for Vitamin A and Carotenoids*, International Life Sciences Institute Research Foundation/Human Nutrition Institute, Washington, DC **2002**.
- [25] M. E. Shils, M. Shike, *Modern Nutrition in Health and Disease*, Lippincott Williams & Wilkins, Philadelphia, PA **2006**.
- [26] J. Folch, M. Lees, G. Sloane Stanley, *J. Biol. Chem.* **1957**, 226, 497.
- [27] Anonymous, *Official Methods and Recommended Practices of the American Oil Chemists' Society: Physical and Chemical Characteristics of Oil, Fats and Waxes*, AOCS PRESS, Urbana, IL **1996**.
- [28] Anonymous, *Official Methods of Analysis of AOAC International: Official Method 940.28. Fatty acids (free) in crude and refined oil*, AOAC, Washington, DC **1990**.
- [29] M. Lebrun, A. Plotto, K. Goodner, M.-N. Ducamp, E. Baldwin, *Postharvest Biol. Technol.* **2008**, 48, 122.
- [30] C. V. Corrales, M. Lebrun, F. Vaillant, M. N. Madec, S. Lortal, A. M. Perez, G. Fliedel, *Food Res. Int.* **2017**, 96, 113.
- [31] O. Olagunju, N. Mchunu, N. Durand, P. Alter, D. Montet, O. Ijabadeniyi, *LWT* **2018**, 98, 533.
- [32] L. K. Massey, *Permeability Properties of Plastics and Elastomers: A Guide to Packaging and Barrier Materials*, William Andrew, Norwich, NY **2003**.
- [33] Anonymous, *Commission Directive 2006/125/CE on Processed Cereal-Based Foods and Baby Foods for Infants and Young Children*, Official Journal of the European Union, Brussels **2006**.
- [34] T. Moyssiadi, A. Badeka, E. Kondyli, T. Vakirtzi, I. Savvaidis, M. G. Kontominas, *Int. Dairy J.* **2004**, 14, 429.
- [35] Anonymous, *Codex Standard for Edible Fats and Oils not Covered by Individual Standards (CODEX STAN 19-1981, Rev. 2017)*, Codex Alimentarius, Geneva **2017**.
- [36] Anonymous, *Commission Regulation EEC 2568/91 on the Characteristics of Olive Oil and Olive-Residue Oil and on the Relevant Methods of Analysis*, Official Journal of the European Union, Brussels **1991**.
- [37] Anonymous, *Commission Regulation EEC 1881/2006 Setting Maximum Levels for Certain Contaminants in Foodstuffs*, Official Journal of the European Union, Brussels **2006**.
- [38] Anonymous, *Interim Summary of Conclusions and Dietary Recommendations on Total Fat & Fatty Acids Summary of Total Fat and Fatty Acid Requirements for Adults, Infants (0-24 months) and Children (2-18 years)*, WHO, Washington, DC **2008**.
- [39] Anonymous, *Vitamin and Mineral Requirements in Human Nutrition*, WHO, Washington, DC **2004**.
- [40] C. Masterjohn, *Med. Hypotheses* **2007**, 68, 1026.
- [41] J. L. Chávez-Servín, A. I. Castellote, M. C. Lopez-Sabater, *Food Chem.* **2008**, 106, 299.
- [42] A. Briend, *Br. J. Nutr.* **2001**, 85, S175.
- [43] E. N. Frankel, *Lipid Oxidation*, Elsevier, Amsterdam **2014**.
- [44] E. Goicoechea, M. D. Guillen, *Eur. J. Lipid Sci. Technol.* **2014**, 116, 395.
- [45] J. L. Chávez-Servín, A. I. Castellote, M. C. López-Sabater, *Food Chem.* **2008**, 107, 558.