



Alternative domestic rodent pest management approaches to address the hazardous use of metal phosphides in low- and middle-income countries

Alexander M. Stuart¹ · Jens Jacob² · Adedayo Michael Awoniyi³ · Federico Costa^{3,4,5} · Luwieke Bosma⁶ · Yonas Meheretu^{7,8} · Nyo Me Htwe² · Stephanie Williamson¹ · Michael Eddleston⁹ · Ambroise Dalecky^{10,11} · Sheila Willis¹

Received: 12 December 2023 / Revised: 24 June 2024 / Accepted: 5 August 2024
© The Author(s) 2024

Abstract

Metal phosphides, particularly aluminium phosphide (AlP) and zinc phosphide (Zn_3P_2), are widely used in low- and middle-income countries (LMICs) as rodenticides in urban and domestic environments due to their low cost and high toxicity to rodent pests. However, they are also highly toxic to humans with no antidote available and have been associated with numerous fatal cases of intentional and accidental poisoning. This paper reviews alternatives to metal phosphide use for rodent pest management in urban and domestic environments, highlights case studies of effective alternative approaches, and provides recommendations for research and policy. This review identifies numerous alternative methods available for managing rodent pests in domestic/urban settings that can replace metal phosphides. These include chemical methods, i.e. rodenticides, and non-chemical methods, e.g. rodent-proofing, sanitation and trapping. However, because the majority of chemical rodenticides qualify as highly hazardous pesticides due to acute human health toxicity, environmental toxicity, and/or bioaccumulation, simply selecting substitute chemical rodenticides to replace metal phosphides are likely to replace one set of hazards with others. Thus, careful risk and hazard assessments are needed when considering substituting with other chemicals. Overall, we need to move away from current levels of rodenticide reliance towards more integrated and ecologically based approaches.

Keywords Ecologically based rodent management · Highly hazardous pesticides · Integrated pest management · Pesticide poisoning · Pesticide regulation · Urban policy

Communicated by Christian Imholt.

✉ Alexander M. Stuart
alex@pan-uk.org

¹ Brighthelm Centre, Pesticide Action Network UK, Brighton, UK

² Julius Kühn-Institute, Federal Research Institute for Cultivated Plants, Institute for Epidemiology and Pathogen Diagnostics, Rodent Research, Münster, Germany

³ Institute of Collective Health, Federal University of Bahia, Canela, Salvador 40110-040, Brazil

⁴ Centro de Pesquisas Gonçalo Moniz, Fundação Oswaldo Cruz, Ministério da Saúde, Rua Waldemar Falcão, 121, Salvador, Bahia, Brasil

⁵ Department of Epidemiology of Microbial Diseases, Yale School of Public Health, New Haven CT06511, USA

⁶ MetaMeta Research, Wageningen, The Netherlands

⁷ Department of Wildlife, Fish and Environmental Studies, Swedish University of Agricultural Sciences, Umeå, Sweden

⁸ Department of Biology and Institute of Mountain Research and Development, Mekelle University, Mekelle, Tigray, Ethiopia

⁹ Centre for Pesticide Suicide Prevention, University of Edinburgh, Edinburgh, UK

¹⁰ IRD, Aix Marseille Univ, LPED, Marseille, France

¹¹ BIOPASS 2, IRD, UGB, Cirad, ISRA, UFR S2ATA, Université Gaston Berger, Saint-Louis, Senegal

Introduction

The metal phosphides, aluminium phosphide (AIP), zinc phosphide (Zn_3P_2 ; herein referred to as ZnP) and to a lesser extent, magnesium phosphide (Mg_3P_2 ; herein referred to as MgP), are widely used as fumigants for stored products and as rodenticides due to their low cost, wide availability and high toxicity to a broad range of insect and rodent pests (Bumrah et al. 2012; Marashi 2015; Trakulsrichai et al. 2017; UNEP et al. 1988). However, metal phosphides are highly toxic to humans and other vertebrates (UNEP et al. 1988). Due to their high toxicity, rapid onset of systemic symptoms (with high risk of mortality within 24 h) and lack of specific antidote, the case fatality is as high as 67% for AIP and 25% for ZnP (Chugh et al. 1998; Gunnell et al. 2017; Shadnia et al. 2010). Thus, metal phosphides have been associated with numerous fatal cases of intentional and accidental poisoning, with cases reported across the world, especially in low- and middle-income countries (LMICs) (Anand et al. 2011; Bhandari and Basnet 2022; Proudfoot 2009; Yogendranathan et al. 2017). For example, in Pakistan, Iran and Nepal, AIP is reported as one of the most frequent agents for fatal self-poisoning (Etemadi-Aleagha et al. 2015; Ghimire et al. 2022). In Zimbabwe, AIP is one of the most common causes of pesticide poisoning, recently causing the death of three children after it was applied to stored maize (NAFDAC 2022). In Thailand, India, and Nepal, ZnP is a common self-poisoning agent as well as frequently involved in accidental poisonings, including young children (Ghimire et al. 2022; Peshin et al. 2014; Trakulsrichai et al. 2017; Varghese and Erickson 2022). In Tanzania, a survey conducted in 2006 revealed that ZnP was one of the most common poisoning agents of children under 18 years old (Lekei et al. 2017).

Metal phosphides are available in a range of formulations designed for different uses. These formulations include solid tablets (3 g or 12 g of 56% AIP), pellets, compressed discs, bag blankets, paste or powder—the last of which is commonly sold in sachets (Murali et al. 2009; Proudfoot 2009; UNEP et al. 1988). Uses include phytosanitary treatment of food and non-food products for import/export, commercial/large-scale insect pest control for stored food products, smallholder grain storage insect pest control, domestic/peri-domestic rodent pest control in and around houses (including grain stores), and agricultural rodent pest control (Buckle 1994; Bumrah et al. 2012; Nayak et al. 2020).

AIP and MgP are unstable solids used as a fumigant via the release of phosphine gas (PH_3) (Marashi 2015; Proudfoot 2009). Phosphine gas is released when AIP or MgP comes into contact with atmospheric moisture, as well as water and other neutral or acid pH liquids. However,

phosphine release is even more vigorous after contact with an acid, such as hydrochloric acid in the stomach (i.e. gastric acid) (Knight 2013; Proudfoot 2009). When used for rodent pest management, either pellets or tablets are applied to rodent burrows which are subsequently closed up with soil with the aim to gas the rodents (Buckle 1994).

ZnP is a relatively stable compound that can be applied as a bait; thus, it is more commonly used for rodent pest management than AIP and MgP (Marashi 2015). In addition, ZnP requires a sufficiently acid pH environment for hydrolysis to occur, releasing phosphine gas upon contact with acids such as gastric acid (Knight 2013; Trakulsrichai et al. 2017). When used against rodent pests, ZnP is typically applied in baits at concentrations of 2%, but concentrations can vary, ranging from 1 to 5% (Buckle 1994; UNEP et al. 1988). In many LMICs, 80% wettable powder formulations are commonly available as a low-cost pre-mix for bait preparation by the user (e.g. Bhandari and Basnet 2022; Buckle 1999; Stuart, A., pers. obs.; CIBRIC 2023; FPA 2023; PPD 2022).

Metal phosphides pose a high risk to human health. Routes of entry into the body include via ingestion and via inhalation of the highly toxic phosphine gas, which is denser than air and can thus travel short distances, e.g. to adjacent buildings (Proudfoot 2009; Yogendranathan et al. 2017). When ingested, metal phosphides are hydrolysed by gastric acid into phosphine, which is absorbed into the bloodstream. Phosphine is a respiratory toxin that inhibits many enzymatic systems, such as cytochrome C oxidase and oxidative respiration, which can result in renal failure and liver failure (Yogendranathan et al. 2017). Other severe clinical symptoms of phosphine poisoning include circulatory collapse, hypotension, pulmonary oedema, congestive heart failure and cardiac arrhythmia (Trakulsrichai et al. 2017). The main uses of concern to human health include i) when metal phosphides are used as a fumigants in or near to human habitation—which can lead to accidental poisoning from phosphine gas to bystanders/ adjacent buildings, ii) occupational exposure to phosphine gas in stores, warehouses or during transport, and iii) when solid formulations are easily accessible during use or when being stored or sold for later use (either against insect or rodent pests)—which can lead to intentional or unintentional poisoning via ingestion (Proudfoot 2009).

To address such risks to human health, many countries have banned, withdrawn or restricted the use of metal phosphides (CILSS 2022; Etemadi-Aleagha et al. 2015; Karunarathne et al. 2021; PAN 2022). For example, ZnP has been banned/withdrawn in at least seven countries and the US Environmental Protection Agency (EPA) only permits consumer products of ZnP and other rodenticides to be sold as block or paste poison bait, with loose bait forms (e.g. pellets, grains, powders) no longer permitted (EPA 2023). Recently, the US EPA also issued a proposed interim

decision to classify ZnP as a “restricted use pesticide” that can only be applied by certified pesticide applicators (EPA 2022). In West Africa and several other countries, AIP, MgP and ZnP are already classified as “restricted use pesticides” (CILSS 2022). In addition, many private voluntary standards (PVS, also known as voluntary sustainability standards) in certified food and fibre supply chains and retailer companies have included AIP and ZnP in their prohibited chemical lists (IPMcoalition 2023).

Despite their hazards, metal phosphides remain widely available and/or are largely unregulated in many LMICs due to lack of bans and restrictions or due to lack of enforcement. To support a transition away from metal phosphides, it is important that regulators, policymakers, the supply chain sector and users are aware of the alternatives (Stuart et al. 2023). To address this knowledge gap, we reviewed available information on alternatives to metal phosphide use for rodent pest management, identify further research needs and make recommendations for decision makers. In addition, we provide two case studies demonstrating less hazardous alternative approaches in LMICs that are at least as effective in reducing rodents as metal phosphides. Defining a universally acceptable threshold or tolerance limit for effective or successful rodent pest management is challenging due to location-specific factors. However, the primary goal of any rodent control intervention is to reduce rodent infestation and its causative factors to a level that no longer adversely affects the community (CDC 2006). Metal phosphide-based interventions typically yield short-term rodent population reductions of 30–90%, and control levels of around 70% are somewhat considered effective (Eisemann et al. 2003). Reviews on alternatives for agricultural rodent pest management are comprehensively covered in the literature (e.g. Brown et al. 2017; Buckle and Smith 2015; Singleton et al. 2021; Swanepoel et al. 2017). In this paper, by contrast, we focus on alternative rodent pest management approaches for domestic/urban environments (from household to municipal scale) where risks of human exposure to metal phosphides are greatest (Md Meftaul et al. 2020).

Alternatives to metal phosphides for domestic rodent pest management

There are a number of chemical and non-chemical tools and methods available that can be used as a direct alternative to metal phosphide rodenticides or as part of an integrated approach to replace their use in urban and domestic settings. These are described in the following section. However, it should be emphasised that all chemical approaches to rodent control may present a risk to human and environmental health. They usually only provide short-term fixes (Lee et al. 2022), while potentially accumulating long-term problems

due to the spread of behavioural or genetic resistance for some compounds (Colvin and Jackson 1999). Sustainable rodent pest management is generally best achieved if a variety of methods are employed (Witmer 2019).

Chemical rodent pest management methods

The chemical rodenticides used to manage rodent pests are numerous and can be grouped into three main categories: 1) acute (fast-acting) rodenticides, 2) first generation anticoagulants (FGARs) and 3) second generation anticoagulants (SGARs).

Acute rodenticides

Acute rodenticides include AIP, MgP and ZnP, as well as alpha-chloralose, alpha-chlorohydrin, carbon dioxide, bromethalin, cholecalciferol (vitamin D3), norbormide, sodium monofluoroacetate (compound 1080), strychnine and yellow phosphorus (Erickson and Urban 2004; Isackson and Irizarry 2023; Jacob and Buckle 2018). Alpha-chloralose kills via hypothermia and is considered to be mostly effective against house mice (*Mus musculus*) at temperatures below 15 °C (Meehan 1984; UFAW 2021). However, it is also highly toxic to birds and aquatic organisms and moderately toxic to mammals, including humans (Table 1; Lewis et al. 2016). Alpha-chlorohydrin is a toxicant sterilant which can cause sterility after a single dose and is highly toxic to birds (Ericsson 1982; Lewis et al. 2016). Carbon dioxide causes asphyxiation by hypoxia. It binds to haemoglobin, which prevents the uptake of oxygen leading to death within minutes (Permentier et al. 2017). For rodent management, carbon dioxide was recently registered in the United States as a solidified dry ice pellet form that should be placed in rodent burrow entrances (DC_Health 2018). Cholecalciferol, the biologically active form of vitamin D, is completely non-toxic in small amounts, but massive single doses or prolonged low-level exposure causes hypercalcemia in rodents leading to death in 3–4 days (Klemann et al. 2023). Bromethalin is a neurotoxic diphenylamine that poisons the central nervous system that ultimately causes respiratory arrest (Hygnstrom et al. 1994). Norbormide is largely *Rattus* species-specific and causes extreme and irreversible vasoconstriction that results in rapid death (Hayes and Laws 1991). To overcome sub-lethal effects and consequent bait shyness, promising prodrug forms of norbormide have been developed to delay the action of the rodenticide and improve palatability (Campbell et al. 2015). Sodium monofluoroacetate (compound 1080) inhibits several metabolic processes, leading to seizures and shock (Cottral et al. 1947). Strychnine acts as an antagonist of glycine receptors and generates uncontrollable muscle spasms (Isackson and Irizarry 2023; Sherley 2004). As with the metal phosphides, the symptoms

Table 1 Hazard profile of acute and anticoagulant rodenticides (Source: Erickson and Urban 2004; Lewis et al. 2016; PAN 2021). Obsolete rodenticides are excluded. Y = yes

Active ingredi- ent	Mammals— Acute oral LD ₅₀ (mg kg ⁻¹) ^a	Birds— Acute oral toxicity	Bio-concentra- tion factor	PPDB assessment on human health	WHO Classification ^b	Fatal if inhaled (H330) ^c	GHS ⁺ repro (1A,1B) ^d	WHO/FAO HHP criteria met ^e	PAN HHP list ^f
<i>Acute rodenti-</i>									
<i>cides</i>									
Alpha-chloralose/ chloralose	400	High	Low risk	Moderately toxic	Narcotic	II			
Alpha-chloro- hydrin	131	High	–	Moderately toxic	Kidney and liver toxicant	1b		Y	Y
Aluminium phosphide	8.7	High	–	Highly toxic	Heart and lung toxicant	FM		Y	Y
Bromethalin	2	High	–	Highly toxic	Both acute and chronic effects can occur follow- ing ingestion	1a		Y	Y
Sodium fluoroacetate (1080)	1.2	High	–	Highly toxic	CNS effects include trem- ulousness, hallucina- tions, convul- sions, and respiratory depression	1a		Y	Y
Carbon dioxide (dry ice)	100,000*	–	Low risk	Harmful by inhalation	Chronic exposure can produce adaptation as well as producing metabolic acidosis	Not listed			
Strychnine	16	Mod	–	Highly toxic	Convulsant	1b		Y	Y
Cholecalciferol (Vit. D3)	1426	Low-mod	Low potential	Very toxic			Y	Y	Y
Zinc phosphide	12	High	–	Highly toxic	May be fatal if swallowed	1b		Y	Y

Table 1 (continued)

Active ingredi- ent	Mammals— Acute oral LD ₅₀ (mg kg ⁻¹) ^a	Birds— Acute oral toxicity	Bio-concentra- tion factor	PPDB assessment on human health	WHO Classification ^b	Fatal if inhaled (H330) ^c	GHS ⁺ repro (1A,1B) ^d	WHO/FAO HHP criteria met ^e	PAN HHP list ^f
Magnesium phosphide	10.4	High	–	Moderately toxic	Y		Y	Y	
Norbormide	> 52.0	Mod	–	Harmful if swallowed	O				
<i>First genera- tion antico- agulants</i>									
Chlorophaci- none	3.15	High	Low risk	Highly toxic	Causes internal haemor- rhages	Suspected car- diovascular toxicant	1a	Y	Y
Coumatetralyl	16.5	Mod	Threshold for concern	Highly toxic	May cause haemorrhag- ing	1b	Y	Y	
Diphacinone	2.3	Low-mod	–	Highly toxic			1a	Y	Y
Pindone	280	Mod	–	–		O		Y	
Warfarin	10.4	Low-mod	Low risk	Highly toxic	Causes internal haemor- rhages	May be absorbed through the skin and cause systemic poisoning	1b	Y	Y
<i>Second genera- tion antico- agulants</i>									
Brodifacoum	0.4	High	High potential	Highly toxic	Severe poison- ings may lead to massive haemorrhag- ing		1a	Y	Y
Bromadiolone	> 0.56	Mod	Threshold for concern	Very toxic	May cause haemorrhag- ing	Slight eye irritant	1a	Y	Y
Difenacoum	1.8	High	High potential	Highly toxic	May cause haemorrhag- ing	Vapour may cause diz- ziness or suffocation	1a	Y	Y

Table 1 (continued)

Active ingredi- ent	Mammals— Acute oral LD ₅₀ (mg kg ⁻¹) ^a	Birds— Acute oral toxicity	Bio-concentra- tion factor	PPDB assessment on human health	WHO Classification ^b	Fatal if inhaled (H330) ^c	GHS ⁺ repro (1A,1B) ^d	WHO/FAO HHP criteria met ^e	PAN HHP list ^f
Difethialone	0.56	High	High potential	Highly toxic	1a	Y	Y	Y	Y
Floucoumafen	0.25	Mod	Threshold for concern	Highly toxic	1a	Y	Y	Y	Y

^aBased on *Rattus norvegicus*—the preferred test species for evaluation of acute toxicity by the oral and inhalation routes. The rat is usually considered to be a good model for human effects (ECHA, 2017)

^bAccording to World Health Organisation (WHO): 1a = extremely hazardous, 1b = highly hazardous, II = moderately hazardous, O = obsolete, FM = fumigant (The solid form of aluminium phosphide is classified by WHO as 'Fatal if swallowed')

^cH330 'Fatal if inhaled', hazard classification according to the EU or Japan Globally Harmonised System (GHS). Phosphine (produced by Aluminium and Magnesium phosphides) is classified as H330

^dKnown or presumed human reproductive toxicant according to EU or Japan GHS

^eAccording to WHO/ Food and Agriculture Organisation (FAO) criteria for highly hazardous pesticides

^fAccording to Pesticide Action Network International (PAN) list of highly hazardous pesticides

*LC50 inhalation value

of yellow (elemental) phosphorus poisoning include multiple organ failure and has been associated with a high number of fatal human poisonings in India (Jeyapal and Sagar 2020; Soni et al. 2020).

These acute rodenticidal compounds vary considerably in mammalian toxicity (Table 1), and their registration status varies among countries. With the exception of norbormide, that is reportedly a selective *Rattus*-specific toxicant (Roszkowski 1965), these acute compounds are not genus or species-specific and, depending on dose and susceptibility, can harm non-target species via primary exposure if they consume bait directly or inhale the gases produced (Shore and Coeurdassier 2018). Rodenticides should thus be placed in tunnels or within secure and purpose-made bait boxes to reduce non-target risk. However—as with placing traps in covers—bait boxes can reduce their efficacy against some species (Buckle and Prescott 2010). For example, Brown rats (*Rattus norvegicus*) are known to be neophobic (i.e. fear of new objects) and it may take days or weeks for a brown rat to enter a bait box. Careful assessment of the infestation, pre-baiting to reduce neophobia, and appropriate placement of bait are recommended. It is essential that measures are taken to minimise human contact and protect non-target species.

Due to the rapid action of acute rodenticides, most, except for bromethalin, have a low risk of bioaccumulation; thus, secondary exposure of non-target species (from consumption of poisoned rodents) is usually not an issue (Table 2). In addition, genetic resistance is unlikely to develop among target species (Erickson and Urban 2004). However, those with moderate to high mammalian toxicity can also pose a significant health risk to humans and caution should be applied when considering these as substitutes for metal phosphides (Table 1). In addition, even though alpha-chloralose is considered to have a low risk to humans and non-target animals when applied correctly and indoors (CRRU 2021), a recent study of human pesticide poisoning cases in Morocco revealed that 33% of the investigated pesticide poisoning cases were due to alpha-chloralose poisoning (Bourekadi et al. 2021; Dalecky et al. 2023).

First generation anticoagulants (FGARs)

Anticoagulant rodenticides (AR) are delayed acting rodenticides that are toxic to all warm-blooded organisms and are classified as WHO Class Ib and thus meet the FAO/WHO defined criteria for Highly Hazardous Pesticides (HHPs) (FAO and WHO 2016). All anticoagulant rodenticides inhibit vitamin K(1)-2,3 epoxide reductase hampering the synthesis of vitamin K and the clotting factor cascade (Berny 2011). Therefore, the term “anti-vitamin-K” (AVK) compounds is sometimes used. Through their action, blood clotting is greatly reduced and rodents die because of excessive internal and external bleeding. However, in the case of

human or accidental animal exposure to an anticoagulant, vitamin K can be administered as an antidote if medical treatment is provided in time (Witmer 2019).

FGARs include chlorophacinone, coumatetralyl, diphacinone, pindone and warfarin (Buckle 1994). These compounds are less toxic, less persistent and less bioaccumulative than SGARs and require the uptake of multiple doses to be effective. However, they are not free from risk to humans and non-target wildlife, especially because larger quantities of these baits need to be applied, as rodents need to continuously feed upon them over several days for effective control. Due to continued and intensive use of FGARs over many years, some rodent populations of some species developed genetic resistance to them (Buckle 1994; Witmer 2019). The need for alternatives to manage FGAR-resistant populations led to the development of the more toxic SGARs.

Second generation anticoagulants (SGARs)

SGARs include bromadiolone, difenacoum, difethialone, flocoumafen and brodifacoum. These compounds are highly toxic by ingestion, presumed reproductive toxins and have long biological half-lives (CRRU 2021); thus, they are more persistent and bioaccumulative than FGARs (Eason et al. 2002; Vandenbroucke et al. 2008). Usually a single dose kills the target rodent, except against some species and anticoagulant resistant individuals (Buckle 1994). Of the SGARs, brodifacoum has the highest mammalian toxicity (lowest LD₅₀ value), and difenacoum the lowest (Table 1). As with FGARs, SGARs are not only highly toxic to rodents but also for other warm-blooded animals, including humans, and are moderately toxic to invertebrates. The presence of SGAR residues in invertebrates (Shore and Coeurdassier 2018), songbirds (Walther et al. 2021) and non-target rodents (Brakes and Smith 2005; Geduhn et al. 2014), along with reported poisonings of domestic pets (Soleng et al. 2022) highlights the risk of primary exposure to non-target organisms.

SGARs have a long soil and water half-life and compounds accumulate in target and non-target organisms, as indicated by a high bio-concentration factor (Table 1). Due to the delayed action of FGARs and SGARs, there is no bait shyness, but poisoned rodents can live and be active for several days after bait uptake, as well as continue to feed on bait if available. This is especially of concern where genetic resistance to the SGAR has occurred. Secondary exposure to non-target animals that consume dead or dying rodents is common and almost impossible to prevent. Furthermore, another secondary exposure route includes the consumption of non-target invertebrates that feed on SGAR baits, such as slugs and snails (Alomar et al. 2018). There are thus numerous reports of AR residues in terrestrial predators/scavengers and avian predators/scavengers (Hindmarch et al. 2019;

López-Perea and Mateo 2018), with multiple AR exposure potentially causing synergistic effects as well as cumulative effects (Lettoof et al. 2020). Also, because anticoagulant rodenticides take several days to kill the rodent, indoor use could potentially result in rodents dying under the house, in the roof or in a wall cavity where it is difficult to remove the carcass.

Aside from SGAR exposure to terrestrial non-target organisms, SGAR exposure and fate in aquatic environments also need attention, especially due to their acute toxicity to fish (Lewis et al. 2016). A recent review by Regenry et al. (2019a) highlighted that very little is known on this topic, but a number of studies have detected the presence of AR residues in fish, as well as other aquatic organisms, raw and treated wastewater, sewage sludge, and estuarine sediments. The authors concluded that the aquatic environment experiences a greater risk of anticoagulant exposure than previously thought, requiring more comprehensive monitoring data. Sewer baiting in particular has been identified as a significant emission source of SGARs in the aquatic environment (Gómez-Canela et al. 2014; Kotthoff et al. 2019; Regnery et al. 2020).

Due to the variability in rodenticide toxicity between different rodent species, the development of rodenticide resistance, including against some of the SGARs, or variability in neophobia between animals, chemical rodenticide treatment alone is unlikely to achieve sustained control in the long term (CRRU 2021). This becomes especially apparent when attempting large-scale municipal level control, where numerous unsuccessful examples exist (Easterbrook et al. 2005; Fernández et al. 2007; Lambropoulos et al. 1999; Lee et al. 2022). Thus, because of these issues, in addition to the environmental and human health risks and public opposition to chemical rodenticide use, more emphasis is needed on applying non-chemical methods of rodent management as described in the next section. Table 2 summarises advantages, disadvantages and effectiveness of the main groups of chemical and non-chemical methods.

Non-chemical rodent pest management methods

Cultural control

Preventative measures should be taken whenever possible to prevent rodent infestation. For household dwellings, food stores and indoor livestock holdings, this can be done to a certain degree by rodent-proofing buildings and blocking holes greater than 6 mm in diameter (UFAW 2021; WHO 1972). Close-fitting doors can be highly successful at keeping rodents out, but may not always be feasible. Other rodent exclusion methods include draft-sealing door strips, using galvanised wire mesh or a mix of small stones with mud or clay in openings, and repairing broken window and door

Table 2 Advantages and disadvantages of different rodent management methods for domestic rodent pest management

Method	Disadvantages	Advantages	Efficacy for short-term rodent control at a local scale (e.g. individual households/ buildings)
Acute rodenticides	High primary non-target exposure risk Sub-lethal doses can lead to conditioned aversion (behavioural resistance) No antidote available Animal welfare concerns	Low secondary exposure risks, except bromethalin Fast acting A single dose is sufficient	Variable, as behavioural resistance can significantly reduce efficacy. (integrated approach with sanitation and proofing also needed to prevent re-invasion)
First generation anticoagulants	High primary non-target exposure risk Multiple doses required Slow acting, can lead to bodies dying in wall/roof cavities High risk of genetic resistance developing among Norway rats and house mice Animal welfare concerns	Low secondary exposure risk (except for coumatetrallyl) Antidote available Bait aversion unlikely to develop (due to delayed effect)	Can be effective if measures are taken to reduce genetic resistance development, e.g. limited use in an integrated approach with other methods such as trapping. (sanitation and proofing also needed to prevent re-invasion)
Second generation anticoagulants	High primary non-target exposure risk High secondary exposure risk Slow acting, can lead to bodies dying in wall/roof cavities High risk of genetic resistance developing among Norway rats and house mice (except for brodifacoum, difethialone, flocoumafen) Animal welfare concerns	Antidote available A single dose is sufficient Bait aversion unlikely to develop (due to delayed effect)	Can be effective if measures are taken to reduce genetic resistance development, e.g. limited use in an integrated approach with other methods such as trapping. (sanitation and proofing also needed to prevent re-invasion)
Cultural control (e.g. rodent proofing and sanitation)		No animal welfare concerns	Can be effective if conducted properly as part of an integrated approach. Sanitation is especially effective when conducted at community/municipal scale
Physical control (e.g. trapping)	Animal welfare concerns Material quality may be poor, e.g. traps that are made with weak spring mechanisms Single capture traps require regular checking/re-setting Lack of evidence on efficacy	No toxic hazards No toxic hazards	Can be effective if used as part of an integrated approach and if materials are of sufficient quality, e.g. traps with spring mechanisms of sufficient strength.
Biological control (e.g. fertility control, predators, biorodenticides)	Lack of availability or under development Potential hazards/animal welfare concerns Predators may kill/injure non-target animals	Fertility control/ predators have no known toxic hazards	Available evidence is limited.

frames (Mari Saez et al. 2018; Mdangi et al. 2013). In situations, where it is not possible to rodent proof a food or grain store, stored food can be protected by keeping it inside closed containers, e.g. tins or claypots, with sealed lids or placing something heavy on top of a metal sheet lids, installing rat guards one metre above ground on the supporting legs of crib platforms, or surrounding grain sacks with wire mesh or fishnetting (Brown et al. 2020; Htwe et al. 2021, 2017; Mdangi et al. 2013).

Rodent-proofing should be also accompanied by sanitation, such as removing potential food, water and harbourage (FWAH), to make the habitat less attractive to rodents and reduce the availability of resources that support rodent population growth. A systematic review of the literature on municipal-scale urban rodent control by Lee et al. (2022) reported that removing FWAH is widely considered to be the most effective method of reducing rats. Within and around buildings, the availability of a wide range of materials will provide habitat for rodents. Rodents often find harbourage within rubbish, debris or other materials lying around buildings, overgrown vegetation or ditches and drainage systems. These areas should be kept as clean and tidy as possible to make them less attractive to rodents as places to live and breed (WHO 1972). Within households, it is important to clean up spills and crumbs, store food in closed containers and cover rubbish bins with tight-fitting lids (UCT 2019). Furthermore, closing sewers, improving drainage, and maintaining sanitation around water bodies are important to reduce available water supply in urban environments (Zepelini et al. 2021).

There are devices available to detect rodent infestation early on (Fuelling et al. 2011). For field rodents, forecast models are available to indicate the risk of rodent outbreaks (Esther et al. 2014; Krebs et al. 2004; Leirs et al. 1996) and associated risk of human infection with rodent-borne pathogens (Kazasidis and Jacob 2023). However, such predictive models are still lacking in urban settings. Forecasting gives the opportunity for timely management action before a rodent problem gets out of hand. In areas where rodents pose a significant problem, factors that contribute to the presence of rats extend beyond individual properties (Lee et al. 2022), thus coordinated community wide campaigns are recommended to remove FWAH and conduct trapping and/or rat hunting. For example, studies in Asia demonstrated significant reductions in rodent numbers and losses to stored grain following intensive daily trapping at the community level together with improved hygiene practices (Belmain et al. 2015). Permanent working groups for rodent management in agriculture (Brown et al. 2024) ensure the involvement of stakeholders and similar structures may serve this purpose in domestic rodent control. In towns and cities, large- or municipal-scale strategies are needed to comprehensively address infestations and their causes (Lee et al. 2022).

Physical control

If rodent infestation occurs in an urban or domestic setting, there are a number of alternatives to rodenticides available. Trapping using live or kill (snap or break-back) traps is often recommended for rodent control in and around buildings and small farms. For example, in household grain stores in Bangladesh, daily trapping was found to be as effective as the use of rodenticides, with the added benefit that the traps could be re-used for many years (Belmain et al. 2015; Krijger et al. 2020). In Uganda, trapping in individual households was highly effective at removing all rodent pests within a 16-day period (Eisen et al. 2018). Following a study in which high quality rat traps were provided to 200 households in poor urban communities in South Africa where poor sanitation, infrequent refuse removal and overcrowded living conditions led to high rodent infestations, over half of study participants that previously used pesticides for rodent control voluntarily stopped using pesticides (Roomaney et al. 2012). Another study across three African countries showed that community household trapping effectively reduced rodent pest populations and damage to stored grains (Taylor et al. 2012). The results of these studies thus indicate that trapping is effective, feasible and acceptable.

Of the many types of rodent traps, good quality snap traps are considered to be the most effective and humane (UFAW 2021). Newer types of kill traps have also been developed such as battery-powered traps that deliver an electric shock or a captive bolt that kill rodents quickly. However, it is essential before setting kill traps, especially those used outdoors for rats, to consider both the target animals and any other animals that may be present and may enter traps accidentally as they can also injure or kill non-target animals (Witmer 2019). To minimise the risk to children, non-target animals, livestock and pets, traps should not be set where they may come into contact with them or tamper-resistant boxes should be used (UFAW 2021).

Using the appropriate trap and bait for the situation and target species is important for achieving high efficacy (Burke et al. 2021; Witmer 2019). Trapping can be cumbersome and difficult to conduct when rodents are trap shy or present in structures that are hard to access for trap placement. Behavioural studies highlight that rodents are creatures of habit and prefer to follow the same runways, thus identifying these runways, such as by sprinkling a fine layer of flour or baby powder, and placing traps along these will improve their success (WHO 1972). Devices have also been developed to assess the technical properties of traps (Baker et al. 2012; Walther et al. 2024). Snap traps that have a double-peg type spring and a wide opening angle are likely to produce a more powerful trap, which is more likely to kill rodents quickly and humanely (Baker et al. 2012; UFAW 2021). Further guidance on how to evaluate the animal welfare impact and

efficacy of snap traps was recently produced by an expert working group in EU, with the goal of establishing a voluntary certification scheme (Schlötterburg et al. 2021). This has subsequently been used by the German Environment Agency to test several types of traps and identify those that meet minimum welfare standards (UFAW 2021).

Another type of trap includes glue (or sticky) boards that captures rodents using a non-toxic sticky substance and holds it until it is either removed or until it dies. However, they are indiscriminate and present serious animal welfare concerns and are hence banned in several countries (CRRU 2021; UFAW 2021). Numerous studies have aimed to identify compounds that repel rodents or to drive them away using (ultra)sound or vibration. This may help to mitigate the problem at the place where repellents are used but simply relocates the problem. However, habituation usually sets in sooner or later, especially if there is an attractive food source for them (Meehan 1984), and the efficacy of repellents and repellent devices is often poor (Hansen et al. 2016; Lund 1988).

Biological control

Biological control is often advocated as an appropriate technique to manage pest rodents. In urban/domestic settings, predators, such as cats, are frequently suggested to reduce the impact of rodents, yet there are very few studies evaluating their impact. One study in Eswatini found that the combined presence of cats and dogs at homesteads can reduce rodent foraging activity at experimental seed trays and movements of rodents (Mahlaba et al. 2017), but it is unclear how rodent abundance, damage and disease risk are affected by biocontrol with predators (Labuschagne et al. 2016; Ostfeld and Holt 2004). A number of parasites and diseases have been trialled against rodents in agricultural settings. However, few are commercially available. A rodent-specific parasite, *Sarcocystis singaporensis*, does not affect humans and has been trialled effectively in towns in Laos and in rice fields in Thailand (Jäkel et al. 2006, 2019). ‘BIORAT’, produced in Cuba, is a combination of warfarin sodium 0.02% + *Salmonella enteritidis* var. Danysz Lysine. However, there are no published data in mainstream literature to support or refute the use of BIORAT. In addition, both warfarin (a FGAR) and *S. enteritidis* are of human health concern (Brown et al. 2017).

Other biological control methods that are being investigated include fertility control to minimise proliferation of rodents by restricting reproduction instead of culling them (Jacoblinnert et al. 2021). This method is favoured by the public over using rodenticides to manage red squirrels (Dunn et al. 2018). Fertility control could minimise recolonisation by infertile residents that defend their territories against (fertile) immigrants. Only few fertility control products

are registered for use against rodents. They include ContraPest® (a combination of 4-vinylcyclohexene diepoxide and triptolide for black rat (*Rattus rattus*) and Norway rat (*R. norvegicus*) in the USA) (Pyzyńska et al. 2016; Siers et al. 2017), EP-1 (a combination of levonorgestrel and quinestrol for multimammate mice (*Mastomys natalensis*) in Tanzania) (Massawe et al. 2018; Selemani et al. 2021), and GonaCon™ (injectable product for black-tailed prairie dogs (*Cynomys ludovicianus*) in the USA) (Shiels et al. 2023). It is important that anti-fertility compounds are effective and safe for non-target species, present no risk to humans and the environment and are economically feasible. Assessments of the environmental fate of quinestrol and levonorgestrel indicate a short half-life in soil (1–2 weeks) and water (a few hours) (Tang et al. 2012a, b). However, a recent review of rodent fertility control by Massei et al. (2023) concluded that studies on the persistence of oral contraceptives in the environment and in the food chain were generally under-reported. For domestic rodent management in sensitive areas where tolerance for rodent infestation is low, fertility control may take too long to achieve the desired management aims. However, recent simulations show that fertility control using contraceptive bait may substantially reduce rodent populations when sustained over multiple rodent generations (Shuster et al. 2023).

Integrated pest management

There is a general dilemma in rodent control. The aim is to reduce the population of a mammalian species considered as pests that has many similarities in morphology and physiology to “wanted” mammalian species including humans. Whatever is bad for the rodent, often is also bad for non-target species. Welfare concerns regarding the use of acute rodenticides are also valid for FGARs and SGARs (Table 2). Welfare concerns may also relate to trapping and biocontrol because traps and predators alike can injure or kill rodents or non-target organisms (Parsons et al. 2018). Some studies have also revealed increased pathogen prevalence/transmission in surviving animals following efforts that focused on lethal control, such as with leptospirosis in rats (Lee et al. 2018), and killing rats in endemic plague foci could lead to the dispersal of rat flea vectors (Belmain 2018; Rahelinirina et al. 2023). Furthermore, reliance on a single method tends to lead to declining effectiveness over time, such as from genetic rodenticide resistance that is a common problem in some urban rodent populations (Witmer 2019). Municipal-scale approaches that focus on lethal control measures rather than addressing causative FWAH factors often fall short of methodological expectations (Lee et al. 2022). This highlights the importance of emphasising preventative action rather than only relying on reactive approaches, such

as rodenticide use (Byers et al. 2019; Colvin and Jackson 1999).

To have the greatest success in the long-term, an integrated approach (i.e. integrated pest management, integrated rodent management or ecologically based rodent management) is needed that considers environmental, ecological and socio-economic factors, with chemical pesticide use as a last resort (Brown et al. 2020; Lambropoulos et al. 1999; Rahelinirina et al. 2023; Witmer 2019). In urban environments, this requires careful planning, cooperation and coordination between local authorities, communities, and pest control operators to take into account the complexity of rat management (Awoniyi et al. 2024; Dalecky et al. 2024; Lambropoulos et al. 1999; Lee et al. 2022). Ongoing resourcing and funding to eliminate factors contributing to FWAH is essential to prevent rodent populations returning to pre-infestation levels (Lee et al. 2022). Community cooperation and awareness raising are also vital in rural communities where rodents have the opportunity to move from house to house. In addition, more attention and research are needed for non-lethal methods, early detection and prediction of rodent problems. Periodic monitoring of rodent populations is an often-overlooked component of rodent management that can instruct appropriate action(s) to be taken before a rodent population builds up (Witmer 2005). The next section provides examples of integrated approaches implemented in LMICs.

Case studies of effective alternatives for domestic rodent pest management

Urban rodent control in Brazil

Household rodent infestation is a major concern in most disadvantaged urban communities often called “favelas” of Brazil (Costa et al. 2014; Masi et al. 2010). Poor sanitation facilities and waste management in addition to dilapidated structures and the continued increase in the numbers of disadvantaged urban dwellers foster the proliferation of rodent populations in these communities. Rodents are reservoirs of important zoonotic pathogens (Costa et al. 2014) and are capable of destroying agricultural products and household properties worth billions of dollars per year (Childs et al. 1991) therefore necessitating regular control efforts.

A number of methods have been used for controlling rodent proliferation in Brazil, e.g. rodent trapping, application of rodenticides, habitat modification, sanitation interventions, education interventions (e.g. “rato fora” or “rats out”) and lately the combination of two or more of these methods, i.e. integrated rodent management—“IRM” (de Masi et al. 2009). Until recently, the application of rodenticides/pesticides was the principal control method (de

Masi et al. 2009; Papini et al. 2019; Pertile et al. 2022). Possibly because it is cheap, readily available and easy to apply. While rodenticides of coumarin derivatives such as brodifacoum and coumatetralyl are legally authorised in Brazil (FUNASA 2002), the application of other illegal rodenticides/pesticides, such as aldicarb (locally known as “chumbinho”), have been well documented in the country (Panis et al. 2021; Papini and Nakagawa 2014).

Reports of bait shyness, rodenticide resistance and the negative results from the sole application of rodenticides in Brazil (with rodent populations quickly returning to the pre-intervention state in 3 months after the chemical application) (Babolin et al. 2016; Pertile et al. 2022), together with the killing of non-targeted species and death of residents (Panis et al. 2021; Papini and Nakagawa 2014) have necessitated a paradigm shift from the sole application of chemical application during rodent control campaigns. For example, the number of poisonings and suicides among the Brazilian population resulting from rodenticides application totalled 26,651 and 16,867 between 2005 and 2011 (Papini and Nakagawa 2014). Additionally, the reporting rate of accidental and intentional human poisonings arising from rodenticides was 64.5 per 100,000 population, and 58.7 per 100,000, respectively, within the same period (Panis et al. 2021).

Therefore, researchers and other stakeholders in Brazil have continued to advocate in favour of IRM techniques during rodent control programmes (Awoniyi et al. 2022; de Masi et al. 2009). One example of an effective IRM programme in Brazil includes a study conducted in Sao Paulo (de Masi et al. 2009). The authors of this study reported that the rodent infestation rate reduced from 40% pre-intervention to 14.4% (64% percentage reduction) post-intervention following a combination of (i) chemical application (coumarin-based rodenticides), (ii) a waste management campaign (improvement of the surrounding environment to limit rodents’ access to food, water and harbourage), and (iii) an education intervention (community awareness programmes on TV, radio, and five local schools about the importance of preventing household rodent infestation and the procedures to achieve this).

Another example in Brazil, conducted by a group from Salvador combined (i) a chemical-based intervention (coumarin-based rodenticides application) that lasted one week, (ii) an education intervention that lasted throughout the study period (involving regular information campaigns to remind residents about the need to practise proper waste disposal and to keep pets’ food/water away from the reach of rodents), and (iii) a local government led infrastructural/environmental intervention that was conducted during the first year of the study and involved the modification of open sewers and pavements in common spaces and sidewalks to reduce rodents’ sources of food, water and harbourage

(Fig. 1) (Awoniyi et al. 2022). As a result of these interventions, there was a significant sustained reduction in the rodent infestation rate from 70% before, to about 20% after the interventions, offering a 71% reduction over approximately 12 months (T3–T5 of Fig. 2). The percentage reduction here is similar, but the effect lasts longer than most metal phosphides applications (Eisemann et al. 2003). To ensure the effectiveness of the intervention, three valleys (1, 2 & 3) with similar socio-environmental features were used for the study. Valleys 1 and 3 served as the chemical-treatment valleys and received all three categories of the interventions, while valley 2 received no chemical intervention. The long-term reduction in rodent infestations observed in all 3 valleys indicates that IRM technique is more effective in controlling the long-term population of rodents than the sole application of a control method (i.e. as practised prior to the intervention), particularly in disadvantaged urban communities. However, supplementing educational and infrastructural interventions with a chemical approach did not provide any noticeably greater benefit.

The two case studies from Sao Paulo and Salvador suggest there are alternative practicable ways of controlling the urban proliferation of rodents without predisposing residents and other non-target species to probable poisonings arising from the exclusive application of hazardous rodenticides/pesticides (Papini and Nakagawa 2014). Brazilian favelas are unplanned, characterised by inadequate sanitation facilities with most residents either unemployed or low-income earners, with most residents unable to afford sustainable IRM techniques, such as rodent-proofing buildings. Given the socio-economic situation of these communities, it is thus important for local governments to drive major rodent control programmes and to continue to support interventions (including infrastructural maintenance) to prevent rodent populations from reverting to pre-intervention levels (Awoniyi et al. 2021; Richardson et al. 2017).

Community ecologically based rodent management (EBRM) in rural households in Ethiopia

In Ethiopia, over 100 species of rodents have been recorded, with only 10 species considered to be significant pests. The main pest species (both field and storage) belong to the following six genera: *Arvicanthis*, *Gerbilliscus*, *Mastomys*, *Stenocephalemys*, *Tachyoryctes*, and *Rattus*. Rodenticide (mainly ZnP) application and trapping are two of the most practised rodent management methods in crop fields and storage areas in northern Ethiopia (Meheretu et al. 2022, 2010). However, the use of rodenticides in storage areas and houses presents a significant health risk to humans and domestic animals (Dalecky et al. 2023; Meheretu et al. 2022). For instance, a retrospective cross-sectional study from Debre Tabor general hospital, a district hospital in

northern Ethiopia, revealed that out of the 102 patients admitted to the hospital presenting acute poisoning between September 2013 and August 2016, rodenticide poisoning accounted for 56.9% of the cases, with a mortality of 18.6% (Endayehu and Shenkutie 2019). In a study in northern Ethiopia, 93% of surveyed cereal farmers reported the use of zinc phosphide rodenticide in rainfed crop fields to manage rodent pests (Meheretu et al. 2010). About 87% of these farmers decided to buy (apply) the rodenticide after noticing intense rodent activities in the fields, suggesting that rodent management is conducted in response to rodent damage, rather than integrated into routine pest management activities.

Traditional storage is often rudimentary and may, for instance, consist of bamboo mats held loosely together. These open storages attract vermin and rodents close to where people live, increasing the risk of zoonotic transmission, as well as causing food losses and contamination. Furthermore, inadequate waste disposal, grain and cattle feed storage methods aid the proliferation of rodent populations in villages thereby heightening public health concerns.

Poor knowledge of rodent management methods, including rodenticides, likely results in weak efficacy of action and treatment failure, as well as a public safety concern, leading to apathy and widespread acceptance of rodent pests in fields and storage areas (Meheretu et al. 2022). However, for the farmers in the highlands of Amhara, the problem became so severe, they reached out to the extension workers for help. This led to the development of a successful community-level EBRM programme that involved extension workers, Meta-Meta, a Dutch social enterprise, and the Amhara Bureau of Agriculture (Dalecky et al. 2024).

Due to a huge knowledge gap on alternative biological and ecological rodent management methods for small-holder farmers in Ethiopia, the programme began by developing a systematic, communal, and ecologically based approach from the ground up. This entailed engaging with farmers and communities to understand the extent of the problem followed by combining their indigenous knowledge with scientific research to work towards a sustainable solution. The steps for the development of community-level EBRM were as follows:

1. Focus group discussions and surveys on the knowledge, attitudes and perceptions were conducted with farmers in the highlands of Amhara.
2. Local knowledge was combined with academic research to develop alternative methods, i.e. a biological rodenticide using locally available botanicals and an EBRM package (Dalecky et al. 2024).
3. Trainings were held twice a year for agricultural experts, community leaders and watershed user associations (WUA) committees. Farmers in the highlands

of Amhara are organised in WUAs, which are registered under the regional government and responsible for watershed management (mainly soil and water conservation practices) as a community. Each WUA has an average area of 500 hectares and 150–200 households. The fact that the watershed community made the EBRM activities part of their collective work was critical to the success of EBRM in the area.

4. The WUA committees thereafter continued action planning with all WUA members. In action planning, they incorporated the timing of the EBRM actions in the lean


season (i.e. when rodent populations are still at lower density) and factored in the importance of collective action (e.g. a by-law was created specifying that any household who did not join in a collective activity such as flooding rodent burrows, would pay a penalty fee).

5. Activities implemented as part of the EBRM package included: measures to expose rodents to predators, removing harbourage, deep ploughing, flooding and plugging rodent burrows, setting up stone traps, using domestic cats (mainly in homesteads), and storing grains

Fig. 1 Infrastructural intervention in Pau da Lima, Salvador-BA, Brazil; **a** typical open sewer in Pau da Lima, **b** cross-sectional area of an aspect of the community during the intervention, and **c** modification of sidewalks to reduce sewer water accumulation



Fig. 2 Mean ($\pm 95\%$ confidence interval) rodent infestation levels in three valleys (valley 1 & 3 = chemical-treatment; valley 2 = non-chemical treatment) from 2014 to 2017. Each survey period, i.e. T1–T7, lasted 3–4 months (adapted from Awoniyi et al. 2022)



in locally made rodent-proof storage structures (e.g. see Fig. 3).

6. These EBRM activities were organised through campaigns, ensuring regular interventions with all households involved.

As well as reducing the pesticide poisoning risk, the EBRM campaign led to a substantial improvement in rodent control in comparison with prior efforts that relied principally on rodenticide use. A post-implementation survey with the WUA committee members from all of the watersheds that were included in the EBRM campaigns indicated that farmers perceived that rodent damages had been cut by a magnitude of up to 50% after two years of implementing EBRM. The survey also indicated that through horizontal learning, the uptake of the EBRM activities increased by up to threefold in the neighbouring watersheds that were not included in the EBRM campaigns. Currently, in 2023, there are at least 65 watersheds implementing EBRM, benefiting up to 13,000 households and covering about 33,000 hectares of land.

This case study focused on rural communities, including their homesteads, in an agricultural landscape. However, the lessons learned from EBRM within and around homesteads also provide useful guidance for rodent management in urban and peri-urban areas. Most notably, the case study highlights the need for community action with a focus on reducing the availability of food and shelter for rodents, for example, via proper garbage disposal, rodent-proofed storage and regular sanitation of houses, storage areas and gardens.

Policy recommendations

Following the literature review, we identify the following policy recommendations (summarised in Table 3).

Recommendation #1: restrict availability

Given the high proportion of poisoning cases associated with metal phosphides in LMICs, and the lack of an effective antidote, we recommend limiting access to this group of rodenticides. Restricting access to the means of suicide is also a globally accepted evidence-based intervention for suicide prevention, by giving persons in distress time for acute crises to pass (Utyasheva and Eddleston 2021; WHO 2021). In descending order of effectiveness, restricting pesticide access can include either 1) an outright ban of the chemical (the most effective and highest level of the industrial Hierarchy of Control, a well-established method of ranking risk

reduction measures (UNEP et al. 2022); 2) a ban of a certain formulation type; 3) limiting access by type of end-user, e.g. via professional certification; 4) limiting access by type of supplier; and/or 5) enforcing restrictions on how/where pesticides are stored (Eddleston and Gunnell 2020; UNEP et al. 2022; Varghese and Erickson 2022).

There are several studies that show that overall suicides have decreased following outright bans of acutely toxic HHPs (Knipe et al. 2017; Yan et al. 2023). With regard to metal phosphides, the results of bans have been mixed. In north India, AIP poisoning was the primary cause of poisoning-related deaths until 3 g 56% AIP tablets were banned from open sale in 2001. After which, AIP-related deaths fell dramatically, indicating a significant public health and pesticide regulatory success (Karunaratne et al. 2021). Preliminary results from Nepal (Ghimire, R. unpublished) also indicate that a switch from 3 to 12 g AIP tablets, which are less easily ingested, has resulted in a reduction in all poisoning deaths. However, in Iran, the ban of 3 g tablet formulations in 2007 did not achieve such success due to the continued availability of 3 g tablets via illegal imports (Etemadi-Aleagha et al. 2015; Hashemi-Domeneh et al. 2016). This highlights that when implementing bans in countries that have porous borders, there is the added need for strengthening border controls and/or a regional approach between neighbouring countries (OECD 2014; Stuart et al. 2023; UNEP et al. 2022).

The availability of ZnP formulations at concentrations as high as 80% also poses an unnecessary and added risk to human health. For example, in the US, most occupational exposure incidents to ZnP between 2012 and 2019 involved exposures to products containing either 62% or 67% ZnP (EPA 2022). In many LMICs, ZnP is commonly available as an 80% wettable powder formulation as a low-cost pre-mix for bait (Bhandari and Basnet 2022). However, 1 grain of 2.5–5% pre-mixed wheat bait is sufficient to kill a 15 g mouse and 3–5 grains of 2.5% bait is sufficient to kill a 150 g rat, demonstrating the effectiveness of much lower dose formulations which pose a reduced human health risk (Staples et al. 2003). In the US, 80% ZnP wettable powder formulations are no longer permitted. In EU countries, the highest concentration of ZnP in registered products is 3% (EPPO 2023), with similar restrictions in place in other countries, such as Australia and New Zealand (Eason et al. 2013; Hinds et al. 2023). Thus, it seems clear that, as a minimum, countries should ban the sale and production of the most hazardous metal phosphide formulations, such as small AIP tablet formulations and high concentration ZnP formulations, as well as prohibit the sale and use of other metal phosphide formulations for household use and for unlicensed users.

Recommendation #2: implement enforcement of restrictions

Unfortunately, introducing bans and restrictions in use is not sufficient if there is no action to enforce such policies. In many LMICs, the illegal street trade in unregistered or restricted-use rodenticides and other pesticides for rodent control is common and some of these pesticides may be just as hazardous as metal phosphides (Dalecky et al. 2023; Rother 2016). For example, in Brazil and South Africa, aldicarb and terbufos, both highly hazardous insecticides, are illegally sold by street vendors and used for rodent control, contributing to frequent fatal human poisoning cases (Davies et al. 2023; Panis et al. 2021; Papini and Nakagawa 2014; Rother 2016). Indomethacin, an anti-inflammatory drug for pain-relief in humans, is also illegally used as a rat poison in several African countries (Dalecky et al. 2023; Donga et al. 2022). It is thus important to follow up bans and restrictions with regular inspections and enforcement, along with public education regarding the dangers of such pesticides (Papini and Nakagawa 2014).

Recommendation #3: improve the availability of less hazardous alternatives

This review identifies effective alternative options for rodent pest management in urban and domestic environments, especially when conducted in an integrated approach. We recognise, however, that these may not always be available or affordable, or the end-user knowledge about such

alternatives and integrated approaches may be lacking (Rother 2016). To address this, government investment is needed in public information/awareness campaigns on alternative approaches to rodent pest management and the hazards of various chemical pesticides that are locally available (including illegal pesticides as indicated above). Due to the hidden costs from human pesticide poisoning, such as medical costs and related burdens on health services and society, there is also an economic argument for government subsidies to help make effective low-risk methods, such as traps, more affordable and available (UNEP et al. 2022). As identified in this review, providing rodent traps to poor urban communities, along with guidance on how to use them, has been proven to reduce demand for illegal as well as legal pesticides for rodent control (Roomaney et al. 2012).

Other important points to consider when promoting less hazardous alternatives include social barriers to adoption and the quality of the alternative options available—such as the strength of trap spring mechanisms, which can affect their efficacy and humaneness (Baker et al. 2012). By understanding such issues, interventions can be better targeted to address them. For example, during a recent household survey in New Zealand, one of the main concerns for household rat trapping for conservation aims was the perception that traps are a cruel and inhumane way to kill animals (Kaine et al. 2023). In response, the authors of the study recommend that promotional campaigns seeking to promote trapping should emphasise on the safety of traps, and the speed and efficacy with which they function. Public and private investment should also prioritise the development of low-risk methods (UNEP et al. 2022), such as supporting local trap producers to produce high quality traps at low cost and implement measures to minimise non-target risk.

If selecting substitute chemical rodenticides to replace metal phosphides, the least toxic rodenticides (including their formulations) should be prioritised over more toxic rodenticides following a careful risk and hazard assessment. The FAO/WHO International Code of Conduct on Pesticide Management recommends that governments should “make every reasonable effort to reduce risks posed by pesticides by making less toxic formulations available” (FAO and WHO 2014). Particular attention to this matter will be necessary in countries where the only alternative rodenticides available are those that also pose high levels of risk to humans and the environment. For example, for West African countries that implement the common regulations set by Sahelian Committee for Pesticides (CSP), the only rodenticides registered are brodifacoum and ZnP (CILSS 2022). When comparing anticoagulant rodenticides, Berny et al. (2014) identify that for rat control, FGARs and less potent SGARs should always be considered as the first choice. SGARS should only be used against rats, where there is evidence that infestations are resistant. However, because FGARs have low efficacy



Fig. 3 A rat-proof storage system developed by Mr Sisay Mengistie, a farmer from Ata Meher watershed. It is a square two-metre high container made of wood and corrugated iron that is sealed with a door and used to store wheat and barley. It is well suited for local adoption: the materials are available locally at a reasonable price, it is easy to build and adaptable, it can store large quantities of grain: up to 1500 kg per storage structure

Table 3 Key recommendations from this literature review

Recommendations	
1 Restrict availability	· Ban the sale and production of the most hazardous metal phosphide formulations, such as small AIP tablet formulations and high concentration ZnP formulations, as well as prohibit the sale and use of other metal phosphide formulations for household use and for unlicensed users. · In countries that have porous borders, strengthen border controls and/ or a regional approach between neighbouring countries.
2 Implement enforcement of restrictions	· Follow up bans and restrictions with regular inspections and enforcement, along with public education regarding the dangers of such pesticides
3 Improve the availability of less hazardous alternatives	· Invest in public information/awareness campaigns on alternative approaches to rodent pest management and the hazards of various chemical pesticides that are locally available. · Public and private investment should also prioritise the development of low-risk methods. · Conduct careful risk and hazard assessments if selecting substitute chemical rodenticides to replace metal phosphides so that the least toxic rodenticides (including their formulations) are prioritised over more toxic rodenticides.
4 Promote and invest in preventative measures, including community action	· Invest in sanitation and infrastructure to mitigate rodent infestations in urban and domestic environments, especially in lower-income areas. · Conduct government-led rodent management interventions and education/awareness campaigns at a community-level.
5 Strengthen research on less hazardous alternatives	· Fund further research to develop less hazardous alternatives that are economically viable and socially acceptable. · Conduct studies for different contexts and social-economic conditions to develop locally adapted approaches that balance risk, cost and efficacy.

against house mice, FGARs should only be used against mice where there is evidence that the local strain is susceptible, otherwise SGARs should be considered as the first choice. With regard to non-target risk, brodifacoum was ranked as posing the greatest overall potential risk to birds and mammals out of 9 rodenticides analysed (Erickson and Urban 2004). With regard to primary non-target risk, ZnP ranked the highest, with brodifacoum second. With regard to secondary non-target risk, brodifacoum ranked the highest, while ZnP ranked the lowest.

Recommendation #4: promote and invest in preventative measures, including community action

Both the case studies presented in this review highlight that investments in sanitation and infrastructure are essential to mitigate rodent infestations in urban and domestic environments. Reactive approaches to urban/domestic rodent management that do not address the underlying features that promote rat abundance are unlikely to be effective in the long term (Byers et al. 2019). Removing sources of food, shelter and water can prevent rodent population build up and thus negate the need for metal phosphides and other forms of lethal rodent pest control. However, this important first step towards sustainable rodent pest management is often neglected and so government investments in such measures are essential, especially in lower-income areas due

to infrastructure disinvestment and lower coping capacity of residents (Awoniyi et al. 2024; Jassat et al. 2013; Peterson et al. 2020). Long-term investments in such measures are also needed so as to prevent rodent populations from reverting to pre-intervention levels.

It should also be highlighted that although the same principles of cultural control apply, the approach of such interventions should be tailored to the local context. For example, the approach needed for a rural community with grain stores and scrub vegetation will be quite different to that needed for an urban poor community with overcrowding and open sewers. Overall, community action and area-wide interventions that consider the ‘total environment’ for sources of FWAH are essential due to the ability of rodents to move between households (Byers et al. 2019; Lee et al. 2022). It is thus important to conduct rodent management interventions and education/awareness campaigns at a community-level and for governments to lead these.

Recommendation #5: strengthen research on less hazardous alternatives

Due to a variety of welfare, hazard and economic concerns from most of the existing rodent pest management methods available, there is clear need to fund further research to develop less hazardous alternatives that are economically viable and socially acceptable. Research to further develop and demonstrate the effectiveness of non-toxic alternatives

such as biological control, fertility control, physical exclusion and trapping should be prioritised, as well as carefully assess and monitor any potential risks. Studies are also needed for different contexts and social-economic conditions to develop locally adapted approaches that balance risk, cost and efficacy (e.g. see Stuart et al. 2020). This requires optimising current methods, developing new methods and using a combination of several methods in an integrated approach.

Conclusions

Due to their high toxicity, rapid onset of systemic symptoms and lack of specific antidote, metal phosphides pose a high risk to human health, particularly if used inappropriately or for intentional poisoning. This is especially apparent for many LMICs where they are widely available to the public, perhaps due to lack of bans and restrictions or lack of enforcement. Our review of the literature, along with case studies, shows that there are numerous alternative methods for managing rodent pests in domestic/urban settings that can replace metal phosphides. These include chemical and non-chemical methods. However, because the majority of chemical rodenticides qualify as HHPs due to acute human health toxicity, environmental toxicity, and/or bioaccumulation, strategies that simply select substitute chemical rodenticides to replace metal phosphides should be approached with caution as these are likely to replace one set of hazards with others. Thus, these will need to be carefully assessed. In addition, there is the added risk of AR resistance development. It is evident that overall, we need to move away from current levels of rodenticide reliance and move towards more integrated and ecologically based approaches.

Author contribution

AMS, ME, SW, SW and JJ conceived the study. AMS and JJ conducted the literature search. YM, LB, AMA and FC provided the case studies. AMS, JJ, AMA, FC, LB and JM wrote the first draft. All authors critically revised the manuscript and approved the final version.

Acknowledgements We thank Prof. Andrea Rother and the anonymous reviewers for their helpful comments on a previous version of this manuscript.

Funding The Centre for Pesticide Suicide Prevention is funded by a grant from Open Philanthropy at the recommendation of GiveWell.

Data availability No datasets were generated or analysed during the current study.

Declarations

Conflict of interest The authors declare no conflict of interests.

Open Access This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

- Alomar H, Chabert A, Coeurdassier M, Vey D, Berny P (2018) Accumulation of anticoagulant rodenticides (chlorophacinone, bromadiolone and brodifacoum) in a non-target invertebrate, the slug, *Deroceras reticulatum*. *Sci Total Environ* 610–611:576–582. <https://doi.org/10.1016/j.scitotenv.2017.08.117>
- Anand R, Binukumar BK, Gill KD (2011) Aluminum phosphide poisoning: an unsolved riddle. *J Appl Toxicol* 31:499–505. <https://doi.org/10.1002/jat.1692>
- Awoniyi AM, Souza FN, Zeppelini CG, Xavier BIA, Barreto AM, Santiago DCC, Santana JO, da Silva EM, Costa F, Begon M, Khalil H (2021) Using Rhodamine B to assess the movement of small mammals in an urban slum. *Methods Ecol Evol* 12:2234–2242. <https://doi.org/10.1111/2041-210x.13693>
- Awoniyi AM, Venegas-Vargas C, Souza FN, Zeppelini CG, Hacker KP, Carvalho-Pereira T, Marins CL, de Santana MC, Pertile AC, Begon M, Ko AI, Diggle PJ, Reis MG, Childs JE, da Silva EM, Costa F, Khalil H (2022) Population dynamics of synanthropic rodents after a chemical and infrastructural intervention in an urban low-income community. *Sci Rep* 12:10109. <https://doi.org/10.1038/s41598-022-14474-6>
- Awoniyi AM, Barreto AM, Argibay HD, Santana JO, Palma FAG, Riviere-Cinnamond A, Dobigny G, Bertherat E, Ferguson L, Belmain S, Costa F (2024) Systematic surveillance tools to reduce rodent pests in disadvantaged urban areas can empower communities and improve public health. *Sci Rep* 14:4503. <https://doi.org/10.1038/s41598-024-55203-5>
- Babolin LdS, Almeida-Silva MJF, Potenza MR, Fava CD, Castro V, Harakava R, Okuda LH, Rebouças MM, Campos AEC (2016) Zoonosis associated to *Rattus rattus* and the impacts of the public actions to control the species. *Arq Inst Biol*. <https://doi.org/10.1590/1808-1657000832014>
- Baker SE, Ellwood SA, Tagarielli VL, Macdonald DW (2012) Mechanical performance of rat, mouse and mole spring traps, and possible implications for welfare performance. *PLoS ONE* 7:e39334. <https://doi.org/10.1371/journal.pone.0039334>
- Belmain SR, Htwe NM, Kamal NQ, Singleton GR (2015) Estimating rodent losses to stored rice as a means to assess efficacy of rodent management. *Wildl Res*. <https://doi.org/10.1071/wr14189>

- Belmain S (2018) A new rat control strategy for madagascar to prevent plague outbreaks. Natural Resources Institute, University of Greenwich. https://www.researchgate.net/publication/330987673_A_New_Rat_Control_Strategy_for_Madagascar_to_Prevent_Plague_Outbreaks#fullTextFileContent. Accessed 10 December 2023
- Berny P (2011) Challenges of Anticoagulant Rodenticides Resistance and Ecotoxicology. In: Margarita S (ed) Pesticides in the Modern World. IntechOpen, Rijeka, p 20. <https://doi.org/10.5772/19916>
- Berny P, Esther A, Jacob J, Prescott C (2014) Risk mitigation measures for anticoagulant rodenticides as biocidal products. Report. European Union, Luxembourg. <https://doi.org/10.2779/241180>
- Bhandari R, Basnet K (2022) A case report on zinc phosphide ingestion resulting to acute pancreatitis. *Annals Med Surg* 78:103859. <https://doi.org/10.1016/j.amsu.2022.103859>
- Bourekadi S, Birich B, El Hajjaji S, Ghandi M, Ait Daoud N, Ouaid M, Badrane N, Soulaymani Bencheikh R, Hani H, Mokhtari A, Slimani K, Soulaymani A (2021) Toxicological analysis of acute pesticides poisoning among Moroccan population. In: E3S web of conferences, p 319 <https://doi.org/10.1051/e3sconf/202131901054>
- Brakes CR, Smith RH (2005) Exposure of non-target small mammals to rodenticides: short-term effects, recovery and implications for secondary poisoning. *J Appl Ecol* 42:118–128. <https://doi.org/10.1111/j.1365-2664.2005.00997.x>
- Brown PR, Giraudoux P, Jacob J, Couval G, Wolff C (2024) Multi-stakeholder working groups to improve rodent management outcomes in agricultural systems. *Int J Pest Manag* 1–17. <https://doi.org/10.1080/09670874.2024.2363877>
- Brown PR, Singleton GR, Belmain SR, Htwe NM, Mulungu L, Mdangi M (2020) Advances in understanding rodent pests affecting cereal grains. In: Maier DE (ed) Advances in postharvest management of cereals and grains. Burleigh Dodds Science Publishing Limited, Cambridge, pp 1–30. <https://doi.org/10.19103/AS.2020.0072.04>
- Brown PR, Giraudoux P, Jacob J, Couval G, Wolff C (in press) Multi-stakeholder working groups to improve rodent management outcomes in agricultural systems.
- Brown PR, Douangboupah B, Htwe NM, Jacob J, Mulungu L, My Phung NT, Singleton GR, Stuart AM (2017) Control of rodent pests in rice cultivation. In: Sasaki T (ed) Achieving sustainable cultivation of rice, vol 2. Burleigh Dodds, Cambridge, UK, pp 343–376. <https://doi.org/10.19103/as.2016.0003.24>
- Buckle AP (1994) Rodent control methods: chemical. In: Buckle AP, Smith RH (eds) Rodent pests and their control. CAB International, Wallingford, pp 127–160. <https://doi.org/10.1079/9781845938178.0123>
- Buckle AP (1999) Rodenticides-their role in rodent pest management in tropical agriculture. In: Singleton G, Hinds L, Zhang Z (eds) Ecologically-based rodent management. Australian Centre for International Agricultural Research, Canberra, pp 163–177
- Buckle AP, Prescott CV (2010) Effects of tamper-resistant bait boxes on bait uptake by Norway rats (*Rattus norvegicus* Berk.). *Int J Pest Manag* 57:77–83. <https://doi.org/10.1080/09670874.2010.537793>
- Buckle AP, Smith RH (2015) Rodent Pests and their Control, 2nd edn. CAB International, Wallingford
- Bumrah GS, Krishan K, Kanchan T, Sharma M, Sodhi GS (2012) Phosphide poisoning: a review of literature. *Forensic Sci Int* 214:1–6. <https://doi.org/10.1016/j.forsciint.2011.06.018>
- Burke CB, Quinn NM, Stapp P (2021) Use of rodenticide bait stations by commensal rodents at the urban-wildland interface: Insights for management to reduce nontarget exposure. *Pest Manag Sci* 77:3126–3134. <https://doi.org/10.1002/ps.6345>
- Byers KA, Lee MJ, Patrick DM, Himsworth CG (2019) Rats about town: a systematic review of rat movement in urban ecosystems. *Front Ecol Evolution*. <https://doi.org/10.3389/fevo.2019.00013>
- Campbell KJ, Beek J, Eason CT, Glen AS, Godwin J, Gould F, Holmes ND, Howald GR, Madden FM, Ponder JB, Threadgill DW, Wegmann AS, Baxter GS (2015) The next generation of rodent eradication: innovative technologies and tools to improve species specificity and increase their feasibility on islands. *Biol Cons* 185:47–58. <https://doi.org/10.1016/j.biocon.2014.10.016>
- CDC (2006) Integrated pest management: conducting urban rodent surveys centres for disease control and prevention, US department of health and human services, Atlanta
- Childs JE, Glass GE, Leduc JW (1991) Rodent sightings and contacts in an inner-city population of Baltimore, Maryland, U.S.A. *Bull Soc Vector Ecol* 16:245–255
- Chugh SN, Aggarwal HK, Mahajan SK (1998) Zinc phosphide intoxication symptoms: analysis of 20 cases. *Int J Clin Pharmacol Ther* 36:406–407
- CIBRIC (2023) Pesticides and formulations registered for use in the country under the insecticides act, 1968 central insecticides board & registration committee (CIBRC), Directorate of plant protection, Quarantine & storage department of agriculture & cooperation India
- CILSS (2022) Liste globale des pesticides autorisés par le Comité Sahélien des Pesticides. Version de Octobre 2022. Permanent inter-state committee for drought control in the sahel (CILSS)
- Colvin BA, Jackson WB (1999) Urban rodent control programs for the 21st century. In: Singleton G, Hinds L, Zhang Z (eds) Ecologically-based Rodent management. Australian Centre for International Agricultural Research, Canberra, pp 243–258
- Costa F, Porter FH, Rodrigues G, Farias H, de Faria MT, Wunder EA, Osikowicz LM, Kosoy MY, Reis MG, Ko AI, Childs JE (2014) Infections by *Leptospira interrogans*, Seoul virus, and Bartonella spp. among norway rats (*Rattus norvegicus*) from the urban slum environment in Brazil. *Vector-Borne Zoonotic Dis* 14:33–40. <https://doi.org/10.1089/vbz.2013.1378>
- Cottral GE, Dibble GD, Winton B (1947) The Effect of sodium fluoroacetate (“1080” rodenticide) on white leghorn chickens. *Poult Sci* 26:610–613. <https://doi.org/10.3382/ps.0260610>
- CRRU (2021) CRRU UK Code of Best Practice. Best practice and guidance for rodent control and the safe use of rodenticides. Campaign for responsible rodenticide use UK (CRRU UK), UK
- Dalecky A, Garba M, Danzabarma AI, Etougbéché J, Badou S, Dossou H-J, Sow I, Niang CT, Diene O, Diallo I, Saghir MSA, Sidatt MEH, Van Steenberg F, Bal AB, Bosma L, Houéménou G, Atteynine SA, Hima K, Dobigny G, Meheretu Y (2023) Rodent proliferation in urban and agricultural settings of sub-Saharan Africa the dark side of synthetic chemical rodenticides. *Environnement, Risques & Santé* 22:205–211. <https://doi.org/10.1684/ers.2023.1731>
- Dalecky A, Sow I, Ibrahim Danzabarma AA, Garba M, Etougbéché J, Badou S, Dossou H-J, Niang CT, Diene O, Diallo I, Saghir MSA, Sidatt MEH, van Steenberg F, Bal AB, Bosma L, Houéménou G, Ag Atteynine S, Hima K, Dobigny G, Meheretu Y (2024) From problem to progress: rodent management in agricultural settings of sub-Saharan Africa and calling for an urban perspective. *Crop Prot*. <https://doi.org/10.1016/j.cropro.2024.106673>
- Davies B, Hlela M, Rother HA (2023) Child and adolescent mortality associated with pesticide toxicity in Cape Town, South Africa, 2010–2019: a retrospective case review. *BMC Public Health* 23:792. <https://doi.org/10.1186/s12889-023-15652-5>
- DC_Health (2018) Dry ice application for rodent abatement. Government of the District of Columbia, U.S. [https://dchealth.dc.gov/sites/default/files/dc/sites/doh/page_content/attachments/Dry%](https://dchealth.dc.gov/sites/default/files/dc/sites/doh/page_content/attachments/Dry%20Ice%20Application.pdf)

- 20Ice%20Flyer%208%2023%2018.pdf. Accessed 30 November 2023
- de Masi E, Vilaça PJ, Razzolini MTP (2009) Evaluation on the effectiveness of actions for controlling infestation by rodents in Campo Limpo region, São Paulo Municipality, Brazil. *Int J Environ Health Res* 19:291–304. <https://doi.org/10.1080/09603120802592723>
- Donga TK, Bosma L, Gawa N, Meheretu Y (2022) Rodents in agriculture and public health in Malawi: farmers' knowledge, attitudes, and practices. *Front Agron*. <https://doi.org/10.3389/fagro.2022.936908>
- Dunn M, Marzano M, Forster J, Gill RMA (2018) Public attitudes towards “pest” management: perceptions on squirrel management strategies in the UK. *Biol Cons* 222:52–63. <https://doi.org/10.1016/j.biocon.2018.03.020>
- Eason CT, Murphy EC, Wright GR, Spurr EB (2002) Assessment of risks of brodifacoum to non-target birds and mammals in New Zealand. *Ecotoxicology* 11:35–48. <https://doi.org/10.1023/a:1013793029831>
- Eason CT, Ross JG, Blackie HM, Fairweather A (2013) Toxicology and ecotoxicology of zinc phosphide as used for pest control in New Zealand. *N Z J Ecol* 37:1–11
- Easterbrook JD, Shields T, Klein SL, Glass GE (2005) Norway rat population in Baltimore, Maryland, 2004. *Vector-Borne Zoonotic Dis* 5:296–299. <https://doi.org/10.1089/vbz.2005.5.296>
- ECHA (2017). “Guidance on the Application of the CLP Criteria - Guidance to Regulation (EC) No 1272/2008 on Classification, Labelling and Packaging of Substances and Mixtures”. (Helsinki, Finland: European Chemicals Agency).
- Eddleston M, Gunnell D (2020) Preventing suicide through pesticide regulation. *Lancet Psychiatry* 7:9–11. [https://doi.org/10.1016/s2215-0366\(19\)30478-x](https://doi.org/10.1016/s2215-0366(19)30478-x)
- Eisemann JD, Petersen BE, Fagerstone KA Efficacy of zinc phosphide for controlling Norway rats, roof rats, house mice, *Peromyscus* spp., prairie dogs and ground squirrels: A literature review (1942–2000). In: Fagerstone KA, Witmer GW (eds) The tenth wildlife damage management conference, Hot Springs, AR, 2003. National wildlife research center, pp 335–349
- Eisen RJ, Atiku LA, Boegler KA, Mpanga JT, Ensore RE, MacMillan K, Gage KL (2018) An evaluation of removal trapping to control rodents inside homes in a plague-endemic region of rural Northwestern Uganda. *Vector Borne Zoonotic Dis* 18:458–463. <https://doi.org/10.1089/vbz.2018.2276>
- Endayehu Y, Shenkutie E (2019) Magnitude of acute poisoning and associated factors in debretabor general hospital, Ethiopia. *J Clin Toxicol* 9:1–4
- EPA (2022) Zinc phosphide proposed interim registration review decision case number 0026. United States Environmental Protection Agency
- EPA (2023) Rodent Control Pesticide Safety Review United States Environmental Protection Agency. <https://www.epa.gov/rodenticides/rodent-control-pesticide-safety-review>. Accessed 30 November 2023
- EPPO (2023) List of databases on registered plant protection products in the EPPO region. European and Mediterranean plant protection organization https://www.eppo.int/ACTIVITIES/plant_protection_products/registered_products. Accessed 11 November 2023
- Erickson W, Urban D (2004) Potential risks of nine rodenticides to birds and nontarget mammals: a comparative approach. United States Environmental Protection Agency
- Ericsson RJ (1982) Alpha-chlorohydrin (EpiBloc®) (1982) : A toxicant-sterilant as an alternative in rodent control. In: Marsh RE (ed) Proceedings tenth vertebrate pest conference, Davis, California, pp 6–9
- Esther A, Imholt C, Perner J, Schumacher J, Jacob J (2014) Correlations between weather conditions and common vole (*Microtus arvalis*) densities identified by regression tree analysis. *Basic Appl Ecol* 15:75–84. <https://doi.org/10.1016/j.baae.2013.11.003>
- Etemadi-Aleagha A, Akhgari M, Irvani FS (2015) Aluminum phosphide poisoning-related deaths in Tehran, Iran, 2006 to 2013. *Medicine* 94:e1637. <https://doi.org/10.1097/md.0000000000001637>
- FAO, WHO, (2014) The International code of conduct on pesticide management World Health Organization Food and Agriculture Organization of the United Nations, Rome
- FAO, WHO (2016) International code of conduct on pesticide management: guidelines on highly hazardous pesticides. Food and Agriculture Organization of the United Nations (FAO) or of the World Health Organization (WHO), Rome
- Fernández MS, Cavia R, Cueto GR, Suárez OV (2007) Implementation and evaluation of an integrated program for rodent control in a shantytown of Buenos Aires City, Argentina. *EcoHealth* 4:271–277. <https://doi.org/10.1007/s10393-007-0122-4>
- FPA (2023) List Of registered agricultural pesticides. Fertilizer and pesticide authority (FPA), Department Of Agriculture, Philippines
- Fuelling O, Klemann N, Endepols S (2011) Automatic and permanent rodent-monitoring: a proper method to evaluate rodenticide effects? In: 8th European Vertebrate Pest Management Conference, Berlin, Germany, vol 432. Julius-Kühn-Archiv, pp 150–151. <https://doi.org/10.5073/jka.2011.432.082>
- FUNASA (2002) Manual de Controle de Roedores. Fundação Nacional de Saúde (FUNASA), Ministério da Saúde, Brasília
- Geduhn A, Esther A, Schenke D, Mattes H, Jacob J (2014) Spatial and temporal exposure patterns in non-target small mammals during brodifacoum rat control. *Sci Total Environ* 496:328–338. <https://doi.org/10.1016/j.scitotenv.2014.07.049>
- Ghimire R, Utyasheva L, Pokhrel M, Rai N, Chaudhary B, Prasad PN, Bajracharya SR, Basnet B, Das KD, Pathak NK, Baral MP, Pande R, Paudel P, Shrestha SK, Bajracharya S, Chaudhary R, Malla GB, Sharma DR, Basnyat B, Maskey MK, Eddleston M (2022) Intentional pesticide poisoning and pesticide suicides in Nepal. *Clin Toxicol (phila)* 60:46–52. <https://doi.org/10.1080/15563650.2021.1935993>
- Gómez-Canela C, Barata C, Lacorte S (2014) Occurrence, elimination, and risk of anticoagulant rodenticides and drugs during wastewater treatment. *Environ Sci Pollut Res* 21:7194–7203. <https://doi.org/10.1007/s11356-014-2714-1>
- Gunnell D, Knipe D, Chang S-S, Pearson M, Konradsen F, Lee WJ, Eddleston M (2017) Prevention of suicide with regulations aimed at restricting access to highly hazardous pesticides: a systematic review of the international evidence. *Lancet Glob Health* 5:e1026–e1037. [https://doi.org/10.1016/s2214-109x\(17\)30299-1](https://doi.org/10.1016/s2214-109x(17)30299-1)
- Hansen SC, Stolter C, Imholt C, Jacob J (2016) Plant secondary metabolites as rodent repellents: a systematic review. *J Chem Ecol* 42:970–983. <https://doi.org/10.1007/s10886-016-0760-5>
- Hashemi-Domeneh B, Zamani N, Hassani-Moghaddam H, Rahimi M, Shadnia S, Erfantalab P, Ostadi A (2016) A review of aluminium phosphide poisoning and a flowchart to treat it. *Arh Hig Rada Toksikol* 67:183–193. <https://doi.org/10.1515/aht-2016-67-2784>
- Hayes WJ, Laws ER (1991) Handbook of pesticide toxicology: classes of pesticides. Academic Press, New York, NY
- Hindmarch S, Rattner BA, Elliott JE (2019) Use of blood clotting assays to assess potential anticoagulant rodenticide exposure and effects in free-ranging birds of prey. *Sci Total Environ* 657:1205–1216. <https://doi.org/10.1016/j.scitotenv.2018.11.485>
- Hinds LA, Henry S, Van De Weyer N, Robinson F, Ruscoe WA, Brown PR (2023) Acute oral toxicity of zinc phosphide: an assessment

- for wild house mice (*Mus musculus*). *Integr Zool* 18:63–75. <https://doi.org/10.1111/1749-4877.12666>
- Htwe NM, Singleton GR, Maw PP (2017) Post-harvest impacts of rodents in Myanmar; how much rice do they eat and damage? *Pest Manag Sci* 73:318–324. <https://doi.org/10.1002/ps.4292>
- Htwe NM, Sarathchandra SR, Sluydts V, Nugaliyadde L, Singleton GR, Jacob J (2021) Small mammal communities, associated damage to rice and damage prevention in smallholder rice storage facilities in Sri Lanka. *Crop Prot*. <https://doi.org/10.1016/j.cropro.2021.105638>
- Hygnsstrom SE, Timm RM, Larsson GE (1994) Prevention and Control of Wildlife Damage. University of Nebraska, Lincoln
- IPMcoalition (2023) Coalition database. Integrated Pest Management (IPM) Coalition. <https://www.ipm-coalition.org/index.php/facet-database>. Accessed 30 November 2023
- Isackson B, Irizarry L (2023) Rodenticide Toxicity. StatPearls Publishing. <https://www.ncbi.nlm.nih.gov/books/NBK554428/>. Accessed 18 October 2023
- Jacob J, Buckle A (2018) Use of Anticoagulant Rodenticides in Different Applications Around the World. In: van den Brink NW, Elliott JE, Shore RF, Rattner BA (eds) *Anticoagulant Rodenticides and Wildlife*. Springer International Publishing, Cham, pp 11–43. https://doi.org/10.1007/978-3-319-64377-9_2
- Jacobinnert K, Hinds L, Zhang Z (2021) The status of fertility control for rodents—recent achievements and future 2 directions. *Integrative Zool* 17:964–980
- Jäkel T, Khoprasert Y, Promkerd P, Hongnark S (2006) An experimental field study to assess the effectiveness of bait containing the parasitic protozoan *Sarcocystis singaporensis* for protecting rice crops against rodent damage. *Crop Prot* 25:773–780. <https://doi.org/10.1016/j.cropro.2005.10.008>
- Jäkel T, Promkerd P, Sitthirath R, Guedant P, Khoprasert Y (2019) Biocontrol of rats in an urban environment in Southeast Asia using *Sarcocystis singaporensis*. *Pest Manag Sci* 75:2148–2157. <https://doi.org/10.1002/ps.5335>
- Jassat W, Naicker N, Naidoo S, Mathee A (2013) Rodent control in urban communities in Johannesburg, South Africa: from research to action. *Int J Environ Health Res* 23:474–483. <https://doi.org/10.1080/09603123.2012.755156>
- Jeyapal S, Sagar H (2020) A true killer among us: a case series of yellow phosphorus poisoning in children at a tertiary care center in South India. *Journal of Pediatric Critical Care* 7:198
- Kaine G, Stronge D, Wright V (2023) Predicting people's motivation to engage in urban predator control. *Urban Ecosyst*. <https://doi.org/10.1007/s11252-023-01430-z>
- Karunarathne A, Bhalla A, Sethi A, Perera U, Eddleston M (2021) Importance of pesticides for lethal poisoning in India during 1999 to 2018: a systematic review. *BMC Public Health* 21:1441. <https://doi.org/10.1186/s12889-021-11156-2>
- Kazasidis O, Jacob J (2023) Machine learning identifies straightforward early warning rules for human Puumala hantavirus outbreaks. *Sci Rep* 13:3585. <https://doi.org/10.1038/s41598-023-30596-x>
- Klemann N, Walther B, Matuschka F-R, Jacob J, Endepols S (2023) The stop-feed effect of cholecalciferol (vitamin D3) and the efficacy of brodifacoum combined with cholecalciferol in Y139C-resistant Norway rats (*Rattus norvegicus*). *J Pest Sci* 96:1331–1340. <https://doi.org/10.1007/s10340-023-01600-0>
- Knight MW (2013) Chapter 85 - Zinc Phosphide. In: Peterson ME, Talcott PA (eds) *Small Animal Toxicology* (Third Edition). W.B. Saunders, Saint Louis, pp 853–864. <https://doi.org/10.1016/B978-1-4557-0717-1.00085-5>
- Knipe DW, Chang SS, Dawson A, Eddleston M, Konradsen F, Metcalfe C, Gunnell D (2017) Suicide prevention through means restriction: impact of the 2008–2011 pesticide restrictions on suicide in Sri Lanka. *PLoS ONE* 12:e0172893. <https://doi.org/10.1371/journal.pone.0172893>
- Kotthoff M, Rüdel H, Jüriling H, Severin K, Hennecke S, Friesen A, Koschorreck J (2019) First evidence of anticoagulant rodenticides in fish and suspended particulate matter: spatial and temporal distribution in German freshwater aquatic systems. *Environ Sci Pollut Res* 26:7315–7325. <https://doi.org/10.1007/s11356-018-1385-8>
- Krebs CJ, Kenney AJ, Singleton GR, Mutze G, Pech RP, Brown PR, Davis SA (2004) Can outbreaks of house mice in south-eastern Australia be predicted by weather models? *Wildl Res* 31:465–474. <https://doi.org/10.1071/WR03131>
- Krijger IM, Gort G, Belmain SR, Groot Koerkamp PWG, Shafali RB, Meerburg BG (2020) Efficacy of management and monitoring methods to prevent post-harvest losses caused by rodents. *Animals* (basel). <https://doi.org/10.3390/ani10091612>
- Labuschagne L, Swanepoel LH, Taylor PJ, Belmain SR, Keith M (2016) Are avian predators effective biological control agents for rodent pest management in agricultural systems? *Biol Control* 101:94–102. <https://doi.org/10.1016/j.biocontrol.2016.07.003>
- Lambropoulos AS, Fine JB, Perbeck A, Torres D, Glass GE, McHugh P, Dorsey EA (1999) Rodent control in urban areas: an interdisciplinary approach. *J Environ Health* 6:12
- Lee MJ, Byers KA, Donovan CM, Bidulka JJ, Stephen C, Patrick DM, Himsforth CG (2018) Effects of culling on *Leptospira interrogans* carriage by rats. *Emerg Infect Dis* 24:356–360. <https://doi.org/10.3201/eid2402.171371>
- Lee MJ, Byers KA, Stephen C, Patrick DM, Corrigan R, Iwasawa S, Himsforth CG (2022) Reconsidering the “war on rats”: what we know from over a century of research into municipal rat management. *Front Ecol Evol*. <https://doi.org/10.3389/fevo.2022.813600>
- Leirs H, Verhagen R, Verheyen W, Mwanjabe P, Mbise T (1996) Forecasting rodent outbreaks in Africa: an ecological basis for Mاستomys control in Tanzania. *J Appl Ecol* 33:937–943
- Lekei E, Ngowi AV, London L (2017) Acute pesticide poisoning in children: hospital review in selected hospitals of Tanzania. *J Toxicol* 2017:4208405. <https://doi.org/10.1155/2017/4208405>
- Lettoof DC, Lohr MT, Busetti F, Bateman PW, Davis RA (2020) Toxic time bombs: Frequent detection of anticoagulant rodenticides in urban reptiles at multiple trophic levels. *Sci Total Environ* 724:138218. <https://doi.org/10.1016/j.scitotenv.2020.138218>
- Lewis KA, Tziliavakis J, Warner DJ, Green A (2016) An international database for pesticide risk assessments and management. *Hum Ecol Risk Assess Int J* 22:1050–1064. <https://doi.org/10.1080/10807039.2015.1133242>
- López-Perea JJ, Mateo R (2018) Secondary exposure to anticoagulant rodenticides and effects on predators. In: van den Brink NW, Elliott JE, Shore RF, Rattner BA (eds) *Anticoagulant rodenticides and wildlife*. Springer International, Cham, pp 159–193
- Lund M (1988) Ultrasound devices. In: Prakash I (ed) *Rodent Pest Management*. CRC Press. <https://doi.org/10.1201/9781351076456>
- Mahlaba TA, Monadjem A, McCleery R, Belmain SR (2017) Domestic cats and dogs create a landscape of fear for pest rodents around rural homesteads. *PLoS ONE* 12:e0171593. <https://doi.org/10.1371/journal.pone.0171593>
- Marashi SM (2015) What really happens after zinc phosphide ingestion? A debate against the current proposed mechanism of phosphine liberation in zinc phosphide poisoning. *Eur Rev Med Pharmacol Sci* 19:4210–4211
- Mari Saez A, Cherif Haidara M, Camara A, Kourouma F, Sage M, Magassouba N, Fichet-Calvet E (2018) Rodent control to fight Lassa fever: evaluation and lessons learned from a 4-year study in Upper Guinea. *PLoS Negl Trop Dis* 12:e0006829. <https://doi.org/10.1371/journal.pntd.0006829>
- Masi E, Pino FA, Santos MdGS, Genehr L, Albuquerque JOM, Bancher AM, Alves JCM (2010) Socioeconomic and environmental risk

- factors for urban rodent infestation in Sao Paulo, Brazil. *J Pest Sci* 83:231–241. <https://doi.org/10.1007/s10340-010-0290-9>
- Massawe AW, Makundi RH, Zhang Z, Mhamphi G, Liu M, Li H-J, Belmain SR (2018) Effect of synthetic hormones on reproduction in *Mastomys natalensis*. *J Pest Sci* 91:157–168. <https://doi.org/10.1007/s10340-017-0894-4>
- Massei G, Jacob J, Hinds LA (2023) Developing fertility control for rodents: a framework for researchers and practitioners. *Integr Zool*. <https://doi.org/10.1111/1749-4877.12727>
- Md Meftaul I, Venkateswarlu K, Dharmarajan R, Annamalai P, Megharaj M (2020) Pesticides in the urban environment: A potential threat that knocks at the door. *Sci Total Environ* 711:134612. <https://doi.org/10.1016/j.scitotenv.2019.134612>
- Mdangi M, Mulungu LS, Massawe AW, Eiseb SJ, Tutjavi V, Kirsten F, Mahlaba T, Malebane P, von Maltitz E, Monadjem A, Dlamini N, Makundi RH, Belmain SR (2013) Assessment of rodent damage to stored maize (*Zea mays*L.) on smallholder farms in Tanzania. *Int J Pest Manag* 59:55–62. <https://doi.org/10.1080/09670874.2012.744495>
- Meehan AP (1984) Rats and Mice: Their Biology and Control. Rentokil Ltd., Felcourt, East Grinstead
- Meheretu Y, Kiros W, Deckers S, Raes D, Makundi R, Leirs H (2010) Farmers' perspectives of rodent damage and management from the highlands of Tigray, Northern Ethiopian. *Crop Prot* 29:532–539. <https://doi.org/10.1016/j.cropro.2009.12.006>
- Meheretu Y, Fanos A, Dawit K, Kiros W (2022) Stored-grain losses and management practices among smallholder cereal farmers in Northern Ethiopia. *Int J Pest Manag* 68:59–68. <https://doi.org/10.1080/09670874.2020.1795301>
- Murali R, Bhalla A, Singh D, Singh S (2009) Acute pesticide poisoning: 15 years experience of a large North-West Indian hospital. *Clin Toxicol* 47:35–38. <https://doi.org/10.1080/15563650701885807>
- NAFDAC (2022) Public Alert No. 009/2022 – warning against the use of aluminium phosphide 560TB pesticides. National agency for food and drug administration and control. <https://www.nafdac.gov.ng/public-alert-no-009-2022-warning-against-the-use-of-aluminium-phosphide-560tb-pesticides/>. Accessed 30 November 2023
- Nayak MK, Daglish GJ, Phillips TW, Ebert PR (2020) Resistance to the fumigant phosphine and its management in insect pests of stored products: a global perspective. *Annu Rev Entomol* 65:333–350. <https://doi.org/10.1146/annurev-ento-011019-025047>
- OECD (2014) Report of the OECD seminar on risk reduction through prevention, detection and control of the illegal International trade in agricultural pesticides. <https://doi.org/10.1787/9789264221574-en>
- Ostfeld RS, Holt RD (2004) Are predators good for your health? Evaluating evidence for top-down regulation of zoonotic disease reservoirs. *Front Ecol Environ* 2:13–20. [https://doi.org/10.1890/1540-9295\(2004\)002\[0013:APGFYH\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2004)002[0013:APGFYH]2.0.CO;2)
- PAN (2021) PAN International list of highly hazardous pesticides. Pesticide action network International, Hamburg, Germany
- PAN (2022) PAN International consolidated list of banned pesticides. 6th edn. Pesticide action network International Penang, Malaysia
- Panis C, Kawassaki ACB, Crestani APJ, Pascotto CR, Bortoloti DS, Vicentini GE, Lucio LC, Ferreira MO, Prates RTC, Vieira VK, Gaboardi SC, Candiotto LZZP (2021) Evidence on human exposure to pesticides and the occurrence of health hazards in the Brazilian population: a systematic review. *Front Public Health* 9:787438. <https://doi.org/10.3389/fpubh.2021.787438>
- Papini S, Nakagawa LE (2014) Current status of rodenticide intoxication in Brazil: a preliminary survey from 2009 to 2011. *Braz Arch Biol Technol* 57:685–688. <https://doi.org/10.1590/s1516-8913201402370>
- Papini S, de Masi E, Nakagawa LE, de Oliveira JCB (2019) Rodenticide efficacy in sewers in São Paulo, Brazil. *Environ Sci Pollut Res Int* 26:35270–35274. <https://doi.org/10.1007/s11356-019-06802-7>
- Parsons MH, Banks PB, Deutsch MA, Munshi-South J (2018) Temporal and space-use changes by rats in response to predation by feral cats in an urban ecosystem. *Front Ecol Evolution*. <https://doi.org/10.3389/fevo.2018.00146>
- Permentier K, Vercammen S, Soetaert S, Schellekens C (2017) Carbon dioxide poisoning: a literature review of an often forgotten cause of intoxication in the emergency department. *Int J Emerg Med* 10:14. <https://doi.org/10.1186/s12245-017-0142-y>
- Pertile AC, Lustosa R, Carvalho-Pereira T, Pedra GG, Panti-May JA, Oliveira U, Zeppelini CG, Souza FN, Oliveira DS, Khalil H, Reis MG, Childs J, Ko AI, Begon M, Costa F (2022) Evaluation of the impact of chemical control on the ecology of *Rattus norvegicus* of an urban community in Salvador, Brazil *Plos One* 17:e0270568. <https://doi.org/10.1371/journal.pone.0270568>
- Peshin SS, Srivastava A, Halder N, Gupta YK (2014) Pesticide poisoning trend analysis of 13 years: a retrospective study based on telephone calls at the national poisons information centre, All India Institute of Medical Sciences, New Delhi. *J Forensic Leg Med* 22:57–61. <https://doi.org/10.1016/j.jflm.2013.12.013>
- Peterson AC, Ghersi BM, Campanella R, Riegel C, Lewis JA, Blum MJ (2020) Rodent assemblage structure reflects socioecological mosaics of counter-urbanization across post-Hurricane Katrina New Orleans. *Landsc Urb Plan*. <https://doi.org/10.1016/j.landurbplan.2019.103710>
- PPD (2022) Registered pesticide lists up to date plant protection division (PPD), Department of Agriculture, Myanmar
- Proudfoot AT (2009) Aluminium and zinc phosphide poisoning. *Clin Toxicol (phila)* 47:89–100. <https://doi.org/10.1080/15563650802520675>
- Pyzyńska B, Whish S, Dyer CA, Mayer LP, Witmer G, Moulton R (2016) Free ranging wild-caught Norway rats have reduced fecundity after consuming liquid oral fertility bait containing 4-vinylcyclohexene diepoxide and triptolide. In: Proceedings of the vertebrate pest conference, pp 314–316 <https://doi.org/10.5070/v427110405>
- Rahelinirina S, Rahajandraibe S, Rakotosamimanana S, Rajerison M (2023) Assessing the effectiveness of intervention to prevent plague through community and animal-based survey. *PLOS Glob Public Health* 3:e0002211. <https://doi.org/10.1371/journal.pgph.0002211>
- Regnery J., Friesen, A., Geduhn, A., Göckener, B., Kotthoff, M., Parrhysius, P., Petersohn, E., Reifferscheid, G., Schmolz, E., Schulz, R.S., Schwarzbauer, J., Brinke, M., 2019a. Rating the risks of anticoagulant rodenticides in the aquatic environment: a review. *Environ. Chem. Lett.* 17(1):215–240. <https://doi.org/10.1007/s10311-018-0788-6>
- Regnery J, Schulz RS, Parrhysius P, Bachtin J, Brinke M, Schaefer S, Reifferscheid G, Friesen A (2020) Heavy rainfall provokes anticoagulant rodenticides' release from baited sewer systems and outdoor surfaces into receiving streams. *Sci Total Environ* 740:139905. <https://doi.org/10.1016/j.scitotenv.2020.139905>
- Richardson JL, Burak MK, Hernandez C, Shirvell JM, Mariani C, Carvalho-Pereira TS, Pertile AC, Panti-May JA, Pedra GG, Serano S, Taylor J, Carvalho M, Rodrigues G, Costa F, Childs JE, Ko AI, Caccone A (2017) Using fine-scale spatial genetics of Norway rats to improve control efforts and reduce leptospirosis risk in urban slum environments. *Evol Appl* 10:323–337. <https://doi.org/10.1111/eva.12449>
- Roomaney R, Ehrlich R, Rother HA (2012) The acceptability of rat trap use over pesticides for rodent control in two poor urban communities in South Africa. *Environ Health* 11:32. <https://doi.org/10.1186/1476-069X-11-32>

- Roszkowski AP (1965) The pharmacological properties of norbormide, a selective rat toxicant. *J Pharmacol Exp Ther* 149:288–299
- Rother H-A (2016) Pesticide vendors in the informal sector: trading health for income. *New Solut: A J Environ Occup Health Policy* 26:241–252. <https://doi.org/10.1177/1048291116651750>
- Schlötelburg A, Geduhn A, Schmolz E, Friesen A, Baker S, Martenson N, Laidier GL, Urzinger M, Klute O, Schröer D, Brigham A, Puschmann M (2021) NoCheRo-Guidance for the Evaluation of Rodent Traps. Part A Break back/Snap traps. . German Environment Agency, Dessau, Germany
- Selemani M, Makundi RH, Massawe AW, Mhamphi G, Mulungu LS, Belmain SR (2021) Impact of contraceptive hormones on the reproductive potential of male and female commensal black rats (*Rattus rattus*). *Integr Zool*. <https://doi.org/10.1111/1749-4877.12563>
- Shadnia S, Mehrpour O, Soltaninejad K (2010) A simplified acute physiology score in the prediction of acute aluminum phosphide poisoning outcome. *Indian J Med Sci* 64:532–539
- Sherley M (2004) The traditional categories of fluoroacetate poisoning signs and symptoms belie substantial underlying similarities. *Toxicol Lett* 151:399–406. <https://doi.org/10.1016/j.toxlet.2004.03.013>
- Shiels AB, Runte J, Ruell EW, Eckery DC, Witmer GW, Salkeld DJ (2023) Treatment with the immunocontraceptive vaccine, Gona-Con, induces temporary fertility control in free-ranging prairie dog populations in Colorado, USA. *Wildl Res*. <https://doi.org/10.1071/WR22135>
- Shore RF, Coeurdassier M (2018) Primary exposure and effects in non-target animals. In: van den Brink NW, Elliott JE, Shore RF, Rattner BA (eds) *Anticoagulant rodenticides and wildlife*. Springer International Publishing, Cham, pp 135–157. https://doi.org/10.1007/978-3-319-64377-9_6
- Shuster SM, Pyzyna B, Ray C, Mayer LP (2023) The demographic consequences of fertility reduction in rats and voles. *J Pest Sci* 96:1313–1329. <https://doi.org/10.1007/s10340-023-01607-7>
- Siers SR, Pyzyna BR, Mayer L, Leinbach CDIL, Sugihara RT, Witmer GW (2017) Laboratory evaluation of the Effectiveness of the fertility control bait ContraPest® on wild-captured black rats (*Rattus rattus*). In: *Proceedings of the Vertebrate Pest Conference*, pp 1–7
- Singleton GR, Lorica RP, Htwe NM, Stuart AM (2021) Rodent management and cereal production in Asia: balancing food security and conservation. *Pest Manag Sci*. <https://doi.org/10.1002/ps.6462>
- Soleng A, Edgar KS, von Krogh A, Seljetun KO (2022) Suspected rodenticide exposures in humans and domestic animals: Data from inquiries to the Norwegian Poison Information Centre, 2005–2020. *PLoS ONE* 17:e0278642. <https://doi.org/10.1371/journal.pone.0278642>
- Soni JP, Ghormade PS, Akhade S, Chavali K, Sarma B (2020) A fatal case of multi-organ failure in acute yellow phosphorus poisoning. *Autops Case Rep* 10:e2020146. <https://doi.org/10.4322/acr.2020.146>
- Staples L, Smith M, Pontin K, Hunt W Use of zinc phosphide to overcome rodent infestations. In: *Stored grain in Australia 2003*. In: *Proceedings of the Australian postharvest technical conference*, Canberra, 2003.
- Stuart AM, Kong P, Then R, Flor RJ, Sathya K (2020) Tailor-made solutions to tackle rodent pests of rice through community-based management approaches in Cambodia. *Crop Prot* 135:104717. <https://doi.org/10.1016/j.cropro.2019.01.023>
- Stuart AM, Merfield CN, Horgan FG, Willis S, Watts MA, Ramirez-Munoz F, Jorge-Sánchez U, Utyasheva L, Eddleston M, Davis ML, Neumeister L, Sanou MR, Williamson S (2023) Agriculture without paraquat is feasible without loss of productivity-lessons learned from phasing out a highly hazardous herbicide. *Environ Sci Pollut Res Int* 30:16984–17008. <https://doi.org/10.1007/s11356-022-24951-0>
- Swanepoel LH, Swanepoel CM, Brown PR, Eiseb SJ, Goodman SM, Keith M, Kirsten F, Leirs H, Mahlaba TA, Makundi RH, Malebane P, von Maltitz EF, Massawe AW, Monadjem A, Mulungu LS, Singleton GR, Taylor PJ, Soarimalala V, Belmain SR (2017) A systematic review of rodent pest research in Afro-Malagasy small-holder farming systems: are we asking the right questions? *PLoS ONE* 12:e0174554. <https://doi.org/10.1371/journal.pone.0174554>
- Tang T, Qian K, Shi T, Wang F, Li P, Li J, Cao Y (2012a) Photodegradation of quinetrol in waters and the transformation products by UV irradiation. *Chemosphere* 89:1419–1425. <https://doi.org/10.1016/j.chemosphere.2012.06.011>
- Tang T, Shi T, Li D, Xia J, Hu Q, Cao Y (2012b) Adsorption properties and degradation dynamics of endocrine-disrupting chemical levonorgestrel in soils. *J Agric Food Chem* 60:3999–4004. <https://doi.org/10.1021/jf300479z>
- Taylor PJ, Downs S, Monadjem A, Eiseb SJ, Mulungu LS, Massawe AW, TaA M, Kirsten F, Von Maltitz E, Malebane P, Makundi RH, Lamb J, Belmain SR (2012) Experimental treatment-control studies of ecologically based rodent management in Africa: balancing conservation and pest management. *Wildl Res* 39:51. <https://doi.org/10.1071/wr11111>
- Trakulsrichai S, Kosanyawat N, Atikawedparit P, Sriapha C, Tongpoo A, Udomsubpayakul U, Rittilert P, Wanankul W (2017) Clinical characteristics of zinc phosphide poisoning in Thailand. *Ther Clin Risk Manag* 13:335–340. <https://doi.org/10.2147/tcrms.129610>
- UCT (2019) Low danger pest control booklet for communities. Division of environmental health, University of Cape Town (UCT), South Africa
- UFAW (2021) Humane management of rats, mice and moles detailed advice. The International Animal Welfare Science Society. <https://www.ufaw.org.uk/rodent-control/humane-rodent-control-detailed-advice>. Accessed 30 November 2023
- UNEP, WHO, FAO (2022) Chapter 6/12: Current pesticide risk reduction and risk management. In: UNEP, WHO, FAO (eds) *Environmental and health impacts of pesticides and fertilizers and ways of minimizing them*. United Nations Environment Programme, World Health Organization, & Food and Agriculture Organization of the United Nations, p 20
- UNEP, WHO, ILO (1988) Phosphine and selected metal phosphides. vol 73.
- Utyasheva L, Eddleston M (2021) Prevention of pesticide suicides and the right to life: the intersection of human rights and public health priorities. *J Hum Rights* 20:52–71. <https://doi.org/10.1080/14754835.2020.1850241>
- Vandenbroucke V, Desmet N, De Backer P, Croubels S (2008) Multi-residue analysis of eight anticoagulant rodenticides in animal plasma and liver using liquid chromatography combined with heated electrospray ionization tandem mass spectrometry. *J Chromatogr B: Anal Technol Biomed Life Sci* 869:101–110. <https://doi.org/10.1016/j.jchromb.2008.05.011>
- Varghese P, Erickson TB (2022) Pesticide poisoning among children in India: the need for an urgent solution. *Glob Pediatr Health* 9:2333794X221086577. <https://doi.org/10.1177/2333794X221086577>
- Walther B, Geduhn A, Schenke D, Jacob J (2021) Exposure of passerine birds to brodifacoum during management of Norway rats on farms. *Sci Total Environ* 762:144160. <https://doi.org/10.1016/j.scitotenv.2020.144160>
- Walther B, Bohot A, Ennen H, Beilmann P, Schaper O, Hantschke P, Werdin S, Jacob J (2024) Technical assessment of mechanical

- and electronic traps to facilitate future improvements in trap efficacy and humaneness. *Pest Manag Sci.* <https://doi.org/10.1002/ps.8011>
- WHO (1972) Vector Control in International Health. World Health Organisation, Geneva
- WHO (2021) Live life: an implementation guide for suicide prevention in countries. World Health Organization, Geneva
- Witmer GW (2005) Wildlife population monitoring: some practical considerations. *Wildl Res* 32:259–263. <https://doi.org/10.1071/WR04003>
- Witmer GW (2019) The changing role of rodenticides and their alternatives in the management of commensal rodents. *Hum-Wildl Interact* 13:186–189
- Yan Y, Jiang Y, Liu R, Eddleston M, Tao C, Page A, Wang L, Feng G, Liu S (2023) Impact of pesticide regulations on mortality from suicide by pesticide in China: an interrupted time series analysis. *Front Psychiatry* 14:1189923. <https://doi.org/10.3389/fpsyt.2023.1189923>
- Yogendranathan N, Herath H, Sivasundaram T, Constantine R, Kulatunga A (2017) A case report of zinc phosphide poisoning: complicated by acute renal failure and tubulo interstitial nephritis. *BMC Pharmacol Toxicol* 18:37. <https://doi.org/10.1186/s40360-017-0144-7>
- Zeppelini CG, Carvalho-Pereira T, Alves RS, Santiago DCC, Santo VFE, Begon M, Costa F, Khalil H (2021) Demographic drivers of Norway rat populations from urban slums in Brazil. *Urban Ecosyst* 24:801–809. <https://doi.org/10.1007/s11252-020-01075-2>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.