



OPEN ACCESS

EDITED AND REVIEWED BY

Eva Sintes,
Spanish Institute of Oceanography
(IEO), Spain

*CORRESPONDENCE

Sophie Bonnet
✉ sophie.bonnet@mio.osupytheas.fr
Cecile Guieu
✉ cecile.guieu@imev-mer.fr

RECEIVED 10 September 2025

ACCEPTED 22 September 2025

PUBLISHED 09 October 2025

CITATION

Bonnet S, Guieu C, Berman-Frank I,
Capone DG, Fitzsimmons JN and Resing JA
(2025) Editorial: Hydrothermal and submarine
volcanic activity: impacts on ocean chemistry
and plankton dynamics.
Front. Microbiol. 16:1703123.
doi: 10.3389/fmicb.2025.1703123

COPYRIGHT

© 2025 Bonnet, Guieu, Berman-Frank,
Capone, Fitzsimmons and Resing. This is an
open-access article distributed under the
terms of the [Creative Commons Attribution
License \(CC BY\)](#). The use, distribution or
reproduction in other forums is permitted,
provided the original author(s) and the
copyright owner(s) are credited and that the
original publication in this journal is cited, in
accordance with accepted academic practice.
No use, distribution or reproduction is
permitted which does not comply with these
terms.

Editorial: Hydrothermal and submarine volcanic activity: impacts on ocean chemistry and plankton dynamics

Sophie Bonnet^{1*}, Cecile Guieu^{2*}, Ilana Berman-Frank³,
Douglas G. Capone⁴, Jessica Nicole Fitzsimmons⁵ and
Joseph A. Resing^{6,7}

¹Aix Marseille University, Université de Toulon, CNRS, IRD, MIO, Marseille, France, ²Sorbonne Université, Centre National de la Recherche Scientifique, Laboratoire d'Océanographie de Villefranche, LOV, Villefranche-sur-Mer, France, ³Department of Marine Biology, Leon H. Charney School of Marine Sciences, University of Haifa, Haifa, Israel, ⁴Department of Biological Sciences, University of Southern California, Los Angeles, CA, United States, ⁵Texas A&M University, College Station, TX, United States, ⁶Cooperative Institute for Climate, Ocean, and Ecosystem Studies, University of Washington, Seattle, WA, United States, ⁷NOAA-Pacific Marine Environmental Laboratory, Seattle, WA, United States

KEYWORDS

hydrothermal activity, iron, plankton, trace metals, ocean

Editorial on the Research Topic

[Hydrothermal and submarine volcanic activity: impacts on ocean chemistry and plankton dynamics](#)

Iron (Fe) is a key limiting factor for primary productivity across vast regions of the global ocean (Tagliabue et al., 2017; Browning and Moore, 2023). Traditionally, atmospheric dust deposition and continental margin fluxes have been considered the dominant sources of Fe to surface waters (Moore et al., 2004). However, deep-sea hydrothermal systems also release substantial amounts of dissolved Fe (dFe) into the overlying water column via hydrothermal plumes. These plumes can be transported over long distances across ocean basins, thereby influencing large-scale ocean chemistry (Nishioka et al., 2013; Saito et al., 2013; Fitzsimmons et al., 2014; Resing et al., 2015). Furthermore, global biogeochemical models estimate that hydrothermal fluxes from mid-ocean ridges contribute nearly 23% of the dFe inventory in the oceanic water column (Tagliabue et al., 2017).

Unlike deep hydrothermal systems associated with mid-ocean ridges (>2,000 m), hydrothermal activity can also occur at much shallower depths, such as in proximity to island arcs and hotspot volcanoes, thereby delivering substantial fluxes of trace metals to the upper ocean waters (ca. 500–1,000 m; Massoth et al., 2007; Hawkes et al., 2014) and in some cases directly to the euphotic zone (0–150 m; Chemine et al., 1991; Resing et al., 2009; Santana-Casiano et al., 2016; Guieu et al., 2018; Tilliette et al., 2022). However, unlike deep hydrothermal plumes, dFe concentrations in these shallow environments decrease rapidly with distance from the source due to the much stronger water mass dynamics that prevail at shallow depths (Tilliette et al., 2022). In deep hydrothermal systems, particulate dFe losses are mitigated through stabilization with organic Fe-binding ligands (Bennett et al., 2008; Toner et al., 2009). In contrast, in shallow hydrothermal systems, both the concentration

and chemical nature of these ligands are poorly characterized, thereby constraining our ability to assess Fe stabilization, residence times, and bioavailability in these environments.

This Research Topic includes, among other contributions, several studies investigating the occurrence, composition, and functional role of ligands in stabilizing Fe near the shallow hydrothermal vents of the Tonga-Kermadec arc (Western subtropical South Pacific). Mahieu et al. identified elevated conditional concentrations of Fe-binding ligands peaking near hydrothermal sites, primarily composed of intermediate-strength L2 types. Despite this, their analysis revealed that ligand concentrations were largely in excess relative to DFe, suggesting limited effectiveness in stabilizing hydrothermal Fe inputs. Consistent with this finding, Portlock et al. reported unusually high concentrations of reduced sulfur substances (phytoplankton-derived biomolecules that associate with trace metals under elevated exposure levels), thereby mitigating toxicity. Complementing these studies, Dulaquais et al. investigated the contribution of soluble humic-like substances produced during phytoplankton degradation to Fe complexation at various sites impacted differently by hydrothermal fluids. The authors showed that the humic-ligands were unsaturated likely due to their inability to access colloidal DFe, Fe(II) and FeOx species. The humic ligands only complexed 1–5.5% of the total DFe pool close to the vent, thereby solubilizing only a small fraction of the hydrothermal Fe released. Beyond Fe, hydrothermal vents release a variety of trace elements and gases. Zhao et al. investigated the impact of hydrothermal activity on the barium cycle. They showed that Ba isotopes in vent waters and sediments are lighter than those in the water column, indicating the preferential removal of lighter isotopes during fluid–seawater mixing and highlighting their value as tracers of hydrothermal influence on sediments. In a shallow Southern Ocean bay, Belyaev et al. demonstrated how hydrothermal activity alters seawater biogeochemistry by linking vent inputs with elevated concentrations of vanadium, cobalt, nickel and Fe, along with methane and carbon dioxide. In addition to the continuous hydrothermal inputs, sporadic volcanic eruptions in many of these systems also deliver material and chemical elements. Chavagnac et al. showed that strontium (Sr) and lithium (Li) isotopic signatures can distinguish between volcanic and hydrothermal sources. Volcanic activity boosts the total Sr and Li through ash dissolution, while hydrothermal inputs drive concurrent increases of both elements in the water column.

These studies suggest that, although shallow hydrothermal systems provide substantial amounts of Fe and other trace elements, limited ligand stabilization may restrict their persistence in surface waters. This, in turn, raises important questions about the residence time, bioavailability, and ecological impact of hydrothermal Fe on plankton communities.

How do these shallow hydrothermal environments (<500 m deep) impact phytoplankton dynamics? The proximity of Fe release to the euphotic layer greatly increases the likelihood of a direct and immediate biological response. Evidence from the Tonga–Kermadec arc indicates that Fe-rich fluids entering surface waters can stimulate intense diazotrophic activity, thereby fueling enhanced net community production, N₂

fixation, and ultimately increased carbon export to the deep ocean (Bonnet et al., 2023). Strikingly, the carbon sequestration efficiency observed in these natural, shallow-vent systems is higher than that of artificial, mesoscale Fe-fertilization experiments, underscoring the ecological significance of shallow hydrothermal inputs as natural “fertilization hotspots” for surface ecosystems.

This Research Topic brings together several complementary studies that explore the complex biological responses elicited by the diverse suite of chemical elements supplied through hydrothermal inputs. Tilliette et al. demonstrated experimentally that increasing fluid inputs initially exerts toxic effects on planktonic communities, but subsequently stimulates net community production, N₂ fixation, and enhanced export relative to controls. Consistent with the findings of Portlock et al., this fertilization effect is likely sustained by planktonic detoxification via thiol-based ligands that bind toxic trace metals (e.g., Cu, Cd, and Hg). Mériguet et al. provided *in situ* evidence that these shallow hydrothermal inputs structure the entire ecosystem’s trophic dynamics. Using imaging and acoustic approaches, the authors demonstrated that the elevated diazotrophic biomass stimulated by hydrothermal fluids propagates through meso- and macrozooplankton communities, which in turn may enhance organic matter export via the production of fast-sinking fecal pellets. Consistent with this, Ababou et al. showed that zooplankton-derived material (fecal aggregates, carcasses, and cylindrical fecal pellets) represents more than 90% of the carbon flux exported below the euphotic zone in the vicinity of hydrothermal vents. In parallel, nitrogen isotope budgets analyzed by Forrer et al. revealed that this export is predominantly fueled by diazotrophy, which requires high Fe availability.

Together, these findings indicate that, in this system, newly fixed nitrogen is efficiently transferred up to zooplankton and repackaged into dense fecal pellets. This highlights an overlooked but highly effective pathway by which diazotroph activity indirectly sustains particulate organic carbon export, strengthening the coupling between nitrogen fixation and the efficiency of the biological carbon pump in shallow hydrothermal regions. In the much deeper environments of the Arctic Ocean, Wegener et al. reported a clear biological response within hydrothermal plumes, characterized by elevated carbon fixation rates compared to surrounding waters, suggesting enhanced chemoautotrophy fueled by hydrogen and sulfide as energy sources.

Taken together, the studies reported in this Research Topic reveal that the fate of hydrothermal Fe in shallow systems is governed by a delicate balance between its limited chemical stabilization by organic ligands and rapid biological utilization in the euphotic zone. While the weak role of binding ligands may constrain the long-term persistence of hydrothermal Fe, its immediate availability fuels primary productivity, diazotroph activity and production, trophic transfer, and greater carbon export. This coupling of trace metal chemistry and biological responses underscores the role of shallow hydrothermal systems as natural laboratories where geochemical processes and ecosystem functioning are tightly interconnected, with potentially significant implications for the oceanic carbon and nitrogen cycles.

Author contributions

SB: Conceptualization, Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Project administration, Supervision, Validation, Writing – original draft, Writing – review & editing. CG: Conceptualization, Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Project administration, Supervision, Validation, Writing – review & editing. IB-F: Writing – review & editing. DC: Writing – review & editing. JF: Writing – review & editing. JR: Writing – review & editing.

Funding

The author(s) declare that financial support was received for the research and/or publication of this article. Several articles of this Research Topic are a contribution of the TONGA project (Shallow hydrothermal sources of trace elements: potential impacts on biological productivity and the biological carbon pump; TONGA cruise doi: 10.17600/18000884) funded by the Agence Nationale de la Recherche (grant TONGA ANR-18-CE01-0016 0014), the LEFE-CyBER program (CNRS-INSU), the A-Midex foundation, the Institut de Recherche pour le Développement (IRD). Joseph Resing was funded by NOAA Ocean Exploration and the Pacific Marine Environmental Laboratory. This publication is CICOES # 2025-1484 and PMEL # 5819.

References

- Bennett, S. A., Achterberg, E. P., Connelly, D. P., Statham, P. J., Fones, G. R., and German, C. R. (2008). The distribution and stabilisation of dissolved Fe in deep-sea hydrothermal plumes. *Earth Planet. Sci. Lett.* 270, 157–167. doi: 10.1016/j.epsl.2008.01.048
- Bonnet, S., Guieu, C., Taillandier, V., Boulart, C., Bouruet-Aubertot, P., Gazeau, F., et al. (2023). Natural iron fertilization by shallow hydrothermal sources fuels diazotroph blooms in the ocean. *Science* 380, 812–817. doi: 10.1126/science.abq4654
- Browning, T. J., and Moore, C. M. (2023). Global analysis of ocean phytoplankton nutrient limitation reveals high prevalence of co-limitation. *Nat. Commun.* 14:5014. doi: 10.1038/s41467-023-40774-0
- Chemine, J.-L., Stoffers, P., McMurtry, G., Richnow, H., Puteanus, D., and Sedwick, P. (1991). Gas-rich submarine exhalations during the 1989 eruption of Macdonald Seamount. *Earth Planet. Sci. Lett.* 107, 318–327. doi: 10.1016/0012-821X(91)90079-W
- Fitzsimmons, J. N., Boyle, E. A., and Jenkins, W. J. (2014). Distal transport of dissolved hydrothermal iron in the deep South Pacific Ocean. *Proc. Nat. Acad. Sci. U. S. A.* 111, 16654–16661. doi: 10.1073/pnas.1418778111
- Guieu, C., Bonnet, S., Petrenko, A., Menkes, C., Chavagnac, V., Desboeufs, K., et al. (2018). Iron from a submarine source impacts the productive layer of the Western Tropical South Pacific (WTSP). *Sci. Rep.* 8:9075. doi: 10.1038/s41598-018-27407-z
- Hawkes, J. A., Connelly, D. P., Rijkenberg, M. J. A., and Achterberg, E. P. (2014). The importance of shallow hydrothermal island arc systems in ocean biogeochemistry. *J. Geophys. Res.* 41, 942–947. doi: 10.1002/2013GL058817
- Massoth, G., Baker, E., Worthington, T., Lupton, J., De Ronde, C., Arculus, R., et al. (2007). Multiple hydrothermal sources along the south Tonga arc and Valu Fa Ridge. *Geochim. Geophys. Geosyst.* 8, 1–26. doi: 10.1029/2007GC001675
- Moore, J. K., Doney, S. C., and Lindsay, K. (2004). Upper ocean ecosystem dynamics and iron cycling in a global three-dimensional model. *Glob. Biogeochem. Cycles* 18, 1–21. doi: 10.1029/2004GB002220
- Nishioka, J., Obata, H., and Tsumune, D. (2013). Evidence of an extensive spread of hydrothermal dissolved iron in the Indian Ocean. *Earth Planet. Sci. Lett.* 361, 26–33. doi: 10.1016/j.epsl.2012.11.040
- Resing, J. A., Baker, E. T., Lupton, J. E., Walker, S. L., Butterfield, D. A., Massoth, G. J., et al. (2009). Chemistry of hydrothermal plumes above submarine volcanoes of the Mariana Arc. *Geochim. Geophys. Geosyst.* 10, 1–23. doi: 10.1029/2008GC002141
- Resing, J. A., Sedwick, P. N., German, C. R., Jenkins, W. J., Moffett, J. W., Sohst, B. M., et al. (2015). Basin-scale transport of hydrothermal dissolved metals across the South Pacific Ocean. *Nature* 523, 200–203. doi: 10.1038/nature14577
- Saito, M. A., Noble, A. E., Tagliabue, A., Goepfert, T. J., Lamborg, C. H., and Jenkins, W. J. (2013). Slow-spreading submarine ridges in the South Atlantic as a significant oceanic iron source. *Nat. Geosci.* 6, 775–779. doi: 10.1038/ngeo1893
- Santana-Casiano, J. M., Fraile-Nuez, E., González-Dávila, M., Baker, E., Resing, J., and Walker, S. (2016). Significant discharge of CO₂ from hydrothermalism associated with the submarine volcano of El Hierro Island. *Sci. Rep.* 6:25686. doi: 10.1038/srep25686
- Tagliabue, A., Bowie, A. R., Boyd, P. W., Buck, K. N., Johnson, K. S., and Saito, M. A. (2017). The integral role of iron in ocean biogeochemistry. *Nature* 543, 51–59. doi: 10.1038/nature21058
- Tilliette, C., Taillandier, V., Bouruet-Aubertot, P., Grima, N., Maes, C., Montanes, M., et al. (2022). Dissolved iron patterns impacted by shallow hydrothermal sources along a transect through the Tonga-Kermadec Arc. *Glob. Biogeochem. Cycles* 36:e2022GB007363. doi: 10.1029/2022GB007363
- Toner, B. M., Fakra, S. C., Manganini, S. J., Santelli, C. M., Marcus, M. A., Moffett, J. W., et al. (2009). Preservation of iron (II) by carbon-rich matrices in a hydrothermal plume. *Nat. Geosci.* 2, 197–201. doi: 10.1038/ngeo433

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Generative AI statement

The author(s) declare that no Gen AI was used in the creation of this manuscript.

Any alternative text (alt text) provided alongside figures in this article has been generated by Frontiers with the support of artificial intelligence and reasonable efforts have been made to ensure accuracy, including review by the authors wherever possible. If you identify any issues, please contact us.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.