

## **Checklist of the shore fishes of New Caledonia**

*Ronald FRICKE<sup>1</sup> & Michel KULBICKI<sup>2</sup>*

<sup>1</sup> Ichthyology, Staatliches Museum für Naturkunde, Rosenstein 1, 70191 Stuttgart, Germany

fricke.smns@naturkundemuseum-bw.de

<sup>2</sup> IRD, UR128, Université de Perpignan, 52, Avenue Paul Alduy, 66860 Perpignan Cedex, France

michel.kulbicki@univ-perp.fr

The present checklist includes the fish species known from the upper 100 m of the New Caledonian seas. Some deep-sea fishes which are occasionally found in shallow water (e.g. Loyalty Islands), high sea species which only rarely enter coastal waters, or freshwater fish species which may be found in estuaries, are excluded from this list. The geographical distribution of the shore fishes of New Caledonia is discussed by Kulbicki (in press).

A detailed annotated checklist of all New Caledonian fish species including distribution data, literature references and material lists is in preparation by R. Fricke. In the present checklist of shore fish species, all records which are verified either by museum specimens or by confirmation by revising authors, are included. Families are arranged systematically according to Nelson (2006), and species alphabetically under the family names. Doubtful records are discussed after the family name. The names which have been applied to New Caledonian shore fish species in the literature are either listed as valid species, or as synonyms or misidentifications in parentheses behind the species name. In the checklist, reference is given to materials in the collections of the Australian Museum Sydney (AMS), the Muséum National d'Histoire Naturelle Paris (MNHN), and the Staatliches Museum für Naturkunde Stuttgart (SMNS), in order to document new records.

The present checklist of the shore fishes of New Caledonia includes a total of 1694 species. Among these, 85 species are recorded from New Caledonia for the first time. As there is still a number of additional undescribed species known to the authors, and as the New Caledonian fish fauna is still incompletely known, with gaps of knowledge mainly around northern Grande Terre and the northern reefs, the Grand Récif Sud, the Ile des Pins, some of the Loyalty Islands, and the islands of Walpole, Matthew and Hunter, additional species are expected to be added to the list in future.

The 10 largest shore fish families are the Gobiidae (153 species, 9.0 % of the total species), Labridae (113 species, 6.7 %), Pomacentridae (110 species, 6.5 %), Serranidae (85 species, 5.0 %), Apogonidae (78 species, 4.6 %), Blenniidae (61 species, 3.6 %), Muraenidae (52 species, 3.1 %), Scorpaenidae (43 species, 2.5 %), Syngnathidae (42 species, 2.5 %), and Carangidae (40 species, 2.4 %). Among the shore fishes of New Caledonia, 39 species (2.3 % of the total shore fish species) are endemic to the territory. The reason for this relatively low endemism rate is that neighbouring island groups like Lord Howe Island or southern Vanuatu are relatively close, so that most species could easily disperse there in the past. If species living in deeper water (100-600 m) were included in the checklist, the figure of endemic species would be much higher. But even for shore fish species, the percentage of endemic species is expected to rise in the future as several apparently endemic species are still undescribed.

## **REFERENCES**

- BORODIN, N.A. 1932. Scientific results of the Yacht "Alva" world cruise, July, 1931 to March, 1932, in command of William K. Vanderbilt. Fishes. Bulletin of the Vanderbilt Marine Museum, **1** (3): 65-101; Huntington, New York, U.S.A.
- CASTELNAU, F.L. 1873. Contribution to the ichthyology of Australia. Nos. III-IX [with subtitles, indexed as one work]. *Proceedings of the Zoological Acclimatisation Society of Victoria*, **2**: 37-158.
- CASTLE, P.H.J. 1963. The systematics, development and distribution of two eels of the genus *Gnathophis* (Congridae) in Australasian waters. *Zoological Publications of the Victoria University of Wellington*, **34**: 15-47.

- CERRO, L. DEL & LLORIS, D. 1997. Gurnard fishes (Scorpaeniformes, Triglidae) from off New Caledonia, with description of five new species. In: SÉRET, B. (ed.): Résultats des campagnes MUSORSTOM, Volume 17. *Mémoires du Muséum National d'Histoire Naturelle*, **174**: 91-124; Paris.
- FOURMANOIR, P. 1971. Listes des espèces de poissons contenues dans les estomacs de thons jaunes, *Thunnus albacares* (Bonnaterre) 1788 et de thons blancs, *Thunnus alalunga* (Bonnaterre) 1788. *Cahiers de l'O.R.S.T.O.M., Série Océanographie*, **9**(2): 109-118.
- FOURMANOIR, P. 1981. La faune marine du lagon. Pp. 24-29. In: Dugas, F. & Debenay, J.P.: Carte sédimentologique et carte annexe du lagon de Nouvelle-Calédonie à l'1:50.000. Paris (O.R.S.T.O.M.: Notice explicative, 91).
- FOURMANOIR, P. & LABOUTE, P. 1976. Poissons des mers tropicales. Nouvelle-Calédonie. Nouvelles Hébrides. Papeete (Éditions du Pacifique), pp. 1-376.
- FOWLER, H.W. 1928. The fishes of Oceania. *Memoirs of the Bernice P. Bishop Museum*, **10**: iii + 540 pp.
- FRICKE, R. 1999. Fishes of the Mascarene Islands (Réunion, Mauritius, Rodriguez). An annotated checklist, with descriptions of new species. *Koenigstein* (Koeltz Scientific Books), **viii** + 759 pp.
- FRICKE, R. 2000. Callionymidae of New Caledonia, with remarks on related species and descriptions of 10 new species from New Caledonia, Australia, New Guinea, and Hawaii (Teleostei). *Stuttgarter Beiträge zur Naturkunde*, (A) **617**: 1-81; Stuttgart.
- GÜNTHER, A.C.L.G. 1877. Andrew Garrett's Fische der Südsee. Band II, Heft. VI. *Journal des Museum Godeffroy*, **4** (13): 169-216, pls. 101-120.
- GÜNTHER, A.C.L.G. 1909. Andrew Garrett's Fische der Südsee. Heft VIII. *Journal des Museums Godeffroy*, **6** (16): i-iv + 261-388, pls 143-160.
- HARRISON, I.J. & SENOU, H. 1999. Mugilidae. Pp. 2069-2108. In: Carpenter, K.E. & Niem, V.H. (eds.): FAO species identification guide for fishery purposes. The living marine resources of the Western Central Pacific. Volume 4. Bony fishes part 2 (Mugilidae to Carangidae). Rome (FAO), pp. 2069-2790, pls 1-7.
- JOUAN, H. 1863. Supplément à la description des poissons de la Nouvelle-Calédonie. *Mémoires de la Société Impériale des Sciences Naturelles de Cherbourg*, **9**: 177-187.
- JOUAN, H. 1879. Quelques mots sur la faune ichthyologique de la côte nord-est d'Australie et du détroit de Torres, comparée à celle de la Nouvelle-Calédonie. *Mémoires de la Société Impériale des Sciences Naturelles de Cherbourg*, **21**: 328-335.
- KUITER, R.H. 1993. Coastal fishes of south-eastern Australia. University of Hawaii Press, Honolulu.
- KUITER, R.H. 2000. Seahorses, pipefishes and their relatives. A comprehensive guide to Syngnathiformes. Chordleywood, U.K. (TMC Publishing), 240 pp.
- KUITER, R.H. 2002. Lippfische Labridae. Stuttgart (E. Ulmer), 208 pp.
- KULBICKI, M. 1988. Correlation between catch from bottom longlines and fish censuses in the SW lagoon of New Caledonia. *Proceedings of the Sixth International Coral Reef Symposium, Townsville, Australia, 8th-12th August 1988*, **2**: 305-312.
- KULBICKI, M. in press. Biogeography of reef fishes of the French territoires in the South Pacific. *Cybium*.
- KULBICKI, M., RANDALL, J.E. & RIVATON, J. 1994. Checklist of the fishes of the Chesterfield Islands (Coral Sea). *Micronesica*, **27** (1-2): 1-43.
- KULBICKI, M. & WILLIAMS, J.T. 1997. Checklist of the shorefishes of Ouvéa Atoll, New Caledonia. *Atoll Research Bulletin*, **444**: 1-26.
- LABOUTE, P. & GRANDPERRIN, R. 2000. Poissons de Nouvelle-Calédonie. Nouméa (Editions Catherine Ledru), 520 pp.
- LOURIE, S.A., VINCENT, A.C. & HALL, H.J. 1999. Seahorses. An identification guide to the world's species and their conservation. London (Project Seahorse), x + 214 pp.
- MYERS, R.F. 1999. Micronesian reef fishes. A comprehensive guide to the coral reef fishes of Micronesia. 3<sup>rd</sup> revised and expanded edition. Barrigada, Guam (Coral Graphics), vi + 330 pp., 192 pls.
- NELSON, J.S. 2006. Fishes of the world. Fourth edition. Hoboken, New Jersey (John Wiley); xix + 601 pp.
- PARENTI, P. & RANDALL, J.E. 2000. An annotated checklist of the species of the labroid fish families Labridae and Scaridae. *Ichthyological Bulletin, J.L.B. Smith Institute of Ichthyology*, **68**: 1-97.
- PLESSIS, Y. & FOURMANOIR, P. 1966. Mission d'étude des récifs de Nouvelle-Calédonie. Liste des poissons récoltés par Yves Plessis en 1961. *Cahiers du Pacifique*, **9**: 123-147.
- RANDALL, J.E. 2002. Surgeonfishes of the world. Honolulu (Mutual Publishing and Bishop Museum Press), i-x + 1-123.
- RANDALL, J.E. 2005. Reef and shore fishes of the South Pacific. New Caledonia to Tahiti and the Pitcairn Islands. Honolulu (University of Hawai'i Press), p. i-xii + 1-707.
- RANDALL J.E. & KULBICKI M. 2005. *Siganus woodlandi*, new species of rabbitfish (Siganidae) from New Caledonia. *Cybium* **29**(2):185-189
- RANDALL J.E. & KULBICKI M. in press. A review of the goatfishes of the genus *Upeneus* (Perciformes: Mullidae) from New Caledonia and the Chesterfield Bank, with a new species, and four new records. *Zoological Studies* **45** (3) (July 2006).

- RIVATON, J. 1989. Premières observations sur la faune ichtyologique des Iles Chesterfield (Mer du Corail). *Cybium*, 13(2): 139-164.
- RIVATON, J. & BOURRET, P. 1999. Les otolithes des poissons de l'Indo-Pacifique. *Documents Scientifiques et Techniques, I.R.D., Centre de Nouméa, II 2, Volume spécial:* 1-378.
- RIVATON, J., FOURMANOIR, P., BOURRET, P. & KULBICKI, M. 1989. Catalogue des poissons de Nouvelle-Calédonie. Checklist of fishes from New Caledonia. Rapport provisoire. *Catalogues, Sciences de la Mer, Biologie Marine, ORSTOM, Centre de Nouméa:* i-iii + 1-170.
- SCHMELTZ, J.D.E. (ed.) 1869. Museum Godeffroy. Catalog IV, nebst einer Beilage, enthaltend: Beschreibung neuer Bryozoen von Senator Dr. Kirchenpauer zu Hamburg und einer neuen Asteriden-Gattung von Dr. Chr. Lütken zu Kopenhagen. Hamburg (W. Mauke Söhne), i-xliii + 1-139 + 2 unnumbered pp.
- THOLLOT, P. & KULBICKI, M. 1988. Overlap between the fish fauna inventories of coral reefs, soft bottoms and mangroves in Saint-Vincent Bay (New Caledonia). *Proceedings of the Sixth International Coral Reef Symposium, Townsville, Australia, 8th-12th August 1988*, 2: 613-618.
- WHITLEY, G.P. 1927. Studies in ichthyology. No. 1. *Records of the Australian Museum*, 15 (5): 289-304, pls 24-25.
- WHITLEY, G.P. 1961. Fishes from New Caledonia. *Proceedings of the Royal Zoological Society of New South Wales, (1958-1959) 1961:* 60-65.

### List of Shore fishes of New Caledonia (0-100 m)

#### STEGOSTOMATIDAE

*Stegostoma fasciatum* (Seba in Hermann, 1783) (Synonym: *Stegostoma varium*).

#### GINGLYMOSMATIDAE

*Nebrius ferrugineus* (Lesson, 1831) (Reported by several authors under the name *Nebrius concolor*)

#### RHINCODONTIDAE

*Rhincodon typus* (Smith, 1828)

#### ODONTASPIDIDAE

*Odontaspis noronhai* (Maul, 1955)

#### PSEUDOCARCHARIIDAE

*Pseudocarcharias kamoharai* (Matsubara, 1936)

#### ALOPIIDAE

*Alopias pelagicus* Nakamura, 1935

*Alopias superciliosus* (Lowe, 1841)

*Alopias vulpinus* (Bonnaterre, 1788)

#### LAMNIDAE

*Carcharodon carcharias* (Linnaeus, 1758)

*Isurus oxyrinchus* Rafinesque-Schmaltz, 1810

*Isurus paucus* Guitart Manday, 1966

#### SCYLIORHINIDAE

*Aulohaelurus kanakorum* Séret, 1990

#### CARCHARHINIDAE

*Carcharhinus albimarginatus* (Rüppell, 1837)

*Carcharhinus amblyrhynchos* (Bleeker, 1856)

*Carcharhinus brevipinna* (Müller & Henle, 1839)

*Carcharhinus falciformis* (Bibron in Müller & Henle, 1839)

*Carcharhinus leucas* (Valenciennes in Müller & Henle, 1839)

*Carcharhinus limbatus* (Valenciennes in Müller & Henle, 1839)

*Carcharhinus longimanus* (Poey, 1861)

*Carcharhinus melanopterus* (Quoy & Gaimard, 1824)

*Carcharhinus obscurus* (LeSueur, 1818)

*Carcharhinus plumbeus* (Nardo, 1827)

*Carcharhinus sorrah* (Valenciennes in Müller & Henle, 1839)

*Galeocerdo cuvier* (Peron & LeSueur in LeSueur, 1822)

*Negaprion acutidens* (Rüppell, 1837)

*Prionace glauca* (Linnaeus, 1758)

*Triaenodon obesus* (Rüppell, 1837)

## SPHYRNIDAE

*Sphyraea lewini* (Griffith & Smith, 1834)

*Sphyraea mokarran* (Rüppell, 1837)

## HEXANCHIDAE

*Hexanchus vitulus* Springer & Waller, 1969

## SQUALIDAE

*Squalus blainville* (Risso, 1827)

*Squalus megalops* (Macleay, 1882)

## RHINIDAE

*Rhina aenoclostoma* Bloch & Schneider, 1801

## RHINOBATIDAE

*Rhinobatus djiddensis* (Forsskål in Niebuhr, 1775)

## DASYATIDAE

*Dasyatis bennetti* (Müller & Henle, 1841)

*Dasyatis fluviorum* Ogilby, 1908

*Dasyatis kuhlii* (Müller & Henle, 1841)

*Himantura fai* Jordan & Seale, 1906

*Himantura granulata* (Macleay, 1883)

*Pastinachus sephen* (Forsskål in Niebuhr, 1775)

*Taeniura lymma* (Forsskål in Niebuhr, 1775)

*Taeniura meyeni* Müller & Henle, 1841 (Synonym: *Taeniura melanospila*)

*Urogymnus asperrimus* (Bloch & Schneider, 1801)

## MYLIOBATIDAE

### - SUBFAMILY MYLIOBATINAE

*Aetobatus narinari* (Euphrasen, 1790)

*Aetomylaeus maculatus* (Gray, 1832)

*Aetomylaeus vespertilio* (Bleeker, 1852)

### - SUBFAMILY MOBULINAE

*Manta birostris* (Walbaum, 1792)

*Mobula tarapacana* (Philippi, 1893)

## ELOPIDAE

*Elops hawaiiensis* Regan, 1909 (Reported by several authors under the name *Elops machnata*).

## MEGALOPIDAE

*Megalops cyprinoides* (Broussonet, 1782)

## ALBULIDAE

*Albula forsteri* Valenciennes in Cuvier & Valenciennes, 1846 (Reported by several authors under the name *Albula vulpes*).

*Albula glossodonta* (Forsskål in Niebuhr, 1775)

## ANGUILLIDAE

*Anguilla australis* Richardson, 1841 (Synonym: *Anguilla australis schmidti*)

*Anguilla marmorata* Quoy & Gaimard, 1824 (Synonym: *Anguilla mauritiana*)

*Anguilla megastoma* Kaup, 1856

*Anguilla obscura* Günther, 1871

*Anguilla reinhardtii* Steindachner, 1867

## MORINGUIDAE

*Moringua ferruginea* Bliss, 1883

*Moringua javanica* (Kaup, 1856)

*Moringua macrochir* Bleeker, 1855

*Moringua microchir* Bleeker, 1853

## CHLOPSIDAE

*Kaupichthys atronasus* Schultz in Schultz, Herald, Lachner, Welander & Woods, 1953

*Kaupichthys brachychirus* Schultz, 1953 (New record from New Caledonia based on SMNS material from northeastern Grande Terre)

*Kaupichthys diodontus* Schultz, 1943

*Kaupichthys hyoprionoides* (Strömmann, 1896)

*Xenoconger fryeri* Regan, 1912

**MURAENIDAE**

A record of *Echidna xanthospilos* (Bleeker, 1859), under the name *Gymnomuraena xanthopleura* needs verification.

*Anarchias cantonensis* (Schultz, 1943)

*Anarchias seychellensis* Smith, 1962

*Echidna amblyodon* (Bleeker, 1856) (**New record** from New Caledonia based on SMNS 22791  
from Tao, northeastern Grande Terre)

*Echidna delicatula* (Kaup, 1856) (**New record** from New Caledonia based on SMNS 22943  
from Lifou, Loyalty Islands)

*Echidna leucotaenia* Schultz, 1943 (**New record** from New Caledonia based on SMNS material  
from Grande Terre and Loyalty Islands)

*Echidna nebulosa* (Ahl, 1789)

*Echidna polyzona* (Richardson, 1845) (Synonym: *Megaderus catenatus*)

*Echidna unicolor* Schultz in Schultz, Herald, Lachner, Welander & Woods, 1953

*Enchelycore bayeri* (Schultz in Schultz, Herald, Lachner, Welander & Woods, 1953)

*Enchelycore pardalis* (Temminck & Schlegel, 1846)

*Gymnomuraena zebra* (Shaw & Nodder, 1797)

*Gymnothorax albimarginatus* (Temminck & Schlegel, 1846) (Reported by Rivaton *et al.* [1989]  
under the name *Gymnothorax hepaticus*)

*Gymnothorax berndti* Snyder, 1904

*Gymnothorax boschii* (Bleeker, 1853)

*Gymnothorax buroensis* (Bleeker, 1857)

*Gymnothorax chilosipilus* Bleeker, 1865

*Gymnothorax eurostus* (Abbott, 1861)

*Gymnothorax fimbriatus* (Bennett, 1832) (Reported by Whitley [1961] under the name *Lycodontis melanospilos*)

*Gymnothorax flavimarginatus* (Rüppell, 1830)

*Gymnothorax fuscomaculatus* (Schultz in Schultz, Herald, Lachner, Welander & Woods, 1953)

*Gymnothorax gracilicauda* Jenkins, 1903 (**New record** from New Caledonia based on SMNS  
22912 from Lifou, Loyalty Islands)

*Gymnothorax javanicus* (Bleeker, 1859)

*Gymnothorax kidako* (Temminck & Schlegel, 1846) (**New record** from New Caledonia; reported  
as *Gymnothorax* sp. 5 by Laboute & Grandperrin [2000: 114])

*Gymnothorax margaritophorus* (Bleeker, 1865) (Synonym: *Gymnothorax talofa*)

*Gymnothorax marshallensis* (Schultz, 1953)

*Gymnothorax melatremus* Schultz in Schultz, Herald, Lachner, Welander & Woods, 1953

*Gymnothorax meleagris* (Shaw in Shaw & Nodder, 1795)

*Gymnothorax monochrous* (Bleeker, 1856)

*Gymnothorax nudivomer* (Günther in Playfair & Günther, 1867) (Synonym: *Gymnothorax xanthostomus*)

*Gymnothorax pictus* (Ahl, 1789) (Reported as *Gymnothorax* sp. 4 by Laboute & Grandperrin  
[2000: 113])

*Gymnothorax pindae* Smith, 1962 (Reported as *Gymnothorax* sp. 3 by Laboute & Grandperrin  
[2000: 113])

*Gymnothorax polyuranodon* (Bleeker, 1853)

*Gymnothorax porphyreus* (Guichenot, 1848) (Reported as *Gymnothorax* sp. 2 and 6 by Laboute &  
Grandperrin [2000: 113, 114])

*Gymnothorax pseudothyrsoides* (Bleeker, 1852)

*Gymnothorax richardsonii* (Bleeker, 1852) (Reported by Laboute & Grandperrin [2000: 112] under  
the name *Gymnothorax reticularis*)

*Gymnothorax rueppellii* (McClelland, 1845) (Name erroneously spelled *Gymnothorax ruepelliae*  
by authors)

*Gymnothorax thyrsoideus* (Richardson, 1845)

*Gymnothorax undulatus* (Commerson & Lacepède in Lacepède, 1803)

- Gymnothorax zonipectis* Seale, 1906  
*Rhinomuraena quaesita* Garman, 1888  
*Scuticaria tigrina* (Lesson, 1828)  
*Siderea prosopeion* (Bleeker, 1853)  
*Strophidon sathete* (Hamilton, 1822) (Synonym: *Thyrsoidea macrura*)  
*Uropterygius alboguttatus* Smith, 1962 (**New record** from New Caledonia based on SMNS 23687 from Lifou, Loyalty Islands)  
*Uropterygius concolor* Rüppell, 1838  
*Uropterygius fuscoguttatus* Schultz in Schultz, Herald, Lachner, Welander & Woods, 1953  
*Uropterygius macrocephalus* (Bleeker, 1865)  
*Uropterygius makatei* Gosline, 1958  
*Uropterygius marmoratus* (Lacepède 1803) (**New record** from New Caledonia based on SMNS 18274 from southeastern Grande Terre)  
*Uropterygius xanthopterus* Bleeker, 1859 (Reported by Schmeltz [1869: 27]; record needs verification)  
*Uropterygius xenodontus* McCosker & Smith, 1997

#### OPHICHTHIDAE

- Apterichtus klazingai* (Weber, 1913)  
*Callechelys catostoma* (Forster in Bloch & Schneider, 1801) (Synonym: *Callechelys melanotaenia*)  
*Callechelys marmorata* (Bleeker, 1853) (**New record** from New Caledonia; reported as *Myrichthys* sp. by Laboute & Grandperrin [2000: 118])  
*Ichthyapus vulturis* (Weber & Beaufort, 1916) (**New record** from New Caledonia based on SMNS 21759 from Chesterfield Islands)  
*Lamnostoma kampeni* (Weber & Beaufort, 1916)  
*Lamnostoma orientalis* (McClelland, 1845)  
*Leiuranus semicinctus* (Lay & Bennett, 1839)  
*Muraenichthys gymnotus* Bleeker, 1857  
*Muraenichthys laticaudatus* (Ogilby, 1898)  
*Muraenichthys macropterus* Bleeker, 1857  
*Muraenichthys schultzi* Bleeker, 1857 (**New record** from New Caledonia based on SMNS 22801 from northern Grande Terre)  
*Myrichthys colubrinus* (Boddaert, 1781)  
*Myrichthys maculosus* (Cuvier, 1816)  
*Myrophis uropterus* (Temminck & Schlegel, 1842)  
*Ophichthus bonaparti* (Kaup, 1856)  
*Ophichthus cephalozona* (Bleeker, 1864)  
*Pisodonophis boro* (Hamilton, 1822)  
*Schismorhynchus labialis* (Seale, 1917)  
*Schultzidia johnstonensis* (Schultz & Woods, 1949)  
*Yirrkala insolitus* McCosker, 1999 (Reported by Rivaton *et al.* [1989] under the name *Yirrkala lumbrioides*)

#### MURAENESOCIDAE

- Muraenesox bagio* (Hamilton-Buchanan, 1822)

#### CONRIDAE

- Ariosoma anagoides* (Bleeker, 1854)  
*Ariosoma anago* (Temminck & Schlegel, 1846)  
*Ariosoma mauritianum* (Pappenheim, 1914)  
*Ariosoma scheelei* (Strömmann, 1896)  
*Conger cinereus* (Rüppell, 1828) (Reported by Jouan [1879] as *Conger marginatus*; reported by Fourmanoir, [1981] under the name *Conger japonicus*)  
*Conger verreauxi* Kaup, 1856  
*Conger wilsoni* (Banning in Bloch & Schneider, 1801)  
*Gnathophis habenatus* (Richardson, 1848) (Reported by Castle [1963] as *Gnathophis habenatus longicaudus*)

*Gnathophis umbrellabius* (Whitley, 1948) (Synonym: *Gnathophis incognitus* Castle, 1963; new record from New Caledonia based on Castle [1963: 37-45])

*Heteroconger hassi* (Klausewitz & Eibl-Eibesfeld, 1959) (Synonym: *Taenioconger hassi neocaledoniensis* Castle, 1967)

*Heteroconger tomberua* Castle & Randall, 1999

#### ENGRAULIDAE

*Encrasicholina devisi* (Whitley, 1940)

*Encrasicholina heteroloba* (Rüppell, 1837)

*Encrasicholina punctifer* (Fowler, 1938)

*Engraulis australis* (White, 1790) (Reported by Fowler [1928: 33] under the name *Engraulis indica*, and by Fourmanoir [1971: 110] as *Engraulis japonicus*).

*Stolephorus indicus* (Hasselt, 1823)

*Stolephorus insularis* Hardenberg, 1933 (Reported by Rivaton et al. [1989] under the name *Stolephorus tri*)

*Stolephorus waitei* Jordan & Seale, 1906

*Thryssa baelama* (Forsskål in Niebuhr, 1775)

#### CHIROCENTRIDAE

*Chirocentrus dorab* (Forsskål in Niebuhr, 1775) (Synonym: *Neosudis vorax* Castelnau, 1873)

#### CLUPEIDAE

*Amblygaster clupeoides* Bleeker, 1849

*Amblygaster sirm* (Walbaum, 1792)

*Anodontostoma chacunda* (Hamilton Buchanan, 1822)

*Dussumieria elopsoides* Bleeker, 1849 (Reported by several authors under the name *Dussumieria acuta*)

*Herklotischthys quadrimaculatus* (Rüppell, 1837) (*Clupea mauritiana* Bennett, 1833 is a senior synonym according to Fricke [1999: 73], but should not be used)

*Nematalosa come* (Richardson, 1846)

*Sardinella fijiense* (Fowler & Bean, 1923) (Reported by Borodin [1932] under the name *Clupea sindensis*)

*Sardinella melanura* (Cuvier, 1829)

*Spratelloides delicatulus* (Bennett, 1832)

*Spratelloides gracilis* (Temminck & Schlegel, 1846)

#### CHANIDAE

*Chanos chanos* (Forsskål in Niebuhr, 1775)

#### GONORYNCHIDAE

*Gonorynchus greyi* (Richardson, 1845) (Reported by several authors under the name *Gonorynchus gonorynchus* with different spelling)

#### PLOTOSIDAE

*Plotosus lineatus* (Thunberg, 1787) (Synonyms: *Plotosus anguillaris*, *Plotosus arab*)

#### AULOPIDAE

*Hime japonica* (Günther, 1880) (Reported by several authors under the name *Aulopus japonicus*)

#### SYNODONTIDAE

A record of *Saurida longimanus* (Norman, 1939) by Rivaton et al. (1989: 74) needs verification.

*Saurida gracilis* (Quoy & Gaimard, 1824)

*Saurida nebulosa* Valenciennes in Cuvier & Valenciennes, 1850 (This species was confused by some authors with *S. gracilis*)

*Saurida tumbil* (Bloch, 1795)

*Saurida undosquamis* (Richardson, 1848) (Reported by Laboute & Grandperrin, 2000: 126 under the name *Saurida* sp.)

*Synodus binotatus* Schultz in Schultz, Herald, Lachner, Welander & Woods, 1953

*Synodus dermatogenys* Fowler, 1912 (This species was confused by several authors with

*Synodus variegatus* and *Synodus englemani*)

*Synodus doaki* Russell & Cressey, 1979

*Synodus hoshionis* (Tanaka, 1917)

*Synodus jaculum* Russell & Cressey, 1979

- Synodus macrocephalus* (Cressey, 1981)  
*Synodus oculatus* Cressey, 1981  
*Synodus rubromarmoratus* Russell & Cressey, 1979  
*Synodus similis* (McCulloch, 1921)  
*Synodus tectus* (Cressey, 1981) (Reported by Rivaton [1989: 142] under the name *Synodus variegatus*)  
*Synodus variegatus* (Commerson in Lacepède, 1803) (Reported by Thollot & Kulbicki [1988: 618] under the name *Synodus englemani*)  
*Trachinocephalus myops* (Forster in Bloch & Schneider, 1801)

#### MYCTOPHIDAE

Some species of Myctophidae, like *Diaphus metopoclampus* (Cocco, 1829) or *Myctophum nitidulum* Garman, 1899, may be encountered inshore at night (e.g. at Baie du Santal, Lifou, Loyalty Islands; identification based on SMNS material).

#### VELIFERIDAE

- Metavelifer multiradiatus* (Regan, 1907)

#### LAMPRIDAE

- Lampris guttatus* (Brünnich, 1788) (Reported by Fourmanoir & Laboute [1976: 326] under the name *Lampris regius*)

#### TRACHIPTERIDAE

- Zu cristatus* (Bonelli, 1819)

#### POLYMXIIDAE

- Polymixia berndti* Gilbert, 1905

#### BREGMACEROTIDAE

- Bregmaceros* sp. (Undescribed species; reported by authors under the name *Bregmaceros maclellandi* or *Bregmaceros mcclellandii*)

- Bregmaceros japonicus* Tanaka, 1908 (New record from New Caledonia based on SMNS 23663 from Lifou, Loyalty Islands)

- Bregmaceros nectabanus* Whitley, 1941

- Bregmaceros rarissimus* Munro, 1950

#### MORIDAE

- Physiculus therosideros* Paulin, 1987

#### CARAPIDAE

- Carapus mourlani* (Petit, 1934)

- Echiodon coheni* Williams, 1984

- Encheliophis homei* (Richardson, 1846)

- Encheliophis vermicularis* (Müller, 1842)

- Onuxodon fowleri* (Smith, 1955) (Reported by authors under the name *Onuxodon margaritiferae*)

- Onuxodon parvibrachium* (Fowler, 1927)

- Pyramodon ventralis* Smith & Radcliffe, 1913

#### OPHIDIIDAE

- Brotula multibarbata* Temminck & Schlegel, 1846

- Brotula townsendi* Fowler, 1901

- Monomitus garmani* (Smith & Radcliffe in Radcliffe, 1913)

- Ophidion muraenolepis* (Günther, 1880)

#### BYTHITIDAE

Several undescribed species are known to occur at New Caledonia; descriptions are under preparation by Schwarzhans, Møller and Nielsen.

- Diancistrus brevirostris* Schwarzhans, Møller & Nielsen, 2005 (Reported by Kulbicki & Williams [1997: 12] under the name *Brosmophyciops pautzkei* [in part])

- Diancistrus longifilis* Ogilby, 1899

- Diancistrus tongensis* Schwarzhans, Møller & Nielsen, 2005 (Reported by authors under the name *Brosmophyciops pautzkei* [in part])

- Dinematichthys randalli* Machida, 1994

- Dinematichthys riukiensis* Aoyagi, 1954

## ANTENNARIIDAE

- Antennarius coccineus* (Cuvier in Lesson, 1831) (Synonym: *Abantennarius neocaledoniensis* Danois, 1964)  
*Antennarius commerson* (Commerson & Lacepède in Anonymus, 1798) (Synonym: *Antennarius moluccensis*; reported by Laboute & Grandperrin [2000: 131] under the name *Antennarius* sp. 1-3)  
*Antennarius duescus* Snyder, 1904  
*Antennarius hispidus* (Bloch & Schneider, 1801) (**New record** from New Caledonia based on SMNS 19848 from Maré, Loyalty Islands)  
*Antennarius maculatus* (Desjardins, 1840)  
*Antennarius nummifer* Cuvier, 1816  
*Antennarius pictus* (Shaw in Shaw & Nodder, 1794) (Reported by Whitley [1961: 64] under the name *Antennarius phymatodes*)  
*Antennarius rosaceus* Smith & Radcliffe in Radcliffe, 1912 (**New record** from New Caledonia based on SMNS 23913 from Lifou, Loyalty Islands)  
*Antennarius striatus* (Shaw in Shaw & Nodder, 1794) (Synonyms: *Chironectes tridens* Temminck & Schlegel, 1845; *Phrynelox zebrinus* Schultz, 1957)  
*Antennatus tuberosus* (Cuvier, 1817) (Reported by Laboute & Grandperrin [2000: 132] under the name *Antennarius* sp. 4; **new record** from New Caledonia, also based on SMNS 23904 from Lifou, Loyalty Islands)  
*Histrio histrio* (Linnaeus, 1758)

## MUGILIDAE

- Mugil broussoneti* Valenciennes in Cuvier & Valenciennes, 1836 was doubtfully recorded from New Caledonia by Harrision & Senou, (1999: 2095).  
*Cestraeus goldiei* (Macleay, 1883)  
*Cestraeus oxyrhyncus* Valenciennes in Cuvier & Valenciennes, 1836  
*Cestraeus plicatilis* Valenciennes in Cuvier & Valenciennes, 1836  
*Chelon macrolepis* (Smith, 1849)  
*Chelon melinopterus* (Valenciennes in Cuvier & Valenciennes, 1836)  
*Chelon planiceps* (Valenciennes in Cuvier & Valenciennes, 1836) (Synonym: *Liza tade*)  
*Chelon subviridis* (Valenciennes in Cuvier & Valenciennes, 1836)  
*Crenimugil crenilabis* (Forsskål in Niebuhr, 1775) (Synonym: *Mugil neocaledonicus* Castelnau, 1873)  
*Crenimugil heterocheilos* (Bleeker, 1855)  
*Ellochelon vaigiensis* (Quoy & Gaimard, 1824) (Synonym: *Liza macrolepidotus*)  
*Moolgarda buchanani* (Bleeker, 1853)  
*Moolgarda cunnesius* (Valenciennes in Cuvier & Valenciennes, 1836)  
*Moolgarda engeli* (Bleeker, 1858)  
*Moolgarda seheli* (Forsskål in Niebuhr, 1775)  
*Mugil cephalus* Linnaeus, 1758 (Synonym: *Mugil catalaram* Whitley, 1951)  
*Oedalechilus labiosus* (Valenciennes in Cuvier & Valenciennes, 1836) (**New record** from New Caledonia based on SMNS 23994 from southern Grande Terre)

## ATHERINIDAE

Record of *Atherinomorus insularum* (Jordan & Evermann, 1903) by Borodin (1932: 76) under the name *Atherina insularum* needs verification.

- Atherinomorus duodecimalis* (Valenciennes in Cuvier & Valenciennes, 1835) (Reported by authors under the name *Atherinomorus endrachtensis*).  
*Atherinomorus lacunosus* (Forster in Bloch & Schneider, 1801) (Reported by Whitley [1961: 65] under the name *Pranesus ogilbyi*, and by Fourmanoir & Laboute [1976: 306] under the name *Pranesus pinguis*).  
*Atherion elymus* Jordan & Starks, 1901  
*Hypoatherina barnesi* Schultz in Schultz, Herald, Lachner, Welander & Woods, 1953  
*Hypoatherina ovalaua* (Herre, 1935)  
*Hypoatherina temminckii* (Bleeker, 1853)  
*Stenatherina panatela* (Jordan & Richardson, 1908)

## EXOCETIDAE

Records of *Cypselurus bruuni* Kotthaus, 1965 by Fourmanoir (1971: 111), and of *Hirundichthys volador* (Jordan, 1884) by Rivaton *et al.* (1989: 38) under the name *Prognichthys rondeletii*, need verification.

*Cheilopogon atrisignis* (Jenkins, 1904)  
*Cheilopogon dorsomacula* (Fowler, 1944)  
*Cheilopogon furcatus* (Mitchill, 1815)  
*Cheilopogon rapanouiensis* Parin, 1962  
*Cheilopogon spilonotopterus* (Bleeker, 1866)  
*Cheilopogon spilopterus* (Valenciennes in Cuvier & Valenciennes, 1846)  
*Cheilopogon unicolor* (Valenciennes in Cuvier & Valenciennes, 1847) (Synonym: *Cypsilurus ogilbyi* Jordan & Snyder in Jordan & Dickerson, 1908)  
*Cypselurus angusticeps* Nichols & Breder, 1935  
*Cypselurus naresii* (Günther, 1889)  
*Cypselurus oligolepis* (Bleeker, 1866)  
*Cypselurus poecilopterus* (Valenciennes in Cuvier & Valenciennes, 1846)  
*Exocoetus obtusirostris* Günther, 1866  
*Exocoetus volitans* Linnaeus, 1758  
*Hirundichthys speculiger* (Valenciennes in Cuvier & Valenciennes, 1847)  
*Oxyporhamphus meristocystis* (Parin, 1961)  
*Parexocoetus brachypterus* (Günther, 1866)

#### HEMIRAMPHIDAE

*Euleptorhamphus viridis* (Russell in Hasselt, 1823)  
*Hemiramphus far* (Forsskål in Niebuhr, 1775) (Reported by Jouan [1879] under the name  
    *Hemiramphus marginatus*, and by Castelnau, [1873] under the name *Hemiramphus commersonii*)  
*Hyporhamphus dussumieri* (Valenciennes in Cuvier & Valenciennes, 1846)  
*Zenarchopterus dispar* (Valenciennes in Cuvier & Valenciennes, 1846)

#### BELONIDAE

*Ablennes hians* (Valenciennes in Cuvier & Valenciennes, 1846)  
*Platybelone argalus platyura* (Bennett, 1832)  
*Strongylura incisa* (Valenciennes in Cuvier & Valenciennes, 1846)  
*Strongylura leiura* (Bleeker, 1851)  
*Strongylura urvillii* (Valenciennes in Cuvier & Valenciennes, 1846)  
*Tylosurus acus melanotus* (Bleeker, 1851)  
*Tylosurus crocodilus crocodilus* (Peron & LeSueur in LeSueur, 1821) (Reported by Seale [1935:  
    345] under the name *Tylosurus indica*)

#### ANOMALOPIDAE

*Anomalops katopron* Bleeker, 1856  
*Photoblepharon palpebratus* (Boddaert, 1781)

#### MONOCENTRIDAE

*Monocentris japonica* (Houttuyn, 1782)

#### TRACHICHYIDAE

*Gephyroberyx darwini* (Johnson, 1866)  
*Paratrachichthys trailli* (Hutton, 1875)  
*Parinoberyx horridus* Kotlyar, 1984

#### BERYCIDAЕ

*Beryx splendens* Lowe, 1834  
*Centroberyx affinis* (Günther, 1859)  
*Centroberyx druzhinini* (Busakhin, 1981)

#### HOLOCENTRIDAE

Records of *Myripristis amaena* by Rivaton *et al.* (1989: 43) and Kulwicki & Williams (1997: 12) are  
probably based on *Myripristis pralinia*.  
*Myripristis adusta* Bleeker, 1853  
*Myripristis berndti* Jordan & Evermann, 1903 (Reported by Fourmanoir & Laboute [1976: 155]  
under the name *Myripristis amaenus*)  
*Myripristis botche* Cuvier, 1829 (Synonym: *Myripristis melanostictus*; reported by Fourmanoir &  
Laboute [1976: 156-157] under the name *Myripristis murdjan*)  
*Myripristis hexagona* (Lacepède, 1802)  
*Myripristis kuntee* Valenciennes in Cuvier & Valenciennes, 1831

- Myripristis murdjan* (Forsskål in Niebuhr, 1775) (Synonym: *Myripristis bowditchae* Woods in Schultz et al., 1953)
- Myripristis pralinia* Cuvier, 1829
- Myripristis violacea* Bleeker, 1851
- Myripristis vittata* Valenciennes in Cuvier & Valenciennes, 1831 (Reported by Borodin [1932: 75] under the name *Myripristis trachyacron*, and by Laboute & Grandperrin [2000: 45, 140] under the name *Myripristis* sp.)
- Myripristis woodsi* Greenfield, 1974 (Reported by Laboute & Grandperrin [2000: 143] under the name *Myripristis* sp.; new record from New Caledonia)
- Neoniphon argenteus* (Valenciennes in Cuvier & Valenciennes, 1831)
- Neoniphon opercularis* (Valenciennes in Cuvier & Valenciennes, 1831)
- Neoniphon sammara* (Forsskål in Niebuhr, 1775)
- Ostichthys hysipterygion* Randall, Shimizu & Yamakawa, 1982
- Ostichthys japonicus* (Cuvier in Cuvier & Valenciennes, 1829)
- Ostichthys kaianus* (Günther, 1880)
- Plectrypops lima* (Valenciennes in Cuvier & Valenciennes, 1831) (Reported by Laboute & Grandperrin [2000: 141] under the name *Myripristis hexagona*)
- Sargocentron caudimaculatum* (Rüppell, 1838)
- Sargocentron cornutum* (Bleeker, 1853)
- Sargocentron diadema* (Commerçon & Lacepède in Lacepède, 1802)
- Sargocentron ensifer* (Jordan & Evermann, 1903) (Reported by Fourmanoir & Laboute [1976: 153] and Laboute & Grandperrin [2000: 138, 145] under the name *Adioryx furcatus* or *Sargocentron furcatum*)
- Sargocentron iota* Randall, 1998
- Sargocentron lepros* (Allen & Cross, 1983)
- Sargocentron melanospilos* (Bleeker, 1858)
- Sargocentron microstoma* (Günther, 1859)
- Sargocentron praslin* (Lacepède, 1802) (New record from New Caledonia based on SMNS 22634 from southern Grande Terre)
- Sargocentron punctatissimum* (Cuvier in Cuvier & Valenciennes, 1829)
- Sargocentron rubrum* (Forsskål in Niebuhr, 1775)
- Sargocentron spiniferum* (Forsskål in Niebuhr, 1775)
- Sargocentron tiere* (Cuvier in Cuvier & Valenciennes, 1829)
- Sargocentron violaceum* (Bleeker, 1853)

## ZEIDAE

- Zenopsis nebulosa* (Temminck & Schlegel, 1845)

## PEGASIDAE

- Eurypegasus draconis* (Linné, 1766)

## SOLENOSTOMIDAE

- Solenostomus cyanopterus* Bleeker, 1854

- Solenostomus paradoxus* (Pallas, 1770)

## SYNGNATHIDAE

- Acentronura breviperula* Fraser-Brunner & Whitley, 1949 (Reported by Rivaton et al. [1989: 73] under the names *Acentronura tentaculata*, *Acentronura australis* and *Acentronura gracilissima*, and by Laboute & Grandperrin [2000: 157, 159] under the names *Halicampus* sp. and *Hippocampus* sp.)

- Corythoichthys* sp. (Undescribed species similar to *Corythoichthys amplexus*)

- Corythoichthys conspicillatus* (Jenyns, 1842) (Reported by Fourmanoir [1971: 112] under the name *Corythoichthys flavofasciatus*).

- Corythoichthys haematopterus* (Bleeker, 1851)

- Corythoichthys intestinalis* (Ramsay, 1881) (Reported by Whitley [1961: 64] under the name

- Corythoichthys flavofasciatus*, by Plessis & Fourmanoir [1966: 125] under the name *Hippichthys intestinalis waitei*, and by Kulbicki et al. [1994: 15] under the name *Corythoichthys amplexus*)

- Corythoichthys nigripectus* Herald in Schultz, Herald, Lachner, Welander & Woods, 1953

- Corythoichthys ocellatus* Herald in Schultz, Herald, Lachner, Welander & Woods, 1953

- Corythoichthys paxtoni* Dawson, 1977

*Corythoichthys schultzi* Herald in Schultz, Herald, Lachner, Welander & Woods, 1953  
*Cosmocampus banneri* (Herald & Randall, 1972)  
*Doryrhamphus melanopleura* (Bleeker, 1858) (Reported by several authors under the name  
    *Doryrhamphus excisus excisus*)  
*Dunckerocampus chapmani* (Herald in Schultz, Herald, Lachner, Welander & Woods, 1953)  
*Dunckerocampus dactyliophorus* (Bleeker, 1853)  
*Festucalex kulwickii* Fricke, 2004 (Reported by Plessis & Fourmanoir [1966: 125] and other  
authors under the name *Ichthyocampus erythraeus* or *Festucalex erythraeus*, by others as *Festucalex*  
*gibbsi*, and by Kuiter [2000: 122] as *Festucalex* sp. 1)  
*Festucalex wassi* Dawson, 1977  
*Halicampus boothae* (Whitley, 1964)  
*Halicampus brocki* (Herald in Schultz, Herald, Lachner, Welander & Woods, 1953)  
*Halicampus dunckeri* (Chabanaud, 1929)  
*Halicampus mataafae* (Jordan & Seale, 1906)  
*Halicampus nitidus* (Günther, 1873)  
*Halicampus spinirostris* (Dawson & Allen, 1981)  
*Hippichthys spicifer* (Rüppell, 1838)  
*Hippocampus bargibanti* Whitley, 1970  
*Hippocampus curvicuspis* Fricke, 2004 (Reported by several authors under the name *Hippocampus*  
*histrix*, by Fourmanoir [1971: 112] as *Hippocampus jayakari*, and by Myers [1999: 89] and Laboute &  
Grandperrin [2000: 158, large fig.] as *Hippocampus kuda*)  
*Hippocampus pusillus* Fricke, 2004  
*Hippocampus semispinosus* Kuiter, 2001 (Reported by several authors under the name  
    *Hippocampus kuda*, and by Lourie, Vincent & Hall [1999: 92-93] as *Hippocampus fisheri*)  
*Hippocampus taeniopterus* Bleeker, 1852  
*Micrognathus brevicorpus* Fricke, 2004 (Reported by several authors under the name *Micrognathus*  
*andersonii*)  
*Micrognathus micronotopterus* (Fowler, 1938)  
*Micrognathus natans* Dawson, 1982  
*Micrognathus pygmaeus* Fritzsche, 1981 (Reported by several authors under the name  
    *Micrognathus brevirostris*)  
*Microphis argulus* (Peters, 1855)  
*Microphis brachyurus* (Bleeker, 1853)  
*Microphis brevidorsalis* (Beaufort, 1913)  
*Microphis leiaspis* (Bleeker, 1853)  
*Microphis retzii* (Bleeker, 1856)  
*Phoxocampus belcheri* (Kaup, 1856)  
*Phoxocampus diacanthus* (Schultz, 1943)  
*Siokunichthys herrei* Herald in Schultz, Herald, Lachner, Welander & Woods, 1953  
*Siokunichthys striatus* Fricke, 2004  
*Syngnathoides biaculeatus* (Bloch, 1785)  
*Trachyrhamphus bicoarctatus* (Bleeker, 1857) (Reported by Jouan [1879: 332] under the name  
*Ichthyocampus maculatus*, by Fourmanoir [1971: 112] and Fourmanoir & Laboute [1976: 272] as *Yozia*  
*intermedia*, and by Rivaton *et al.* [1989: 74] and Rivaton & Bourret [1999: 46] as *Trachyrhamphus longirostris*)

#### AULOSTOMIDAE

*Aulostomus chinensis* (Linné, 1766)

#### FISTULARIIDAE

*Fistularia commersonii* Rüppell, 1838 (Confused with *Fistularia petimba* by Fowler [1928: 117-118])

*Fistularia petimba* Lacepède, 1803 (Records from New Caledonia need verification; they may be  
    based on *F. commersonii* according to Fritzsche & Thiesfeld [1999: 2279]; synonym: *Fistularia villosa*)

#### MACRORAMPHOSIDAE

*Macroramphosus scolopax* (Linnaeus, 1758)

## CENTRISCIDAE

*Aeoliscus strigatus* (Günther, 1860) (Reported by Jouan [1863: 183] and Fowler [1928: 118] under the name *Amphisile scutatum* or *Centriscus scutatus*)

## DACTYLOPTERIDAE

*Dactyloptena orientalis* (Cuvier in Cuvier & Valenciennes, 1829)

## SCORPAENIDAE

### - SUBFAMILY SCORPAENINAE

*Dendrochirus bellus* (Jordan & Hubbs, 1925)

*Dendrochirus biocellatus* (Fowler, 1938)

*Dendrochirus brachypterus* (Cuvier in Cuvier & Valenciennes, 1829)

*Dendrochirus zebra* (Cuvier in Cuvier & Valenciennes, 1829) (Reported by Whitley [1961: 65] under the name *Pterois volitans*)

*Iracundus signifer* Jordan & Evermann, 1903

*Parascorpaena mcadamsi* (Fowler, 1938)

*Parascorpaena mossambica* (Peters, 1855)

*Parascorpaena picta* Kuhl & Hasselt in Cuvier, 1829 (Synonym: *Sebastapistes bynoensis*)

*Phenacoscorpius megalops* Fowler, 1938

*Pterois antennata* (Bloch, 1787)

*Pterois lunulata* Temminck & Schlegel, 1844

*Pterois mombasae* (Smith, 1957)

*Pterois radiata* Cuvier, 1829

*Pterois volitans* (Linnaeus, 1758)

*Rhinopias aphanes* Eschmeyer, 1973

*Scorpaena neglecta* Temminck & Schlegel, 1844

*Scorpaenodes albaiensis* (Evermann & Seale, 1907)

*Scorpaenodes corallinus* Smith, 1957

*Scorpaenodes guamensis* (Quoy & Gaimard, 1824) (Reported by Laboute & Grandperrin [2000: 167] under the name *Scorpaenodes* sp. 2)

*Scorpaenodes hirsutus* (Smith, 1957)

*Scorpaenodes kelloggi* (Jenkins, 1903) (Reported by Laboute & Grandperrin [2000: 167] under the name *Scorpaenodes* sp. 1)

*Scorpaenodes minor* (Smith, 1958) [*Scorpaenodes brocki* (Schultz, in Schultz, Woods & Lachner, 1966) is a synonym]

*Scorpaenodes parvipinnis* (Garrett, 1864)

*Scorpaenodes scaber* (Ramsay & Ogilby, 1885)

*Scorpaenodes varipinnis* Smith, 1957

*Scorpaenopsis diabolus* Cuvier, 1829

*Scorpaenopsis eschmeyeri* Greenfield & Randall, 2004

*Scorpaenopsis macrochir* Ogilby, 1910 (Reported by several authors under the name *Scorpaenopsis gibbosa*, and by Laboute & Grandperrin [2000: 169] under the name *Scorpaenopsis* sp.)

*Scorpaenopsis neglecta* Heckel, 1837

*Scorpaenopsis papuensis* (Cuvier in Cuvier & Valenciennes, 1829) (Reported by Fourmanoir & Laboute [1976: 158] under the name *Scorpaenopsis cirrhosa*, and by other authors as *Scorpaenopsis oxycephala*)

*Scorpaenopsis possi* Randall & Eschmeyer, 2001

*Scorpaenopsis ramaraoi* Randall & Eschmeyer, 2001

*Scorpaenopsis venosa* (Cuvier in Cuvier & Valenciennes, 1829)

*Scorpaenopsis vittapinna* Randall & Eschmeyer, 2001 (Reported by Kulbicki *et al.* [1994: 17] under the name *Scorpaenopsis brevifrons*)

*Sebastapistes cyanostigma* (Bleeker, 1856) (Synonym: *Scorpaena albobrunnea* Günther, 1874)

*Sebastapistes coniorta* Jenkins, 1903 (New record from New Caledonia based on SMNS 23749 from Lifou, Loyalty Islands)

*Sebastapistes fowleri* (Pietschmann, 1934)

*Sebastapistes galactacma* Jenkins, 1903 (New record from New Caledonia based on SMNS material from Lifou, Loyalty Islands)

*Sebastapistes mauritiana* (Cuvier, 1829)

*Sebastapistes strongia* (Cuvier, 1829) (Synonym: *Kantapus oglinus* Smith, 1947)

*Sebastapistes tinkhami* (Fowler, 1946)

*Taenianotus triacanthus* Lacepède, 1802

- SUBFAMILY TETRAROGINAE

*Ablabys taenianotus* (Cuvier in Cuvier & Valenciennes, 1829)

*Ocosia apia* Poss & Eschmeyer, 1975

*Paracentropogon longispinis* (Cuvier, 1829)

*Richardsonichthys leucogaster* (Richardson, 1848)

*Tetraoge barbata* (Cuvier, 1829)

- SUBFAMILY SYNANCEIINAE

*Erosa erosa* (Langsdorf in Cuvier & Valenciennes, 1829)

*Inimicus caledonicus* (Sauvage, 1878)

*Inimicus didactylus* (Pallas, 1769)

*Minous monodactylus* (Bloch & Schneider, 1801)

*Minous pusillus* Temminck & Schlegel, 1843

*Minous trachycephalus* (Bleeker, 1854)

*Synanceia horrida* (Linné, 1766)

*Synanceia verrucosa* (Bloch & Schneider, 1801)

## CARACANTHIDAE

*Caracanthus maculatus* (Gray, 1831)

*Caracanthus unipinna* (Gray, 1831)

## APLOACTINIDAE

*Aploactis aspera* (Richardson, 1844)

*Cocotropus dermacanthus* (Bleeker, 1852)

*Cocotropus richeri* Fricke, 2004

*Erisphex pottii* (Steindachner, 1896)

*Neoaploactis tridorsalis* Eschmeyer & Allen, 1978

*Paraploactis* sp. (Very similar to *Paraploactis kagoshimensis* from Japan and *P. obbesi* from Indonesia; specimens under study by Stuart G. Poss according to Randall [2005: 130])

*Paraploactis trachyderma* Bleeker, 1865

## TRIGLIDAE

*Pterygotrigla andertoni* Waite, 1910 (Reported by Cerro & Lloris [1997: 15-117] under the name

*Pterygotrigla picta*)

*Satyrichthys rieffeli* (Kaup, 1859)

## PLATYCEPHALIDAE

Records of *Papilloculiceps longiceps* (Ehrenberg in Cuvier & Valenciennes, 1829) by Rivaton *et al.* [1989: 61] and Rivaton & Bourret [1999: 50] are erroneous, as the species is restricted to the Red Sea and Western Indian Ocean.

*Cociella crocodila* (Tilesius, 1812)

*Cymbacephalus beauforti* (Knapp, 1973) (Reported by Whitley [1961: 64] under the name *Cociella crocodilus*)

*Cymbacephalus staigeri* (Castelnau, 1875)

*Eurycephalus arenicola* (Schultz, in Schultz, Woods & Lachner, 1966)

*Eurycephalus otaitensis* (Parkinson in Cuvier & Valenciennes, 1829)

*Onigocia bimaculata* Knapp, Imamura & Sakashita, 2000

*Onigocia macrolepis* (Bleeker, 1854)

*Onigocia pedimacula* (Regan, 1908) (Reported by several authors under the name *Onigocia spinosa*; records from New Caledonia need verification, may be based on *O. bimaculata*)

*Rogadius patriciae* Knapp, 1987

*Rogadius pristiger* (Cuvier in Cuvier & Valenciennes, 1829) (Reported by Rivaton *et al.* [1989: 61] under the name *Rogadius asper*)

*Rogadius serratus* (Cuvier in Cuvier & Valenciennes, 1829)

*Rogadius welanderi* (Schultz, in Schultz, Woods & Lachner, 1966) (Reported by Kulbicki *et al.* [1994: 18] under the name *Rogadius* sp.; new record from New Caledonia based on SMNS 21758 from

Chesterfield Islands)

*Thysanophrys celebicus* (Bleeker, 1854)

*Thysanophrys chiltonae* Schultz, in Schultz, Woods & Lachner, 1966

## AMBASSIDAE

*Ambassis buruensis* Bleeker, 1856

*Ambassis interruptus* Bleeker, 1852

*Ambassis miops* Günther, 1871

## SYMPHYSANODONTIDAE

*Symphsanodon maunaloaee* Anderson, 1970

## SERRANIDAE

### - SUBFAMILY ANTHIINAE

*Caprodon schlegelii* (Günther, 1859)

*Luzonichthys waitei* (Fowler, 1931) (Synonym: *Luzonichthys robustus* Fourmanoir, 1977)

*Luzonichthys whitleyi* (Smith, 1955)

*Luzonichthys williamsi* Randall & McCosker, 1992

*Plectranthias* sp. 1 (Undescribed species, based on SMNS material from Lifou, Loyalty Islands)

*Plectranthias* sp. 2 (Undescribed species, based on SMNS material from Lifou, Loyalty Islands)

*Plectranthias fourmanoiri* Randall, 1980

*Plectranthias kamii* Randall, 1980 (Reported by Rivaton *et al.* [1989: 70] under the name

*Plectranthias anthiooides*)

*Plectranthias kelloggi* (Jordan & Evermann, 1903)

*Plectranthias longimanus* (Weber, 1913)

*Plectranthias nanus* Randall, 1980

*Plectranthias randalli* Fourmanoir & Rivaton, 1980

*Plectranthias retrofasciatus* Fourmanoir & Randall, 1979

*Plectranthias rubrifasciatus* Fourmanoir & Randall, 1979

*Plectranthias wheeleri* Randall, 1980 (New record from New Caledonia based on SMNS 24052

from Lifou, Loyalty Islands)

*Plectranthias winniensis* (Tyler, 1966)

*Pseudanthias bicolor* (Randall, 1979)

*Pseudanthias carlsoni* Randall & Pyle, 2001

*Pseudanthias cichlops* (Bleeker, 1853)

*Pseudanthias cooperi* (Regan, 1902) (Reported by Laboute & Grandperrin [2000: 203] under the name 'Anthiinae indéterminée 3')

*Pseudanthias elongatus* (Franz, 1910)

*Pseudanthias flavicauda* Randall & Pyle, 2001 (Reported by Laboute & Grandperrin [2000: 203]

under the name Anthiinae indéterminée 2; new record for New Caledonia)

*Pseudanthias hypselosoma* (Bleeker, 1878)

*Pseudanthias lori* (Lubbock & Randall, 1976)

*Pseudanthias pascalus* (Jordan & Tanaka, 1927)

*Pseudanthias pictilis* (Randall & Allen, 1978)

*Pseudanthias pleurotaenia* (Bleeker, 1857)

*Pseudanthias rubrizonatus* (Randall, 1983)

*Pseudanthias squamipinnis* (Peters, 1855)

*Pseudanthias ventralis ventralis* (Randall, 1979)

*Sacura margaritacea* (Hilgendorf, 1879)

*Serranocirrhitus latus* Watanabe, 1949 (Synonym: *Dactylanthias mcmichaeli* Whitley, 1962)

### - SUBFAMILY EPINEPHELINAE

#### - TRIBE EPINEPHELINI

A record of *Epinephelus corallicola* (Valenciennes in Cuvier & Valenciennes, 1828) by Plessis & Fourmanoir (1966: 127), and the record of *Epinephelus melanostigma* Schultz in Schultz, Herald, Lachner, Welander & Woods, 1953 and *Plectropomus areolatus* (Rüppell, 1830) by Rivaton *et al.* (1989: 69, 70) need verification.

*Aethaloperca rogaa* (Forsskål in Niebuhr, 1775)

*Anyperodon leucogrammicus* (Valenciennes in Cuvier & Valenciennes, 1828)

*Cephalopholis argus* Bloch & Schneider, 1801  
*Cephalopholis aurantia* (Valenciennes in Cuvier & Valenciennes, 1828)  
*Cephalopholis boenak* (Bloch, 1790) (Reported by Fourmanoir & Laboute [1976: 52, 53] and other authors under the names *Cephalopholis formosanus* (non Shaw & Nodder, 1812) and *Cephalopholis pachycentron*)  
*Cephalopholis igarashiensis* Katayama, 1957  
*Cephalopholis leopardus* (Lacepède, 1801)  
*Cephalopholis microprion* (Bleeker, 1852)  
*Cephalopholis miniata* (Forsskål in Niebuhr, 1775)  
*Cephalopholis sonnerati* (Valenciennes in Cuvier & Valenciennes, 1828)  
*Cephalopholis spiloparaea* (Valenciennes in Cuvier & Valenciennes, 1828)  
*Cephalopholis urodeta* (Forster in Bloch & Schneider, 1801)  
*Cromileptes altivelis* (Valenciennes in Cuvier & Valenciennes, 1828)  
*Epinephelus areolatus* (Forsskål in Niebuhr, 1775) (Synonym: *Serranus angularis* Valenciennes in Cuvier & Valenciennes, 1828)  
*Epinephelus coeruleopunctatus* (Bloch, 1790) (Synonym: *Serranus hoevenii* Bleeker, 1849)  
*Epinephelus chlorostigma* (Valenciennes in Cuvier & Valenciennes, 1828)  
*Epinephelus coioides* (Hamilton, 1822) (Synonym: *Serranus suillus* Valenciennes in Cuvier & Valenciennes, 1828)  
*Epinephelus cyanopodus* (Richardson, 1846) (Synonyms: *Serranus hoedtii* Bleeker, 1855; *Epinephelus kohleri* Schultz in Schultz et al., 1953)  
*Epinephelus fasciatus* (Forsskål in Niebuhr, 1775)  
*Epinephelus fuscoguttatus* (Forsskål in Niebuhr, 1775) (Synonym: *Serranus horridus* Valenciennes in Cuvier & Valenciennes, 1828)  
*Epinephelus hexagonatus* (Bloch & Schneider, 1801)  
*Epinephelus howlandi* (Günther, 1873) (Reported by Whitley [1961: 64] under the name *Epinephelus corallicola*; synonym: *Epinephelus spilotus* Schultz in Schultz et al., 1953)  
*Epinephelus lanceolatus* (Bloch, 1790)  
*Epinephelus macropsilos* (Bleeker, 1855)  
*Epinephelus maculatus* (Bloch, 1790)  
*Epinephelus magniscutis* Postel, Fourmanoir & Guézé, 1963  
*Epinephelus malabaricus* (Bloch & Schneider, 1801) (Reported by Borodin [1932: 79] under the name *Serranus stoliczkae* [non Day, 1875]; synonym: *Epinephelus cylindricus* Postel, 1965)  
*Epinephelus merra* Bloch, 1793  
*Epinephelus morrhua* (Valenciennes in Cuvier & Valenciennes, 1833)  
*Epinephelus ongus* (Bloch, 1790) (Reported by several authors under the name *Epinephelus summana* [non Forsskål in Niebuhr, 1775])  
*Epinephelus polyphekadion* (Bleeker, 1849) (Synonym: *Epinephelus microdon*)  
*Epinephelus retouti* (Bleeker, 1868)  
*Epinephelus rivulatus* (Valenciennes in Cuvier & Valenciennes, 1830) (Synonym: *Serranus rhyncholepis* Bleeker, 1852)  
*Epinephelus spilotoceps* Schultz in Schultz, Herald, Lachner, Welander & Woods, 1953 (Reported by Laboute & Grandperrin [2000: 189] under the name *Epinephelus* sp.; new record from New Caledonia)  
*Epinephelus tauvina* (Forsskål in Niebuhr, 1775)  
*Gracila albomarginata* (Fowler & Bean, 1930)  
*Plectropomus laevis* (Lacepède, 1802) (Synonym: *Plectropomus melanoleucus*)  
*Plectropomus leopardus* (Lacepède, 1802) (Reported by Whitley [1961: 65] under the name *Plectropomus maculatus*)  
*Variola albimarginata* Baissac, 1952  
*Variola louti* (Forsskål in Niebuhr, 1775)  
- TRIBE DIPLOPRIONI  
*Belonoperca chabanaudi* Fowler & Bean, 1930  
*Diploprion bifasciatum* Kuhl & Hasselt in Cuvier & Valenciennes, 1828  
- TRIBE LIOPROPOMATINI  
*Liopropoma susumi* (Jordan & Seale, 1906)

*Liopropoma tonstrinum* Randall & Taylor, 1988

- TRIBE GRAMMISTINI

*Aporops bilinearis* Schultz, 1943

*Grammistes sexlineatus* (Thunberg, 1792)

*Grammistops ocellatus* Schultz in Schultz, Herald, Lachner, Welander & Woods, 1953

*Pogonoperca punctata* (Valenciennes in Cuvier & Valenciennes, 1830)

*Pseudogramma astignum* Randall & Baldwin, 1997 (New record from New Caledonia based on SMNS material from Lifou, Loyalty Islands)

*Pseudogramma polyacanthum* polyacanthum (Bleeker, 1856)

*Suttonia lineata* Gosline, 1960

**CALLANTHIIDAE**

*Callanthias australis* Ogilby, 1900

*Grammatonotus laysanus* Gilbert, 1905

*Grammatonotus surugaensis* Katayama, Yamakawa & Suzuki, 1980

**PSEUDOCHROMIDAE**

*Cypho purpurascens* (DeVis, 1884) (Reported by Fourmanoir & Laboute [1976: 285] under the name *Pseudochromis mccullochi*, and by Laboute & Grandperrin [2000: 205] as 'Pseudochromidae indéterminée')

*Ogilbyina salvati* (Plessis & Fourmanoir, 1966) (Reported by Whitley [1961: 65] under the name *Pseudochromis novaehollandiae*)

*Pictichromis coralensis* Gill, 2004 (Reported by several authors under the name *Pseudochromis paccagnellae*)

*Pseudochromis cyanotaenia* Bleeker, 1857

*Pseudochromis fuscus* Müller & Troschel, 1849 (Synonym: *Pseudochromis aurea* Seale, 1910)

*Pseudochromis jamesi* Schultz, 1943

*Pseudochromis kolythrus* Gill & Winterbottom, 1993

*Pseudochromis marshallensis* Schultz in Schultz, Herald, Lachner, Welander & Woods, 1953

*Pseudochromis tapeinosoma* Bleeker, 1853

*Pseudoplesiops howensis* Allen, 1987

*Pseudoplesiops immaculatus* Gill & Edwards, 2002

*Pseudoplesiops multisquamatus* Allen, 1987

*Pseudoplesiops rosae* Schultz, 1943

**PLESIOPIDAE**

A record of *Plesiops corallicola* (Bleeker, 1853) by Rivaton *et al.* (1989: 61) needs verification

*Assessor macneilli* Whitley, 1935

*Belonepterygion fasciolatum* (Ogilby, 1889)

*Callopleiops altivelis* (Steindachner, 1903)

*Plesiops coeruleolineatus* Rüppell, 1835 (Synonym: *Plesiops melas* Bleeker, 1849)

*Plesiops insularis* Mooi & Randall, 1991 (Reported by Rivaton *et al.* [1989: 61] under the name *Plesiops oxycephalus*)

*Plesiops verecundus* Mooi, 1995

**OPISTOGNATHIDAE**

*Opistognathus* sp. 1 (To be described by W.F. Smith-Vaniz)

*Opistognathus* sp. 2 (To be described by W.F. Smith-Vaniz)

*Stalix histrio* Jordan & Snyder, 1902

**BANJOSIDAE**

*Banjos banjos* (Richardson, 1846)

**PRIACANTHIDAE**

*Cookeolus japonicus* (Cuvier, 1829)

*Heteropriacanthus cruentatus* (Lacepède, 1802)

*Priacanthus hamrur* (Forsskål in Niebuhr, 1775) (Synonym: *Priacanthus longipinnis* Borodin, 1932)

*Priacanthus macracanthus* Cuvier, 1829

*Priacanthus sagittarius* Starnes, 1988

*Pristigenys niphonia* (Cuvier, 1829)

## APOGONIDAE

A record of *Ostorrhinchus poecilopterus* Cuvier, 1828 by Rivaton *et al.* (1989) under the name *Apogon poecilopterus* needs verification.

*Apogon amboinensis* Bleeker, 1853  
*Apogon caudicinctus* Randall & Smith, 1988  
*Apogon crassiceps* Garman, 1903 (Reported by Plessis & Fourmanoir [1966: 128] under the name *Apogon erythrinus*, and by other authors under the name *Apogon coccineus*)  
*Apogon doryssa* (Jordan & Seale, 1906)  
*Apogon ellioti* Day, 1878  
*Apogon fasciatus* (White, 1790)  
*Apogon gilberti* (Jordan & Seale, 1905)  
*Apogon hyalosoma* Bleeker, 1853  
*Apogon indicus* Greenfield, 2001 (Reported by Kulbicki & Williams [1997: 14] under the name *Apogon erythrinus*)  
*Apogon melas* Bleeker, 1848  
*Apogon notatus* (Houttuyn, 1782)  
*Apogon noumeae* Whitley, 1958  
*Apogon semiornatus* Peters, 1876  
*Apogon susanae* Greenfield, 2001  
*Apogon talboti* Smith, 1961  
*Apogonichthys ocellatus* (Weber, 1913)  
*Apogonichthys perdist* Bleeker, 1854  
*Archamia fucata* (Cantor, 1850)  
*Archamia leai* Waite, 1916  
*Archamia macroptera* (Cuvier in Cuvier & Valenciennes, 1828) (Reported by several authors under the name *Archamia lineolata* or *A. lineolatus*; **new record** from New Caledonia)  
*Archamia zosterophora* (Bleeker, 1856)  
*Cercamia cladara* Randall & Smith, 1988  
*Cheilodipterus artus* Smith 1961  
*Cheilodipterus isostigmus* (Schultz, 1940)  
*Cheilodipterus macrodon* (Lacepède, 1802) (Reported by several authors under the name *Cheilodipterus lachneri*)  
*Cheilodipterus quinquelineatus* Cuvier in Cuvier & Valenciennes, 1828  
*Cheilodipterus singapurensis* Bleeker, 1859 (Synonym: *Cheilodipterus subulatus* Weber, 1909; **new record** from New Caledonia)  
*Foa fo* Jordan & Seale, 1906 (Reported by several authors under the name *Foa brachygramma*, and by Laboute & Grandperrin [2000: 219] as 'Apogonidae indéterminé 1')  
*Fowleria aurita* (Valenciennes, 1831)  
*Fowleria isostigma* (Jordan & Seale, 1906)  
*Fowleria marmorata* (Alleyne & Macleay, 1877)  
*Fowleria vaiulae* (Jordan & Seale, 1906) (Synonym: *Fowleria abocellata* Goren & Karplus, 1980; reported by Laboute & Grandperrin [2000: 218] under the name *Fowleria* sp.)  
*Fowleria variegata* (Valenciennes, 1832)  
*Gymnapogon philippinus* (Herre, 1939)  
*Gymnapogon urospilotus* Lachner in Schultz, Herald, Lachner, Welander & Woods, 1953  
*Neamia catalai* (Fourmanoir, 1973)  
*Neamia octospina* Smith & Radcliffe in Radcliffe, 1912  
*Ostorrhinchus angustatus* (Smith & Radcliffe, 1911)  
*Ostorrhinchus apogonides* (Bleeker, 1856)  
*Ostorrhinchus aureus* (Lacepède, 1802)  
*Ostorrhinchus bandanensis* (Bleeker, 1854) (Reported by Laboute & Grandperrin [2000: 216] under the name *Apogon* sp. 6)  
*Ostorrhinchus capricornis* (Allen & Randall, 1993)  
*Ostorrhinchus compressus* (Smith & Radcliffe in Radcliffe, 1911)  
*Ostorrhinchus cookii* (Macleay, 1881)

- Ostorrhinchus cyanosoma* (Bleeker, 1853)
- Ostorrhinchus diversus* (Smith & Radcliffe in Radcliffe 1912)
- Ostorrhinchus doederleini* (Jordan & Snyder, 1901)
- Ostorrhinchus endekataenia* (Bleeker, 1852)
- Ostorrhinchus euspilotus* (Fraser, 2006)
- Ostorrhinchus flavus* (Allen & Randall, 1993)
- Ostorrhinchus fuscus* (Quoy & Gaimard, 1825)
- Ostorrhinchus guamensis* (Bleeker, 1856) (Synonym: *Apogon nubilus* Garman, 1903; reported by Whitley [1961: 64] under the name *Aspiscis savayensis*, by Rivaton *et al.* [1989: 24] as *Apogon ocellatus*, and by Laboute & Grandperrin [2000: 219] as 'Apogonidae indéterminé 2')
- Ostorrhinchus kiensis* (Jordan & Snyder, 1901) (Reported by Laboute & Grandperrin [2000: 215] under the name *Apogon* sp. 3)
- Ostorrhinchus lateralis* (Valenciennes, 1832) (Synonym: *Apogon ceramensis* Bleeker, 1852)
- Ostorrhinchus nigrofasciatus* (Lachner in Schultz, Herald, Lachner, Welander & Woods, 1953)
- Ostorrhinchus norfolkensis* (Ogilby, 1888) (Reported by Laboute & Grandperrin [2000: 215] under the names *Apogon* sp. 2 and *Apogon* sp. 4)
- Ostorrhinchus novemfasciatus* (Cuvier in Cuvier & Valenciennes, 1828)
- Ostorrhinchus rubrimacula* (Randall & Kulbicki, 1988)
- Ostorrhinchus savayensis* (Günther, 1872)
- Ostorrhinchus sealei* (Fowler, 1918) (New record from New Caledonia based on SMNS material from southern Grande Terre)
- Ostorrhinchus selas* (Randall & Hayashi, 1990)
- Ostorrhinchus septemstriatus* (Günther, 1880)
- Ostorrhinchus taeniophorus* (Regan, 1905)
- Pristiapogon exostigma* (Jordan & Starks, 1906)
- Pristiapogon fraenatus* (Valenciennes, 1832)
- Pristiapogon kallopterus* (Bleeker, 1856) (Reported by Fourmanoir & Laboute [1976: 290] under the names *Apogon* sp. and *Apogon snyderi*; reported by Laboute & Grandperrin [2000: 216] under the name *Apogon* sp. 7)
- Pristiapogon taeniopterus* (Bennett, 1835)
- Pristicon trimaculatus* (Cuvier in Cuvier & Valenciennes, 1828) (Reported by Rivaton *et al.* [1989: 24] under the names *Apogon taeniatus* and *A. koilomatodon*)
- Pseudamia gelatinosa* Smith, 1955 (Reported by Laboute & Grandperrin [2000: 218] under the name *Pseudamia* sp.)
- Pseudamia zonata* Randall, Lachner & Fraser, 1985
- Pseudamiops gracilicauda* (Lachner in Schultz, Herald, Lachner, Welander & Woods, 1953)
- Rhabdamia cypselura* Weber, 1909 (Reported by Laboute & Grandperrin [2000: 66] under the name 'Apogonidae indéterminé')
- Rhabdamia gracilis* (Bleeker, 1856)
- Siphania versicolor* (Smith & Radcliffe, 1909) (Reported by Laboute & Grandperrin [2000: 218] under the name *Siphania* sp.)
- Sphaeramia nematoptera* (Bleeker, 1856)
- Sphaeramia orbicularis* (Kuhl & Hasselt in Cuvier & Valenciennes, 1828)
- Zoramia fragilis* (Smith, 1961) (Reported by Laboute & Grandperrin [2000: 215] under the name *Apogon* sp. 1)
- Zoramia leptacantha* (Bleeker, 1856) (reported by several authors under the name *Apogon graeffei*)
- SILLAGINIDAE**
- Sillago ciliata* Cuvier in Cuvier & Valenciennes, 1829 (Synonym: *Sillago insularis* Castelnau, 1873)
- Sillago sihama* (Forsskål in Niebuhr, 1775)
- MALACANTHIDAE**
- Branchiostegus wardi* Whitley, 1932
- Hoplolatilus chlupatyi* Klausewitz, McCosker, Randall & Zetsche, 1978
- Hoplolatilus cuniculus* Randall & Dooley, 1974
- Hoplolatilus fronticinctus* Günther, 1887
- Hoplolatilus starcki* Randall & Dooley, 1974

*Malacanthus brevirostris* Guichenot, 1848 (Synonyms: *Malacanthus hoedti* Bleeker, 1859; *Dikellorhynchus incredibilis* Smith, 1956)

*Malacanthus latovittatus* (Lacepède, 1802)

## CORYPHAEINIDAE

*Coryphaena equiselis* Linnaeus, 1758

*Coryphaena hippurus* Linnaeus, 1758

## ECHENEIDAE

*Echeneis naucrates* Linnaeus, 1758

*Phtheirichthys lineata* (Menzies, 1791)

*Remora osteochir* (Cuvier, 1829)

*Remorina albescens* (Temminck & Schlegel, 1845)

## CARANGIDAE

Records of *Carangoides malabaricus* (Bloch & Schneider, 1801) by Jouan (1879: 334), *Trachinotus rhomboides* by Borodn (1932: 77) of *Caranx malabaricus* and of *Decapterus macrosoma* Bleeker, 1951 by Rivaton *et al.* [1989: 31] needs verification.

*Alectis ciliaris* (Bloch, 1787) (Reported by several authors under the name *Alectis indicus*)

*Alepes apercna* Smith-Vaniz in Grant, 1987 (Reported by Fourmanoir & Laboute [1976: 176] under the name *Alepes kalla*, and by several authors under the name *Alepes djeddaba*)

*Alepes vari* (Cuvier, 1833)

*Atule mate* (Cuvier in Cuvier & Valenciennes, 1833)

*Carangoides coeruleopinnatus* (Rüppell, 1830) (Synonyms: *Carangoides caeruleopinnatus*; *Carangoides uii*)

*Carangoides chrysophrys* (Cuvier in Cuvier & Valenciennes, 1833)

*Carangoides dinema* Bleeker, 1851

*Carangoides ferdau* (Forsskål in Niebuhr, 1775) (Synonym: *Carangoides gilberti*)

*Carangoides fulvoguttatus* (Forsskål in Niebuhr, 1775) (Synonym: *Carangoides emburyi*)

*Carangoides gymnostethus* (Cuvier in Cuvier & Valenciennes, 1833)

*Carangoides hedlandensis* (Whitley, 1934) (Reported by Fourmanoir [1971: 113] under the name *Caranx armatus*).

*Carangoides orthogrammus* Jordan & Gilbert, 1881

*Carangoides plagiotaenia* (Bleeker, 1857)

*Caranx ignobilis* (Forsskål in Niebuhr, 1775)

*Caranx lugubris* Poey, 1860

*Caranx melampygus* Cuvier in Cuvier & Valenciennes, 1833

*Caranx papuensis* Alleyne & Macleay, 1877 (Synonym: *Caranx celetus* Smith, 1968)

*Caranx sexfasciatus* (Quoy & Gaimard, 1824) (Reported by Plessis & Fourmanoir [1966: 129] under the name *Caranx hippos*)

*Caranx tille* Cuvier, 1833

*Decapterus macarellus* (Valenciennes in Cuvier & Valenciennes, 1833)

*Decapterus muroadsi* (Temminck & Schlegel, 1844)

*Decapterus russelli* (Rüppell, 1829) (Synonym: *Decapterus lajang* Bleeker, 1855)

*Decapterus tabl* Berry, 1967

*Elagatis bipinnulata* (Quoy & Gaimard, 1825)

*Gnathanodon speciosus* (Forsskål in Niebuhr, 1775)

*Megalaspis cordyla* (Linnaeus, 1758)

*Naucrates ductor* (Linnaeus, 1758)

*Pseudocaranx dentex* (Bloch & Schneider, 1801) (Synonym: *Carangoides georgianus*)

*Scomberoides commersonianus* Lacepède, 1801

*Scomberoides lysan* (Forsskål in Niebuhr, 1775)

*Scomberoides tol* (Cuvier in Cuvier & Valenciennes, 1832)

*Selar boops* (Cuvier in Cuvier & Valenciennes, 1833)

*Selar crumenophthalmus* (Bloch, 1793)

*Seriola dumerili* (Risso, 1810)

*Seriola lalandi* Valenciennes in Cuvier & Valenciennes, 1833 (Synonym: *Seriola aureovittata*)

*Seriola rivoliana* Valenciennes in Cuvier & Valenciennes, 1833

*Trachinotus anak* Ogilby, 1909  
*Trachinotus baillonii* (Lacepède, 1801)  
*Trachinotus blochii* (Lacepède, 1801)  
*Uraspis uraspis* (Günther, 1860)

#### MENIDAE

A record of *Mene maculata* (Bloch & Schneider, 1801) by Rivaton *et al.* (1989: 52) needs verification.

#### LEIOGNATHIDAE

A record of *Leiognathus lineolatus* (Valenciennes in Cuvier & Valenciennes, 1835) by Rivaton *et al.* (1989: 48) needs verification.  
*Gazza minuta* (Bloch, 1795) (Synonym: *Gazza equulaeformis* Rüppell, 1835)  
*Leiognathus bindus* (Valenciennes in Cuvier & Valenciennes, 1835)  
*Leiognathus equulus* (Forsskål in Niebuhr, 1775)  
*Leiognathus fasciatus* (Lacepède, 1803)  
*Leiognathus leuciscus* (Günther, 1860)  
*Leiognathus splendens* (Cuvier, 1829)  
*Photoplacios rivulatus* (Temminck & Schlegel, 1845)  
*Secutor insidiator* (Bloch, 1787)  
*Secutor ruconius* (Hamilton Buchanan, 1822)

#### BRAMIDAE

*Brama dussumieri* Cuvier, 1831  
*Brama myersi* Mead, 1972  
*Brama orcinii* Cuvier in Cuvier & Valenciennes, 1831 (Synonym: *Collybus drachme*)  
*Pteraclis aesticola* (Jordan & Snyder, 1901)  
*Pterycombus petersii* (Hilgendorf, 1878)  
*Taractes asper* Lowe, 1843  
*Taractes rubescens* (Jordan & Jordan, 1887)  
*Taractichthys steindachneri* (Döderlein, 1883) (Reported by several authors under the name  
    *Taractichthys longipinnis*)

#### EMMELICHTHYIDAE

*Emmelichthys nitidus nitidus* Richardson, 1845  
*Erythrocles taeniatus* Randall & Rivaton, 1992 (Reported by several authors under the name  
    *Erythrocles schlegelii*)

#### LUTJANIDAE

Records of *Lutjanus erythropterus* Bloch, 1790 and *Paracaesio gonzalezi* Fourmanoir & Rivaton, 1979 by Rivaton *et al.* (1989: 49) need verification.  
*Aphareus furca* (Lacepède, 1801)  
*Aphareus rutilans* Cuvier in Cuvier & Valenciennes, 1830  
*Aprion virescens* Valenciennes in Cuvier & Valenciennes, 1830  
*Etelis carbunculus* Cuvier in Cuvier & Valenciennes, 1828  
*Etelis coruscans* Valenciennes, 1862 (Synonym: *Etelis oculatus lifuensis* Fourmanoir, 1971)  
*Etelis radiosus* Anderson, 1981  
*Lutjanus adetii* (Castelnau, 1873) (Synonyms: *Lutjanus amabilis*; *Lutjanus parvavitta* Postel, 1965)  
*Lutjanus argenticulatus* (Forsskål in Niebuhr, 1775)  
*Lutjanus bohar* (Forsskål in Niebuhr, 1775)  
*Lutjanus fulviflamma* (Forsskål in Niebuhr, 1775)  
*Lutjanus fulvus* (Bloch & Schneider, 1801)  
*Lutjanus fuscescens* (Valenciennes in Cuvier & Valenciennes, 1830)  
*Lutjanus gibbus* (Forsskål in Niebuhr, 1775)  
*Lutjanus kasmira* (Forsskål in Niebuhr, 1775)  
*Lutjanus lutjanus* (Bloch, 1790) (Reported by Fourmanoir & Laboute [1976: 76] under the name  
    *Lutjanus lineolatus*)  
*Lutjanus malabaricus* (Bloch & Schneider, 1801)  
*Lutjanus monostigma* (Cuvier in Cuvier & Valenciennes, 1828)  
*Lutjanus quinquelineatus* Bloch, 1790  
*Lutjanus rivulatus* (Cuvier in Cuvier & Valenciennes, 1828)

*Lutjanus rufolineatus* (Valenciennes in Cuvier & Valenciennes, 1830) (Reported by several authors under the name *Lutjanus boutton*; new record from New Caledonia)

*Lutjanus russelli* (Bleeker, 1849)

*Lutjanus sebae* (Cuvier, 1816)

*Lutjanus semicinctus* Quoy & Gaimard, 1824

*Lutjanus vitta* (Quoy & Gaimard, 1824)

*Macolor macularis* Fowler, 1931

*Macolor niger* (Forsskål in Niebuhr, 1775)

*Paracaesio caerulea* (Katayama, 1934)

*Paracaesio kusakarii* Abe, 1960

*Paracaesio sordida* Abe & Shinohara, 1962

*Paracaesio xanthura* (Bleeker, 1869)

*Pristipomoides argyrogrammicus* (Valenciennes, 1831)

*Pristipomoides filamentosus* (Valenciennes in Cuvier & Valenciennes, 1830)

*Pristipomoides flavipinnis* Shinohara, 1963

*Pristipomoides multidens* (Day, 1870) (Reported by several authors under the name *Pristipomoides typus*)

*Pristipomoides zonatus* (Valenciennes in Cuvier & Valenciennes, 1830)

*Syphorichthys spilurus* (Günther, 1874)

*Syphorus nematophorus* (Bleeker, 1860)

## CAESIONIDAE

*Caesio caerulea* Lacepède, 1802

*Caesio cuning* (Bloch, 1791) (Synonym: *Caesio erythrogaster* Cuvier in Cuvier & Valenciennes, 1830)

*Caesio lunaris* Ehrenberg & Cuvier in Cuvier in Cuvier & Valenciennes, 1830

*Caesio teres* Seale, 1906 (Synonym: *Caesio pulcherrimus*; reported by Rivaton *et al.* [1989: 30] under the name *Caesio xanthonota*)

*Dipteronotus balteatus* (Valenciennes in Cuvier & Valenciennes, 1830)

*Gymnoaesio gymnoptera* (Bleeker, 1856)

*Pterocaesio chrysazona* (Cuvier in Cuvier & Valenciennes, 1830)

*Pterocaesio digramma* (Bleeker, 1865)

*Pterocaesio marri* Schultz in Schultz, Herald, Lachner, Welander & Woods, 1953 (Reported by Laboute & Grandperrin [2000: 259] under the name *Pterocaesio* sp. 1)

*Pterocaesio pisang* (Bleeker, 1853)

*Pterocaesio tessellata* Carpenter, 1987

*Pterocaesio tile* (Cuvier in Cuvier & Valenciennes, 1830)

*Pterocaesio trilineata* Carpenter, 1987

## LOBOTIDAE

*Lobotes surinamensis* (Bloch, 1790)

## GERREIDAE

*Gerres filamentosus* Cuvier, 1829

*Gerres longirostris* (Lacepède, 1801) (Synonym: *Gerres acinaces*)

*Gerres oblongus* Cuvier in Cuvier & Valenciennes, 1830 (Synonym: *Gerres macrosoma*)

*Gerres ovatus* Günther, 1859

*Gerres oyena* (Forsskål in Niebuhr, 1775) (Synonym: *Gerres argyreus*)

## HAEMULIDAE

*Diagramma pictum* (Thunberg, 1792) (Reported by Borodin [1932: 83] under the name *Diagramma punctatum*)

*Plectorhinchus albovittatus* (Rüppell, 1835) (Synonym: *Plectorhinchus harrawayi*, taxonomic decision of Randall & Johnson ([2000: 479]).

*Plectorhinchus chaetodonoides* Lacepède, 1800

*Plectorhinchus chrysotaenia* (Bleeker, 1855)

*Plectorhinchus flavomaculatus* (Ehrenberg in Cuvier & Valenciennes, 1830) (Reported by Whitley [1961: 65] under the name *Plectorhinchus roughleyi*)

*Plectorhinchus gibbosus* (Lacepède, 1802)

*Plectorhinchus lessonii* (Cuvier in Cuvier & Valenciennes, 1830) (Reported by Rivaton *et al.*)

[1989: 43] under the name *Plectorhinchus diagrammus*)

*Plectorhinchus lineatus* (Linnaeus, 1758) (Synonym: *Plectorhinchus goldmanni*)

*Plectorhinchus obscurus* (Günther, 1871)

*Plectorhinchus picus* (Cuvier in Cuvier & Valenciennes, 1830)

*Plectorhinchus vittatus* (Linnaeus, 1758) (Synonym: *Plectorhinchus orientalis*)

*Pomadasys argenteus* (Forsskål in Niebuhr, 1775) (Synonym: *Pomadasys hasta*; reported by Borodin [1932: 83] under the name *Pristipoma operculare*)

#### NEMIPTERIDAE

A record of *Parascolopsis inermis* (Temminck & Schlegel, 1843) from New Caledonia by Rivaton *et al.* (1989: 57) needs verification.

*Nemipterus balinensisoides* (Popa, 1918) (Reported by Borodin [1932: 84] under the name *Synagris striatus*)

*Nemipterus furcosus* (Valenciennes in Cuvier & Valenciennes, 1830) (Reported by Laboute & Grandperrin [2000: 274, 275] under the name *Nemipterus peronii*)

*Nemipterus peronii* (Valenciennes in Cuvier & Valenciennes, 1830) (Synonym: *Dentex tolu*)

*Nemipterus zyson* (Bleeker, 1856) (Synonym: *Nemipterus metopias*)

*Pentapodus aureofasciatus* Russell, 2001 (Reported by various authors under the names

*Pentapodus microdon* [non Bleeker, 1853] and *Pentapodus nagasakiensis* [non Tanaka, 1915])

*Pentapodus caninus* (Cuvier in Cuvier & Valenciennes, 1830) (Synonym: *Pentapodus macrurus*)

*Pentapodus paradiseus* (Günther, 1859)

*Scolopsis affinis* Peters, 1877 (Reported by Fourmanoir [1971: 112] under the name *Scolopsis personatus*)

*Scolopsis bilineata* (Bloch, 1793) (Reported by Whitley [1961: 65] under the name *Scolopsis cancellatus*)

*Scolopsis ciliata* (Lacepède, 1802)

*Scolopsis lineatus* Quoy & Gaimard, 1824

*Scolopsis taeniopterus* (Kuhl & Hasselt in Cuvier & Valenciennes, 1830)

*Scolopsis temporalis* (Cuvier in Cuvier & Valenciennes, 1830) (Reported by Randall, Allen & Steene [1997: 207] under the name *Scolopsis monogramma* [part].)

*Scolopsis trilineatus* Kner, 1868

#### LETHRINIDAE

*Gnathodentex aureolineatus* (Lacepède, 1802)

*Gymnocranius* sp. (Undescribed species, reported by several authors under the name *Gymnocranius lethrinoides* [non Bleeker, 1850])

*Gymnocranius audleyi* Ogilby, 1916 (Synonym: *Gymnocranius bitorquatus*)

*Gymnocranius elongatus* Senta, 1973

*Gymnocranius euanus* Günther, 1879 (Synonym: *Gymnocranius japonicus* Akazaki, 1961)

*Gymnocranius grandoculis* (Valenciennes in Cuvier & Valenciennes, 1830) (Synonym:

*Gymnocranius rivulatus*; reported by Kulbicki [1988: 306] under the name *Gymnocranius robertsi*)

*Lethrinus atkinsoni* (Seale, 1909) (Reported by several authors under the name *Lethrinus mahsena*; reported by Laboute & Grandperrin [2000: 272] under the name *Lethrinus* sp.)

*Lethrinus erythracanthus* Valenciennes in Cuvier & Valenciennes, 1830 (Synonym: *Lethrinus kallopterus* Bleeker, 1856)

*Lethrinus genivittatus* Valenciennes in Cuvier & Valenciennes, 1830 (Frequently used synonym: *Lethrinus nematacanthus*)

*Lethrinus harak* (Forsskål in Niebuhr, 1775) (Reported by Whitley [1961: 64] under the name *Lethrinus glyphodon*)

*Lethrinus laticaudis* Alleyne & Macleay, 1877 (Synonym: *Lethrinus anarhynchus* Postel, 1965)

*Lethrinus lentjan* (Lacepède, 1802)

*Lethrinus miniatus* (Forster in Bloch & Schneider, 1801) (Synonym: *Lethrinus chrysostomus* Richardson, 1848)

*Lethrinus nebulosus* (Forsskål in Niebuhr, 1775) (Reported by Borodin [1932: 83] under the name *Lethrinus hematopterus* [non Temminck & Schlegel, 1844])

*Lethrinus obsoletus* (Forsskål in Niebuhr, 1775) (Synonym: *Lethrinus ramak*)

*Lethrinus olivaceus* Valenciennes in Cuvier & Valenciennes, 1830 (Synonym: *Lethrinus rostratus* Valenciennes in Cuvier & Valenciennes, 1830)

- Lethrinus ravus* Carpenter & Randall, 2003  
*Lethrinus rubrioperculatus* Sato, 1978  
*Lethrinus semicinctus* Valenciennes in Cuvier & Valenciennes, 1830  
*Lethrinus variegatus* Ehrenberg in Valenciennes in Cuvier & Valenciennes, 1830  
*Lethrinus xanthochilus* Klunzinger, 1870  
*Monotaxis grandoculis* (Forsskål in Niebuhr, 1775)  
*Monotaxis heterodon* (Bleeker, 1854)  
*Wattsia mossambica* (Smith, 1957)

#### SPARIDAE

- Acanthopagrus berda* (Forsskål in Niebuhr, 1775)

#### POLYNEMIDAE

- Eleutheronema tetradactylum* (Shaw, 1804)  
*Polydactylus microstomus* (Bleeker, 1851)  
*Polydactylus plebeius* (Broussonet, 1782)  
*Polydactylus sexfilis* (Valenciennes in Cuvier & Valenciennes, 1831)

#### MULLIDAE

A record of *Mulloidichthys pfluegeri* (Steindachner, 1901) from New Caledonia by Rivaton *et al.* (1989: 53) under the name *Mulloides pflugeri* needs verification.

- Mulloidichthys flavolineatus* (Lacepède, 1801) (Synonym: *Mulloides samoensis*)  
*Mulloidichthys vanicolensis* (Valenciennes in Cuvier & Valenciennes, 1831)  
*Parupeneus barberinoides* (Bleeker, 1852)  
*Parupeneus barberinus* (Lacepède, 1801)  
*Parupeneus ciliatus* (Lacepède, 1801) (Reported by Fourmanoir & Laboute [1976: 199] under the name *Parupeneus porphyreus*, and by several authors under the name *Parupeneus dispilurus* or *Parupeneus dispirlus*)  
*Parupeneus crassilabris* (Valenciennes in Cuvier & Valenciennes, 1831) (Reported by several authors under the name *Parupeneus bifasciatus*)  
*Parupeneus cyclostomus* (Lacepède, 1802) (Synonym: *Parupeneus chryserydros*)  
*Parupeneus heptacanthus* (Lacepède, 1801) (Synonym: *Parupeneus pleurospilos*; reported by Borodin [1932: 84] under the name *Upeneus cinnaborinus*)  
*Parupeneus indicus* (Shaw, 1803) (Reported by Whitley [1961: 65] under the name *Pseudupeneus filamentosus*)  
*Parupeneus multifasciatus* (Quoy & Gaimard, 1824) (Reported by several authors under the name *Parupeneus trifasciatus*)  
*Parupeneus pleurostigma* (Bleeker, 1853)  
*Parupeneus spilurus* (Bleeker, 1854) (Synonyms: *Parupeneus signatus*; *Pseudupeneus jeffi* Ogilby, 1908; reported by several authors under the name *Parupeneus signatus* [Günther, 1867], with the latter name considered as valid for the New Caledonian populations by Kuiter [1993], which was not accepted by subsequent authors like Randall [2005: 299])  
*Upeneus australiae* Kim & Nakaya, 2002  
*Upeneus filifer* (Ogilby, 1910)  
*Upeneus moluccensis* (Bleeker, 1855)  
*Upeneus mouthami* Randall & Kulbicki, 2005  
*Upeneus sulphureus* Cuvier in Cuvier & Valenciennes, 1829 (Reported by Rivaton *et al.* [1989: 53] under the name *Upeneus bensasi*)  
*Upeneus tragula* Richardson, 1845  
*Upeneus vittatus* Lacepède, 1801

#### PEMPHERIDAE

- Parapriacanthus dispar* (Herre, 1935)  
*Parapriacanthus marei* Fourmanoir, 1971  
*Parapriacanthus ransonneti* Steindachner, 1870 (Reported by Fourmanoir & Laboute [1976: 292-293] under the name *Parapriacanthus beryciformes*)  
*Pempheris oualensis* Cuvier in Cuvier & Valenciennes, 1831 (Synonym: *Pempheris otaitensis*)  
*Pempheris schwenkii* Bleeker, 1855

## MONODACTYLIDAE

*Monodactylus argenteus* (Linnaeus, 1758)

## KYPHOSIDAE

*Kyphosus cinerascens* (Forsskål in Niebuhr, 1775)

*Kyphosus pacificus* Sakai & Nakabo, 2004 (Reported by several authors from Grande Terre under the name *Kyphosus bigibbus*; reported by Laboute & Grandperrin [2000: 288] under the name *Kyphosus* sp. 1; **new record** from New Caledonia)

*Kyphosus sydneyanus* (Günther, 1886) (Reported by Laboute & Grandperrin [2000: 289] under the name *Kyphosus* sp. 2)

*Kyphosus vaigiensis* (Quoy & Gaimard, 1825)

*Microcanthus strigatus* (Cuvier in Cuvier & Valenciennes, 1831)

## DREPANIDAE

*Drepane punctata* (Linnaeus, 1758)

## CHAETODONTIDAE

*Amphichactodon howensis* (Waite, 1903)

*Chaetodon auriga* Forsskål in Niebuhr, 1775

*Chaetodon baronessa* Cuvier in Cuvier & Valenciennes, 1831

*Chaetodon bennetti* Cuvier in Cuvier & Valenciennes, 1831

*Chaetodon citrinellus* Cuvier in Cuvier & Valenciennes, 1831

*Chaetodon ephippium* Cuvier in Cuvier & Valenciennes, 1831

*Chaetodon flavirostris* Günther, 1874

*Chaetodon guentheri* Ahl, 1923

*Chaetodon kleinii* Bloch, 1790

*Chaetodon lineolatus* Cuvier in Cuvier & Valenciennes, 1831

*Chaetodon lunula* (Lacepède, 1802)

*Chaetodon lunulatus* Quoy & Gaimard, 1825 (Reported by several authors under the name *Chaetodon trifasciatus*)

*Chaetodon melanotus* Bloch & Schneider, 1801

*Chaetodon mertensi* Cuvier in Cuvier & Valenciennes, 1831 (Reported by Whitley [1961: 64] under the name *Chaetodon dixsoni*)

*Chaetodon meyeri* Bloch & Schneider, 1801

*Chaetodon ornatissimus* Cuvier in Cuvier & Valenciennes, 1831

*Chaetodon pelewensis* Kner, 1868

*Chaetodon plebeius* Cuvier in Cuvier & Valenciennes, 1831

*Chaetodon rafflesii* Bennett, 1830

*Chaetodon reticulatus* Cuvier in Cuvier & Valenciennes, 1831

*Chaetodon semeion* Bleeker, 1855

*Chaetodon speculum* Cuvier in Cuvier & Valenciennes, 1831

*Chaetodon trifascialis* (Quoy & Gaimard, 1824)

*Chaetodon ulietensis* Cuvier in Cuvier & Valenciennes, 1831 (Reported by Plessis & Fourmanoir [1966: 130] under the name *Chaetodon falcula*):

*Chaetodon unimaculatus* Bloch, 1787

*Chaetodon vagabundus* Linnaeus, 1758

*Coradion altivelis* McCulloch, 1916

*Forcipiger flavissimus* Jordan & McGregor, 1898

*Forcipiger longirostris* (Broussonet, 1782)

*Hemitaurichthys polylepis* (Bleeker, 1857) (Reported by Whitley [1961: 64] under the name *Hemitaurichthys zoster*)

*Heniochus acuminatus* (Linnaeus, 1758)

*Heniochus chrysostomus* Cuvier in Cuvier & Valenciennes, 1831 (Reported by Whitley [1961: 64] and Fourmanoir & Laboute [1976: 228-229] under the name *Heniochus permutteratus*)

*Heniochus monoceros* Cuvier in Cuvier & Valenciennes, 1831

*Heniochus singularis* Smith & Radcliffe, 1911

*Heniochus varius* (Cuvier in Cuvier & Valenciennes, 1829)

*Prognathodes guyotensis* (Yamamoto & Tameka in Okamura *et al.*, 1982)

## POMACANTHIDAE

- Apolemichthys trimaculatus* (Cuvier in Cuvier & Valenciennes, 1831)  
*Centropyge bicolor* (Bloch, 1787)  
*Centropyge bispinosa* (Günther, 1860)  
*Centropyge flavicauda* Fraser-Brunner, 1933  
*Centropyge flavissima* (Cuvier in Cuvier & Valenciennes, 1831)  
*Centropyge heraldi* Woods & Schultz in Schultz, Herald, Lachner, Welander & Woods, 1953  
*Centropyge loricula* (Günther, 1874)  
*Centropyge multifasciata* (Smith & Radcliffe, 1911)  
*Centropyge nigriocella* Woods & Schultz in Schultz, Herald, Lachner, Welander & Woods, 1953  
*Centropyge nox* (Bleeker, 1853)  
*Centropyge tibicen* (Cuvier in Cuvier & Valenciennes, 1831)  
*Centropyge vrolikii* (Bleeker, 1853)  
*Chaetodontoplus conspicillatus* (Waite, 1900)  
*Genicanthus melanospilos* (Bleeker, 1857)  
*Genicanthus watanabei* (Yasuda & Tominaga, 1970)  
*Pomacanthus annularis* (Bloch, 1787)  
*Pomacanthus imperator* (Bloch, 1787)  
*Pomacanthus semicirculatus* (Cuvier in Cuvier & Valenciennes, 1831)  
*Pomacanthus sexstriatus* (Cuvier in Cuvier & Valenciennes, 1831)  
*Pomacanthus xanthometopon* (Bleeker, 1853)  
*Pygoplites diacanthus* (Boddaert, 1772)

## PENTACEROTIDAE

- Eviotas acutirostris* (Temminck & Schlegel, 1844)  
*Pentaceros decacanthus* Günther, 1883  
*Pseudopentaceros richardsoni* (Smith, 1849)

## TERAPONTIDAE

- Mesopristes kneri* (Bleeker, 1873)  
*Terapon jarbua* (Forsskål in Niebuhr, 1775) (Synonym: *Therapon servus*)  
*Terapon theraps* (Cuvier in Cuvier & Valenciennes, 1829)

## KUHLIIDAE

- Kuhlia marginata* (Cuvier in Cuvier & Valenciennes, 1829)  
*Kuhlia mugil* (Forster in Bloch & Schneider, 1801)  
*Kuhlia munda* (De Vis, 1885) (Synonym: *Kuhlia humilis*)  
*Kuhlia rupestris* (Lacepède, 1802)

## CIRRhitidae

- Amblycirrhitus bimacula* (Jenkins, 1903)  
*Cirrhitichthys falco* Randall, 1963  
*Cirrhitichthys oxycephalus* (Bleeker, 1855)  
*Cirrhitus pinnulatus* (Forster in Bloch & Schneider, 1801)  
*Cyprinocirrhites polyactis* (Bleeker, 1875)  
*Neocirrhites armatus* Castelnau, 1873 (New record from New Caledonia based on AMS IB.7081  
from southern Grande Terre)  
*Oxycirrhites typus* Bleeker, 1857  
*Paracirrhites arcatus* (Parkinson in Cuvier & Valenciennes, 1829)  
*Paracirrhites forsteri* (Bloch & Schneider, 1801)  
*Paracirrhites hemistictus* (Günther, 1874)

## CHEILODACTYLIDAE

- Goniistius francisi* (Burridge, 2004) (Reported from New Caledonia by authors under the name  
*Cheilodactylus vittatus*)  
*Goniistius vestitus* (Castelnau, 1878) (Reported from New Caledonia by authors under the name  
*Cheidodactylus gibbosus*, or *Goniistius gibbosus*)

## CICHLIDAE

- Oreochromis mossambicus* (Peters, 1852) (Introduced species)

## POMACENTRIDAE

A record of *Amblyglyphidodon ternatensis* (Bleeker, 1853) from New Caledonia by Rivaton *et al.* (1989: 62), and records of *Amblypomacentrus breviceps* (Schlegel & Müller, 1839) (under the name *Pomacentrus breviceps*), *Chromis elerae* Fowler & Bean, 1928 and *Chrysiptera unimaculata* (Cuvier in Cuvier & Valenciennes, 1830) (under the name *Abudefdup unimaculatus*) from Nouméa by Borodin (1932: 92, 93), need verification. Kulbicki & Williams (1997: 19) recorded *Stegastes cf. apicalis* from Ouvéa, Loyalty Islands; this also needs verification.

*Abudefdup septemfasciatus* (Cuvier in Cuvier & Valenciennes, 1830)

*Abudefdup sexfasciatus* (Lacepède, 1801) (Synonym: *Glyphisodon coelestinus* Cuvier in Cuvier & Valenciennes, 1830)

*Abudefdup sordidus* (Forsskål in Niebuhr, 1775) (Synonym: *Glyphidodon leucopleura* Day, 1877)

*Abudefdup vaigiensis* (Quoy & Gaimard, 1825) (Reported by Rivaton *et al.* [1989: 62] under the name *Abudefdup saxatilis*)

*Abudefdup whitleyi* Allen & Robertson, 1974

*Amblyglyphidodon aureus* (Cuvier in Cuvier & Valenciennes, 1830)

*Amblyglyphidodon curacao* (Bloch, 1787)

*Amblyglyphidodon leucogaster* (Bleeker, 1847)

*Amblyglyphidodon orbicularis* (Hombron & Jacquinot, 1853) (Reported by Laboute & Grandperrin [2000: 318] from New Caledonia under the name *Amblyglyphidodon* sp.)

*Amphiprion akindynos* Allen, 1972

*Amphiprion chrysopterus* Cuvier in Cuvier & Valenciennes, 1830 (**New record** from New Caledonia based on SMNS material from Maré, Loyalty Islands)

*Amphiprion clarkii* (Bennett, 1830) (Reported by Whitley [1961: 64] under the name *Amphiprion bicinctus*)

*Amphiprion melanopus* Bleeker, 1852 (Reported by Whitley [1961: 64] and subsequent authors under the name *Amphiprion ephippium*)

*Amphiprion perideraion* Bleeker, 1855

*Amphiprion tricinctus* Schultz & Welander in Schultz, 1953 (Reported by several authors from the northern lagoon of Grande Terre and Îles Bélep)

*Cheiloprion labiatus* (Day, 1877) (Reported by Rivaton *et al.* [1989: 63] on the basis of specimen MHN 1980-0347 from Grande Terre)

*Chromis acares* Randall & Swerdlow, 1973

*Chromis agilis* Smith, 1960

*Chromis alpha* Randall, 1988

*Chromis amboinensis* (Bleeker, 1873)

*Chromis analis* (Cuvier in Cuvier & Valenciennes, 1830)

*Chromis atripectoralis* Welander & Schultz, 1951

*Chromis atripes* Fowler & Bean, 1928

*Chromis caudalis* Randall, 1988

*Chromis chrysura* (Bliss, 1883)

*Chromis flavomaculata* Kamohara, 1960 (Synonym: *Chromis kennensis* Whitley, 1964)

*Chromis fumea* (Tanaka, 1917)

*Chromis iomelas* Jordan & Seale, 1906

*Chromis lepidolepis* Bleeker, 1877

*Chromis leucura* Gilbert, 1905

*Chromis margaritifer* Fowler, 1946

*Chromis mirationis* Tanaka, 1917

*Chromis nitida* (Whitley, 1928)

*Chromis notata* (Temminck & Schlegel, 1843)

*Chromis retrofasciata* Weber, 1913

*Chromis ternatensis* (Bleeker, 1856) [*Chromis caerulea* (Cuvier in Cuvier & Valenciennes, 1830)]

is the senior synonym, but was suppressed by the International Commission on Zoological Nomenclature, Opinion 1563]

*Chromis vanderbilti* (Fowler, 1941)

*Chromis viridis* (Cuvier in Cuvier & Valenciennes, 1830)

- Chromis weberi* Fowler & Bean, 1928  
*Chromis xanthochira* (Bleeker, 1851)  
*Chromis xanthura* (Bleeker, 1854)  
*Chrysiptera biocellata* (Quoy & Gaimard, 1825) (Synonym: *Glyphisodon zonatus* Cuvier in Cuvier & Valenciennes, 1830)  
*Chrysiptera brownriggii* (Bennett, 1828) (Synonym: *Chrysiptera leucopoma*; reported from Baie de Saint-Vincent/Grande Terre by Plessis & Fourmanoir [1966: 134] under the name *Pomacentrus albofasciatus*)  
*Chrysiptera cyanea* (Quoy & Gaimard, 1825) (Synonym: *Glyphisodon uniocellatus* Quoy & Gaimard, 1825)  
*Chrysiptera flavipinnis* (Allen & Robertson, 1974)  
*Chrysiptera glauca* (Cuvier in Cuvier & Valenciennes, 1830)  
*Chrysiptera notialis* (Allen, 1975)  
*Chrysiptera rex* (Snyder, 1909)  
*Chrysiptera rollandi* (Whitley, 1961)  
*Chrysiptera starcki* (Allen, 1973)  
*Chrysiptera talboti* (Allen, 1975) (**New record** from New Caledonia based on SMNS 21623 from Maré, Loyalty Islands)  
*Chrysiptera taupou* (Jordan & Seale, 1906)  
*Chrysiptera tricincta* (Allen & Randall, 1974)  
*Dascyllus aruanus* (Linnaeus, 1758)  
*Dascyllus flavicaudus* Randall & Allen, 1977 (**New record** from New Caledonia based on SMNS 24048 from Grand Récif Sud)  
*Dascyllus melanurus* Bleeker, 1854  
*Dascyllus reticulatus* (Richardson, 1846) (Reported by Whitley [1961: 65] under the name *Pelochromis marginatus*)  
*Dascyllus trimaculatus* (Rüppell, 1828)  
*Dischistodus fasciatus* (Cuvier in Cuvier & Valenciennes, 1830) (**New record** from New Caledonia based on SMNS 19727 from southern Grande Terre)  
*Dischistodus prosopotaenia* (Bleeker, 1852) (**New record** from New Caledonia based on SMNS material from southern Grande Terre)  
*Lepidozygus tapeinosoma* (Bleeker, 1856)  
*Neoglyphidodon carlsoni* (Allen, 1975)  
*Neoglyphidodon melas* (Cuvier in Cuvier & Valenciennes, 1830)  
*Neoglyphidodon nigroris* (Cuvier in Cuvier & Valenciennes, 1830)  
*Neoglyphidodon polyacanthus* (Ogilby, 1889)  
*Neopomacentrus anabatoides* (Bleeker, 1847) (Reported from Grande Terre, New Caledonia by Whitley [1961: 65] under the name *Parapomacentrus bankieri*)  
*Neopomacentrus azysron* (Bleeker, 1877)  
*Neopomacentrus bankieri* (Richardson, 1846) (**New record** from New Caledonia based on SMNS 22661 from southern Grande Terre)  
*Neopomacentrus cyanomos* (Bleeker, 1856)  
*Neopomacentrus filamentosus* (Macleay, 1883)  
*Neopomacentrus nemurus* (Bleeker, 1857)  
*Neopomacentrus taeniurus* (Bleeker, 1856)  
*Neopomacentrus violascens* (Bleeker, 1848)  
*Parma polylepis* Günther, 1862  
*Plectroglyphidodon dickii* (Liénard, 1839)  
*Plectroglyphidodon imparipennis* Vaillant & Sauvage, 1875  
*Plectroglyphidodon johnstonianus* Fowler & Ball, 1924  
*Plectroglyphidodon lacrymatus* (Quoy & Gaimard, 1825)  
*Plectroglyphidodon leucozonus* (Bleeker, 1859)  
*Pomacentrus adelus* Allen, 1991  
*Pomacentrus amboinensis* Bleeker, 1868  
*Pomacentrus arenarius* Allen, 1987

*Pomacentrus aurifrons* Allen, 2004 (Reported by several authors including Laboute & Grandperrin [2000: 326] under the name *Pomacentrus smithi*)  
*Pomacentrus bankanensis* Bleeker, 1853  
*Pomacentrus brachialis* Cuvier in Cuvier & Valenciennes, 1830 (Synonym: *Pomacentrus melanopterus* Bleeker, 1852)  
*Pomacentrus chrysurus* Cuvier in Cuvier & Valenciennes, 1830 (Synonym: *Pomacentrus rhodonotus* Bleeker, 1853)  
*Pomacentrus coelestis* Jordan & Starks, 1901  
*Pomacentrus grammorhynchus* Fowler, 1918  
*Pomacentrus imitator* (Whitley, 1964)  
*Pomacentrus lepidogenys* Fowler & Bean, 1928  
*Pomacentrus moluccensis* Bleeker, 1853 (Synonyms: *Pomacentrus popei* Evermann & Seale, 1907; *Pomacentrus sufflavis* Whitley, 1927)  
*Pomacentrus nagasakiensis* Tanaka, 1909 (Synonym: *Pomacentrus arenarius* Allen, 1987 Reported from New Caledonia by Fourmanoir [1981: 27] under the name *Pomacentrus wardi*, and Laboute & Grandperrin [2000: 327] under the name *Pomacentrus* sp.)  
*Pomacentrus nigromanus* Weber, 1913 (New record from New Caledonia based on SMNS 24117 from Lifou, Loyalty Islands)  
*Pomacentrus pavo* (Bloch, 1787)  
*Pomacentrus philippinus* Evermann & Seale, 1907  
*Pomacentrus reidi* Fowler & Bean, 1928  
*Pomacentrus simsianus* Bleeker, 1856  
*Pomacentrus spilotoceps* Randall, 2002 (New record from New Caledonia based on SMNS material from Grande Terre and Lifou/Loyalty Islands).  
*Pomacentrus taeniometopon*: Bleeker, 1852  
*Pomacentrus tripunctatus* Cuvier in Cuvier & Valenciennes, 1830  
*Pomacentrus vaiuli* Jordan & Seale, 1906  
*Pomachromis richardsoni* (Snyder, 1909)  
*Pristotis obtusirostris* (Günther, 1862) (Synonym: *Pristotis jerdoni*; reported from New Caledonia by Laboute & Grandperrin [2000: 327] under the name *Pristotis* sp.)  
*Stegastes albifasciatus* (Schlegel & Müller, 1839) (Synonym: *Pomacentrus eclipticus* Jordan & Seale, 1906)  
*Stegastes apicalis* (DeVis, 1885)  
*Stegastes aureus* (Fowler, 1927)  
*Stegastes fasciolatus* (Ogilby, 1889)  
*Stegastes gascoynei* (Whitley, 1964)  
*Stegastes nigricans* (Lacepède, 1803) (Synonym: *Pomacentrus subniger* DeVis, 1885)  
*Stegastes punctatus* (Quoy & Gaimard, 1825) (Reported from New Caledonia by Myers [1999: 187] under the name *Stegastes lividus*)

## LABRIDAE

Records of *Bodianus unimaculatus* (Günther, 1862) under the name *Bodianus oxycephalus* (non Bleeker, 1862) by Rivaton *et al.* (1989: 45) and *Labroides rubrolabiatus* Randall, 1958 by Fourmanoir & Laboute (1976: 127) need verification.  
*Anampsese caeruleopunctatus* Rüppell, 1828  
*Anampsese femininus* Randall, 1972  
*Anampsese geographicus* Valenciennes in Cuvier & Valenciennes, 1840  
*Anampsese neoguinaicus* Bleeker, 1878  
*Anampsese twistii* Bleeker, 1856  
*Bodianus anthoides* (Bennett, 1831)  
*Bodianus axillaris* (Bennett, 1831)  
*Bodianus bimaculatus* Allen, 1973  
*Bodianus busellatus* Gomon, 2006 (New record from New Caledonia; previously reported from New Caledonia by several authors including Laboute & Grandperrin [2000: 350] and Randall [2005: 392] under the name *Bodianus bilunulatus*)

- Bodianus cylindriatus* (Tanaka, 1930)
- Bodianus dictynna* Gomon, 2006 (Previously reported from New Caledonia by authors under the name *Bodianus diana*)
- Bodianus izuensis* Araga & Yoshino in Masuda, Araga & Yoshino, 1975
- Bodianus loxozonus* (Snyder, 1908) (Reported from New Caledonia by Whitley [1961: 64] under the name *Bodianus hirsutus*)
- Bodianus masudai* Araga & Yoshino, 1975
- Bodianus mesothorax* (Bloch & Schneider, 1801)
- Bodianus paraleucosticticus* Gomon, 2006
- Bodianus perditio* (Quoy & Gaimard, 1824)
- Cheilinus chlorourus* (Bloch, 1791)
- Cheilinus fasciatus* (Bloch, 1791)
- Cheilinus oxycephalus* Bleeker, 1853
- Cheilinus trilobatus* (Lacepède, 1801)
- Cheilinus undulatus* Rüppell, 1835
- Cheilio inermis* (Forsskål in Niebuhr, 1775)
- Choerodon anchorago* (Bloch, 1791)
- Choerodon fasciatus* (Günther, 1867)
- Choerodon graphicus* (De Vis, 1885) (Synonym: *Choerodon transversalis* Whitley, 1956)
- Choerodon jordani* (Snyder, 1908) (Reported from New Caledonia by several authors under the name *Choerodon melanostigma*)
- Choerodon marginatus* Fowler & Bean, 1928
- Cirrhilabrus bathophilus* Randall & Nagareda, 2002
- Cirrhilabrus laboutei* Randall & Lubbock, 1982
- Cirrhilabrus lineatus* Randall & Lubbock, 1982
- Cirrhilabrus punctatus* Randall & Kuiter, 1989
- Cirrhilabrus roseofascia* Randall & Lubbock, 1982
- Coris aygula* Lacepède, 1801 (Synonyms: *Coris cingulum*; *Coris angulatus*; *Coris variegata* Ramsay & Ogilby, 1887)
- Coris batuensis* (Bleeker, 1856) (Synonym: *Coris schroederi*)
- Coris dorsomacula* Fowler, 1908 (Reported from New Caledonia by Rivaton & Bourret [1999: 158] under the name *Coris multicolor*)
- Coris gaimardi* (Quoy & Gaimard, 1824)
- Coris picta* (Bloch & Schneider, 1801)
- Coris pictoides* Randall & Kuiter, 1982
- Coris sandeyeri* (Hector, 1884) (Reported from Nouméa by Whitley [1961: 64] under the name *Guntheria trimaculata*; new record from New Caledonia based on AMS IB.4152).
- Cymolutes praetextatus* (Quoy & Gaimard, 1834) (Reported from New Caledonia by Whitley [1961: 64] under the name *Cymolutes lecluse*)
- Cymolutes torquatus* (Valenciennes in Cuvier & Valenciennes, 1840)
- Diproctacanthus xanthurus* (Bleeker, 1856) (New record from New Caledonia based on AMS IB.7799 from southern Grande Terre)
- Epibulus insidiator* (Pallas, 1770)
- Gomphosus varius* Lacepède, 1801 (Synonym: *Thalassoma stuckiae* Whitley, 1959)
- Halichoeres argus* (Bloch & Schneider, 1801) (Synonym: *Halichoeres leparensis*)
- Halichoeres biocellatus* Schultz in Schultz, Chapman, Lachner & Woods, 1960
- Halichoeres chrysus* Randall, 1981
- Halichoeres chloropterus* (Bloch, 1791)
- Halichoeres hortulanus* (Lacepède, 1801)
- Halichoeres lamarii annularis* (Valenciennes in Cuvier & Valenciennes, 1839) (Reported from New Caledonia by several authors under the name *Halichoeres marginatus*; new record from New Caledonia based on these records and SMNS material from Grande Terre and Lifou/Loyalty Islands)
- Halichoeres margaritaceus* (Valenciennes in Cuvier & Valenciennes, 1839)
- Halichoeres melanurus* (Bleeker, 1851) (Synonym: *Halichoeres hoevenii*; reported by Rivaton et al. [1989: 46] under the name *Halichoeres timorensis*)

*Halichoeres miniatus* (Kuhl & Hasselt in Valenciennes in Cuvier & Valenciennes, 1839)  
*Halichoeres nebulosus* (Valenciennes in Cuvier & Valenciennes, 1839)  
*Halichoeres ornatissimus* (Garrett, 1863) (Reported from New Caledonia by Rivaton *et al.* [1989: 46] under the name *Halichoeres vrolikii*)  
*Halichoeres prosopeion* (Bleeker, 1853)  
*Halichoeres scapularis* (Bennett, 1832) (New record from New Caledonia based on SMNS 21560 from southern Grande Terre)  
*Halichoeres trimaculatus* (Quoy & Gaimard, 1834)  
*Halichoeres zeylonicus* (Bennett, 1833)  
*Hemigymnus fasciatus* (Bloch, 1792)  
*Hemigymnus melapterus* (Bloch, 1791)  
*Hologymnosus annulatus* (Lacepède, 1801)  
*Hologymnosus doliatus* (Lacepède, 1802)  
*Hologymnosus longipes* (Günther, 1862)  
*Iniistius aneitensis* (Günther, 1862)  
*Iniistius celebicus* (Bleeker, 1856) (New record from New Caledonia based on *Xyrichthys* sp. of Kulbicki, Randall & Rivaton [1994: 30] from Chesterfield Islands).  
*Iniistius pavo* (Valenciennes in Cuvier & Valenciennes, 1839)  
*Labrichthys unilineatus* (Guichenot, 1847)  
*Labroides bicolor* Fowler & Bean, 1928  
*Labroides dimidiatus* (Valenciennes in Cuvier & Valenciennes, 1839)  
*Labroides pectoralis* Randall & Springer, 1975  
*Labropsis australis* Randall, 1981  
*Labropsis xanthonota* Randall, 1981  
*Macropharyngodon choati* Randall, 1978  
*Macropharyngodon kuiteri* Randall, 1978  
*Macropharyngodon meleagris* (Valenciennes in Cuvier & Valenciennes, 1839)  
*Macropharyngodon negrosensis* Herre, 1932  
*Novaculichthys taeniourus* (Lacepède, 1802)  
*Oxycheilinus bimaculatus* (Valenciennes in Cuvier & Valenciennes, 1839)  
*Oxycheilinus celebicus* (Bleeker, 1853)  
*Oxycheilinus digrammus* (Lacepède, 1801)  
*Oxycheilinus nigromarginatus* Randall, Westneat & Gomon, 2003  
*Oxycheilinus orientalis* (Günther, 1862)  
*Oxycheilinus unifasciatus* (Streets, 1877) (Reported from New Caledonia by Fourmanoir & Laboute [1976: 117] under the name *Cheilinus rhodocrous*)  
*Pseudocheilinus evanidus* Jordan & Evermann, 1903  
*Pseudocheilinus hexataenia* (Bleeker, 1857)  
*Pseudocheilinus ocellatus* Randall, 1999  
*Pseudocheilinus octotaenia* Jenkins, 1900  
*Pseudocheilinus tetraetaenia* Schultz in Schultz, Chapman, Lachner & Woods, 1960  
*Pseudocoris bleekeri* (Hubrecht, 1876) (Reported from New Caledonia by authors under the name *Coris philippina*)  
*Pseudocoris yamashiroi* (Schmidt, 1931)  
*Pseudodax moluccanus* (Valenciennes in Cuvier & Valenciennes, 1839)  
*Pseudojuloides cerasinus* (Snyder, 1904)  
*Pteragogus* sp. (Undescribed species, which has been reported from New Caledonia by several authors under the name *Pteragogus cryptus*, a species which is restricted to the Red Sea according to Kuiter [2002: 56])  
*Pteragogus enneacanthus* (Bleeker, 1853)  
*Pteragogus flagellifer* (Valenciennes in Cuvier & Valenciennes, 1839) (Reported from New Caledonia by authors under the names *Pteragogus opercularis* and *P. flagellifera*)  
*Stethojulis bandanensis* (Bleeker, 1851) (Reported from New Caledonia by Whitley [1961: 65] under the name *Stethojulis axillaris*)

*Stethojulis notialis* Randall, 2000 (Reported from New Caledonia by various authors under the name *Stethojulis interrupta*)  
*Stethojulis strigiventer* (Bennett, 1832)  
*Suezichthys arquatus* Russell, 1985  
*Suezichthys devisi* (Whitley, 1941) (Reported from New Caledonia by various authors under the name *Suezichthys gracilis*)  
*Terelabrus rubrovittatus* Randall & Fourmanoir, 1998  
*Thalassoma amblycephalum* (Bleeker, 1856) (Reported from New Caledonia by Whitley [1961: 65] under the name *Thalassoma melanochir*)  
*Thalassoma hardwicke* (Bennett, 1830)  
*Thalassoma lunare* (Linnaeus, 1758)  
*Thalassoma lutescens* (Lay & Bennett, 1839) (Reported from New Caledonia by Whitley [1961: 65] under the name *Thalassoma aneitense*)  
*Thalassoma nigrofasciatum* Randall, 2003 (Reported from New Caledonia by several authors under the name *Thalassoma jansenii*)  
*Thalassoma purpureum* (Forsskål in Niebuhr, 1775)  
*Thalassoma quinquevittatum* (Lay & Bennett, 1839) (Reported from Lifou/Loyalty Islands by Günther [1909: 294-295] under the name *Julis güntheri*)  
*Thalassoma trilobatum* (Lacepède, 1801)  
*Wetmorella albofasciata* Schultz & Marshall, 1954  
*Wetmorella nigropinnata* Seale, 1901 (Reported from New Caledonia by Fourmanoir & Laboute [1976: 128] under the name *Wetmorella ocellata*)  
*Xiphocelius typus* Bleeker, 1856 (Reported from New Caledonia by Parenti & Randall [2000: 48] under the name *Xiphocelius quadrimaculatus*)

## SCARIDAE

Records by Plessis & Fourmanoir (1966: 136) of *Scarus tricolor* Bleeker, 1847 (under the name *Scarus mus*) need verification.  
*Bolbometopon muricatum* (Valenciennes in Cuvier & Valenciennes, 1839)  
*Calotomus carolinus* (Valenciennes in Cuvier & Valenciennes, 1840)  
*Calotomus spinidens* (Quoy & Gaimard, 1824)  
*Cetoscarus ocellatus* (Valenciennes in Cuvier & Valenciennes, 1840) (Reported from New Caledonia by various authors under the name *Cetoscarus bicolor*; **new record** from New Caledonia, also based on AMS IB.3830)  
*Chlorurus bleekeri* (Beaufort, 1940)  
*Chlorurus frontalis* (Valenciennes in Cuvier & Valenciennes, 1839)  
*Chlorurus microrhinos* (Bleeker, 1854) (Reported from New Caledonia by several authors under the name *Scarus gibbus*, and by Rivaton & Bourret [1999: 16] confused with *Scarus rhoduropterus*)  
*Chlorurus sordidus* (Forsskål in Niebuhr, 1775)  
*Hipposcarus longiceps* (Valenciennes in Cuvier & Valenciennes, 1840)  
*Leptoscarus vaigiensis* (Quoy & Gaimard, 1824)  
*Scarus altipinnis* (Steindachner, 1879)  
*Scarus chameleon* Choat & Randall, 1986  
*Scarus dimidiatus* Bleeker, 1859  
*Scarus flavipectoralis* Schultz, 1958  
*Scarus forsteni* (Bleeker, 1861)  
*Scarus frenatus* Lacepède, 1802  
*Scarus ghobban* (Forsskål in Niebuhr, 1775) (Reported from New Caledonia by Laboute & Grandperrin [2000: 383] under the name *Scarus* sp. 3)  
*Scarus globiceps* Valenciennes in Cuvier & Valenciennes, 1839  
*Scarus longipinnis* Randall & Choat, 1980 (Reported from New Caledonia by Laboute & Grandperrin [2000: 383] under the name *Scarus* sp. 1)  
*Scarus niger* (Forsskål in Niebuhr, 1775)  
*Scarus oviceps* Valenciennes in Cuvier & Valenciennes, 1839  
*Scarus psittacus* Forsskål in Niebuhr, 1775  
*Scarus quoyi* Valenciennes in Cuvier & Valenciennes, 1840

*Scarus rivulatus* Valenciennes in Cuvier & Valenciennes, 1840 (Reported from New Caledonia by several authors under the name *Scarus fasciatus*)

*Scarus rubroviolaceus* Bleeker, 1847

*Scarus schlegeli* (Bleeker, 1861) (Reported from New Caledonia by various authors under the name *Scarus venosus*)

*Scarus spinus* (Kner, 1868)

#### PINGUIPEDIDAE

*Parapercis australis* Randall, 2003 (Reported by various authors under the name *Parapercis cylindrica*)

*Parapercis binivirgata* (Waite, 1904)

*Parapercis clathrata* Ogilby, 1910

*Parapercis colemani* Randall & Francis, 1993 (New record from New Caledonia based on SMNS 23850 from Lifou, Loyalty Islands)

*Parapercis hexophtalma* (Cuvier in Cuvier & Valenciennes, 1829) (Reported from New Caledonia by various authors under the name *Parapercis polyophthalma*)

*Parapercis lineopunctata* Randall, 2003

*Parapercis millepunctata* (Günther, 1860)

*Parapercis multiplicata* Randall, 1984

*Parapercis nebulosa* (Quoy & Gaimard, 1825) (New record from New Caledonia, based on SMNS 23673 from Lifou, Loyalty Islands)

*Parapercis schauinslandii* (Steindachner, 1900)

*Parapercis snyderi* Jordan & Starks, 1905 (Reported from New Caledonia by Laboute & Grandperrin [2000: 391] under the name *Parapercis* sp.)

*Parapercis xanthozona* (Bleeker, 1849)

#### TRICHONOTIDAE

*Trichonotus elegans* Shimada & Yoshino, 1984

*Trichonotus filamentosus* (Steindachner, 1867)

*Trichonotus setiger* (Bloch & Schneider, 1801)

#### CREDIIDAE

*Chalixodutes tauensis* Schultz, 1943 (Synonym: *Chalixodutes chameleontoculis*)

*Limnichthys nitidus* Smith, 1958 (Synonym: *Limnichthys donaldsoni* Schultz in Schultz et al., 1960)

#### AMMODYTIDAE

*Ammodytoides vagus* (McCulloch & Waite, 1916) (Reported from New Caledonia by Laboute & Grandperrin [2000: 387] under the name *Ammodytes* sp.)

*Bleekeria mitsukurii* Jordan & Evermann, 1902

#### URANOSCOPIDAE

A record of *Ichthyscopus lebeck* Bloch & Schneider, 1801 by two authors needs verification; New Caledonian populations may be referable to *Ichthyscopus sannio* Whitley, 1936.

*Uranoscopus oligolepis* Bleeker, 1858

*Uranoscopus sulphureus* Valenciennes in Cuvier & Valenciennes, 1831

#### TRIPTYGYIIDAE

*Ceratobregma helenae* Holleman, 1987 (Synonym: *Ceratobregma striata* Fricke, 1994)

*Enneapterygius elegans* (Peters, 1877)

*Enneapterygius flavoccipitis* Shen & Wu, 1994 (Synonym: *Enneapterygius bichrous* Fricke, 1994)

*Enneapterygius hemimelas* (Kner & Steindachner, 1867) (Reported from Chesterfield Islands by Kulbicki et al. [1994: 32] under the name *Enneapterygius semilarvatus*, a nomen nudum)

*Enneapterygius howensis* Fricke, 1997

*Enneapterygius nanus* (Schultz in Schultz, Chapman, Lachner & Woods, 1960)

*Enneapterygius niger* Fricke, 1994

*Enneapterygius paucifasciatus* Fricke, 1994

*Enneapterygius philippinus* (Peters, 1869) (Synonym: *Enneapterygius minutus*; reported from New Caledonia by Whitley [1961: 65] under the name *Vauclusella rufopileum*)

*Enneapterygius rhabdotus* Fricke, 1994

*Enneapterygius rhothion* Fricke, 1997

*Enneapterygius rubicauda* Shen & Wu, 1994

*Enneapterygius rufopileus* (Waite, 1904)  
*Enneapterygius similis* Fricke, 1997  
*Enneapterygius triserialis* Fricke, 1994  
*Enneapterygius trisignatus* Fricke, 2001  
*Enneapterygius tutuilae* Jordan & Seale, 1906  
*Enneapterygius williamsi* Fricke, 1997  
*Helcogramma* sp. (To be described by J.T. Williams) (Reported by several authors from New Caledonia under the name *Helcogramma ellioti*)  
*Helcogramma hudsoni* (Jordan & Seale, 1906)  
*Helcogramma novaecaledoniae* Fricke, 1994  
*Helcogramma trigloides* (Bleeker, 1858)  
*Norfolkia brachylepis* (Schultz in Schultz, Chapman, Lachner & Woods, 1960)  
*Norfolkia squamiceps* (McCulloch & Waite, 1916)  
*Norfolkia thomasi* Whitley, 1964  
*Springerichthys kulwickii* (Fricke & Randall in Fricke, 1994)  
*Ucla xenogrammus* Holleman, 1993

#### BLENNIIDAE

Records of *Andamia reyi* (Sauvage, 1880) by Rivaton *et al.* (1989: 27), *Ecsenius aequalis* Springer, 1988 and *Salarias sinuosus* Snyder, 1908 by Rivaton *et al.* (1989: 28, 29) need verification. Records of *Cirripectes variolosus* (Valenciennes in Cuvier & Valenciennes, 1836) (under the name *C. sebae*) by several authors also need verification.  
*Alticus* sp. (To be described by J.T. Williams)  
*Alticus sertatus* (Garman, 1903)  
*Andamia amphibius* (Walbaum, 1792) (**New record** from New Caledonia, based on SMNS 23579 from Lifou, Loyalty Islands)  
*Aspidontus dussumieri* (Valenciennes in Cuvier & Valenciennes, 1836)  
*Aspidontus taeniatus* Quoy & Gaimard, 1836  
*Atrosalarias fuscus holomelas* (Günther, 1872)  
*Blenniella chrysospilos* (Bleeker, 1857)  
*Blenniella paula* (Bryan & Herre, 1903) (**New record** from New Caledonia based on SMNS 22752 from northeastern Grande Terre)  
*Blenniella periophthalmus* (Valenciennes in Cuvier & Valenciennes, 1836)  
*Cirripectes alboapicalis* (Ogilby, 1899) (Reported from New Caledonia by Laboute & Grandperrin [2000: 397] under the name *Cirripectes* sp.; **new record** from New Caledonia, also based on SMNS 23810 from Lifou, Loyalty Islands)  
*Cirripectes castaneus* (Valenciennes in Cuvier & Valenciennes, 1836)  
*Cirripectes chelomatus* Williams & Maugé, 1983  
*Cirripectes filamentosus* (Alleyne & Macleay, 1877) (**New record** from New Caledonia, based on SMNS 22769 from northern Grande Terre)  
*Cirripectes fuscoguttatus* Strasburg & Schultz, 1953 (**New record** from New Caledonia, based on SMNS material from northern Grande Terre and Lifou/Loyalty Islands)  
*Cirripectes polyzonus* Bleeker, 1868  
*Cirripectes stigmaticus* Strasburg & Schultz, 1953  
*Crossosalarias macrospilus* Smith-Vaniz & Springer, 1971  
*Ecsenius bicolor* (Day, 1888)  
*Ecsenius fourmanoiri* Springer, 1972  
*Ecsenius isos* McKinney & Springer, 1976 (Reported from New Caledonia by Laboute & Grandperrin [2000: 399] under the name *Ecsenius* sp.)  
*Ecsenius midas* Starck, 1969  
*Ecsenius nalolo* Smith, 1959  
*Ecsenius stictus* Springer, 1988 (Reported by several authors under the name *Ecsenius yaeyamaensis*)  
*Ecsenius tessera* Springer, 1988 (Reported from New Caledonia by Laboute & Grandperrin [2000: 399] under the name *Ecsenius oculus*)  
*Enchelyurus ater* (Günther, 1877)  
*Enchelyurus kraussii* (Klunzinger, 1871)

*Entomacrodus caudofasciatus* (Regan, 1909)  
*Entomacrodus decussatus* (Bleeker, 1858)  
*Entomacrodus sealei* Bryan & Herre, 1903  
*Entomacrodus striatus* (Quoy & Gaimard, 1836)  
*Entomacrodus thalassinus* (Jordan & Seale, 1906) (Reported from New Caledonia by Laboute & Grandperrin [2000: 400] under the name 'Blenniidae indéterminé'; **new record** from New Caledonia, also based on SMNS 19796 from southern Grande Terre)  
*Exallias brevis* (Kner, 1868)  
*Istiblennius dussumieri* (Valenciennes in Cuvier & Valenciennes, 1836) (Reported by Günther [1877: 208] under the name *Salarias meleagris*, by Whitley [1927: 304] as *Salarias geminatus*, and by Plessis & Fourmanoir [1966: 140] under the name *Halmablenius striatomaculatus*) Plessis & Fourmanoir, 1966: 140 (Nouvelle-Calédonie/Grande Terre, New Caledonia).  
*Istiblennius edentulus* (Forster in Bloch & Schneider, 1801)  
*Istiblennius lineatus* (Valenciennes in Cuvier & Valenciennes, 1836)  
*Meiacanthus anema* (Bleeker, 1852)  
*Meiacanthus atrodorsalis* (Günther, 1877)  
*Meiacanthus ditrema* Smith-Vaniz, 1976  
*Meiacanthus grammistes* (Valenciennes in Cuvier & Valenciennes, 1836)  
*Meiacanthus phaeus* Smith-Vaniz, 1976  
*Nannosalarias nativitatus* (Regan, 1909) (**New record** from New Caledonia based on SMNS material from southern Grande Terre)  
*Omobranchus germaini* (Sauvage, 1883) (Synonym: *Graviceps alexanderi* Whitley, 1945)  
*Omobranchus obliquus* (Garman, 1903) (Reported from New Caledonia by Rivaton *et al.* [1989: 28] under the name *Omobranchus banditus*)  
*Omox biporus* Springer, 1972  
*Parablennius tasmanianus* (Richardson, 1849) (Synonym: *Parablennius tasmanianus caledoniensis* Bath, 1989; taxonomic decision of H. Bath, personal communication, June 2006)  
*Petroscirtes breviceps* (Valenciennes in Cuvier & Valenciennes, 1836)  
*Petroscirtes lupus* (De Vis, 1886) (Synonym: *Dasson icelii*)  
*Petroscirtes mitratus* Rüppell, 1830  
*Petroscirtes variabilis* Cantor, 1850  
*Petroscirtes xestus* Jordan & Seale, 1906  
*Plagiotremus laudandus* (Whitley, 1961)  
*Plagiotremus rhinorhynchos* (Bleeker, 1852)  
*Plagiotremus tapeinosoma* (Bleeker, 1857) (Reported from New Caledonia by Laboute & Grandperrin [2000: 68] under the name *Omox* sp.)  
*Praealticus bilineatus* (Peters, 1868)  
*Rhabdoblennius snowi* (Fowler, 1928) (Reported from New Caledonia by Kulbicki *et al.* [1994: 33] under the name *Rhabdoblennius ellipes*)  
*Salarias alboguttatus* Kner, 1867  
*Salarias fasciatus* (Bloch, 1786) (Reported from New Caledonia by Fourmanoir & Laboute [1976: 137] under the name *Petroscirtes mitratus*)  
*Salarias guttatus* Valenciennes in Cuvier & Valenciennes, 1836 (**New record** from New Caledonia based on SMNS 19777 from southeastern Grande Terre)  
*Stanulus seychellensis* Smith, 1959 (**New record** from New Caledonia based on SMNS 23616 from Lifou, Loyalty Islands)  
*Stanulus talboti* Springer, 1968 (**New record** from New Caledonia based on SMNS 23617 from Lifou, Loyalty Islands)  
*Xiphasia setifer* Swainson, 1839

## CLINIDAE

*Springeratus* sp. (Undescribed species) (Reported from New Caledonia under the names *Petraites roseus*, *Petraites sellularius* and *Petraites nasutus*)

## GOBIESOCIDAE

*Conidens samoensis* (Steindachner, 1906)  
*Diademichthys lineatus* (Sauvage, 1883)

*Discotrema crinophila* Briggs, 1976

*Lepadichthys* sp. (Undescribed species based on SMNS material from southern Grande Terre)

*Lepadichthys frenatus* Waite, 1904

*Lepadichthys minor* Briggs, 1955

*Pherallodus indicus* (Weber, 1913) (Reported by Kulbicki & Williams [1997: 12] under the name

*Pherallodus* sp.; **new record** from New Caledonia, also based on SMNS material from Lifou, Loyalty Islands)

#### CALLIONYMIDAE

*Callionymus* sp. (Chesterfield Islands; to be described by R. Fricke; recorded by Kulbicki *et al.* [1994: 33] under the name *Calliurichthys japonicus*, and by Fricke [2000: 20-24, part] under the name *Callionymus scaber*)

*Callionymus brevianalis* Fricke, 1983

*Callionymus corallinus* Gilbert, 1905

*Callionymus enneactis* Bleeker, 1879

*Callionymus keeleyi* Fowler, 1941 (Reported from New Caledonia by several authors under the name *Repomucenus virgus*)

*Callionymus pleurostictus* Fricke, 1982

*Callionymus rivatoni* Fricke, 1993 (Reported from New Caledonia by several authors under the name *Repomucenus huguenini*; reported by Laboute & Grandperrin [2000: 400] as *Diplogrammus goramensis*)

*Callionymus simplicicornis* Valenciennes in Cuvier & Valenciennes, 1837

*Callionymus tethys* Fricke, 1993

*Diplogrammus goramensis* (Bleeker, 1858)

*Synchiropus circularis* Fricke, 1984

*Synchiropus morrisoni* Schultz in Schultz, Chapman, Lachner & Woods, 1960 (Reported from New Caledonia by Laboute & Grandperrin [2000: 401, left fig.] under the name *Synchiropus ocellatus* [part])

*Synchiropus ocellatus* (Pallas, 1770) (Reported by Laboute & Grandperrin [2000: 401] under the name *Synchiropus* sp.)

*Synchiropus rameus* (McCulloch, 1926)

*Synchiropus rubrovinctus* (Gilbert, 1905) (**New record** from New Caledonia based on SMNS material from Lifou, Loyalty Islands)

*Synchiropus sechellensis* Regan, 1908

*Synchiropus splendidus* (Herré, 1927)

*Synchiropus springeri* Fricke, 1983 (Reported from New Caledoniy by Rivaton *et al.* [1989: 31] under the name *Synchiropus postulus*, and by Kulbicki & Williams [1997: 21] under the name *Minysynchiropus laddi*)

#### ELEOTRIDAE

*Butis amboinensis* (Bleeker, 1853)

*Calumia godeffroyi* (Günther, 1877)

*Eleotris acanthopoma* Bleeker, 1853

*Eleotris fusca* (Bloch & Schneider, 1801) (Reported from New Caledonia by Whitley [1961: 64] under the name *Culius melanosoma*)

*Eleotris melanosoma* Bleeker, 1852

#### XENISTHMIDAE

*Xenisthmus eirospilus* Gill & Hoese, 2004 (Reported from Chesterfield Islands by Kulbicki *et al.* [1994: 35] under the name *Xenisthmus* sp.; **new record** from New Caledonia)

*Xenisthmus polyzonatus* (Klunzinger, 1871)

#### KRAEMERIIDAE

*Kraemeria samoensis* Steindachner, 1906

#### GOBIIDAE

Records of *Drombus triangularis* (Weber, 1909), under the name *Acentrogobius bontii*, by several authors including Fourmanoir (1981: 27), *Ambyeleotris japonica* Takagi, 1957 by Fourmanoir & Laboute (1976: 140), *Bathygobius petrophilus* (Bleeker, 1853) by Plessis & Fourmanoir [1966: 139], *Parapocryptes serperaster* (Richardson, 1846) (under the name *Apocryptes* s.) by Borodin (1932: 96), and *Amblygobius hectori* (Smith,

1957), *Bryaninops ridens* Smith, 1959 and *Valenciennea sexguttata* (Valenciennes in Cuvier & Valenciennes, 1837) by Rivaton *et al.* [1989: 40, 42], need verification.

*Amblyeleotris bellicauda* Randall, 2004

*Amblyeleotris biguttata* Randall, 2004

*Amblyeleotris diagonalis* Polunin & Lubbock, 1979 (Reported by Laboute & Grandperrin [2000: 405] under the name *Amblyeleotris* sp. 1)

*Amblyeleotris fasciata* (Herre, 1953) (Synonym: *Amblyeleotris wheeleri*)

*Amblyeleotris fontanesii* (Bleeker, 1852)

*Amblyeleotris guttata* (Fowler, 1938)

*Amblyeleotris novacaledoniae* Goren, 1981

*Amblyeleotris ogasawarensis* Yanagisawa, 1978 (Reported from New Caledonia by Laboute & Grandperrin [2000: 7, 8] under the names *Amblyeleotris* sp. 7 and *Amblyeleotris* sp. 8)

*Amblyeleotris periophthalma* (Bleeker, 1853) (Synonym: *Amblyeleotris exilis*)

*Amblyeleotris randalli* Hoese & Steene, 1978

*Amblyeleotris rubrimarginata* Mohlmann & Randall, 2002 (Reported from New Caledonia by Laboute & Grandperrin [2000: 406] under the name *Amblyeleotris* sp. 9)

*Amblyeleotris steinitzi* (Klausewitz, 1974)

*Amblyeleotris stenotaeniata* Randall, 2004 (Reported from New Caledonia by Laboute & Grandperrin [2000: 406] under the names *Amblyeleotris* sp. 4 and *Amblyeleotris* sp. 5)

*Amblygobius bynoensis* (Richardson, 1844)

*Amblygobius decussatus* (Bleeker, 1855)

*Amblygobius linki* (Herre, 1927)

*Amblygobius nocturnus* (Herre, 1945)

*Amblygobius phalaena* (Valenciennes in Cuvier & Valenciennes, 1837) (Reported from New Caledonia by several authors under the name *Amblygobius albimaculatus*)

*Amblygobius sphynx* (Valenciennes in Cuvier & Valenciennes, 1837)

*Asterropteryx ensifera* (Bleeker, 1874)

*Asterropteryx semipunctata* Rüppell, 1830

*Asterropteryx spinosa* (Goren, 1981)

*Barbuligobius* sp. (Record based on SMNS 23775 from Lifou, Loyalty Islands)

*Bathygobius coalitus* (Bennett, 1832) (Reported from Grande Terre by several authors under the name *Bathygobius albopunctatus*, which is a synonym; new record from New Caledonia, also based on MNHN material from Grande Terre)

*Bathygobius cocosensis* (Bleeker, 1854)

*Bathygobius cotticeps* (Steindachner, 1880) (New record from New Caledonia based on SMNS material from northeastern, southwestern and southern Grande Terre and Lifou/Loyalty Islands)

*Bathygobius cyclopterus* (Valenciennes in Cuvier & Valenciennes, 1837)

*Bathygobius fuscus* (Rüppell, 1830) (Synonyms: *Gobius caledonicus* Sauvage, 1880; *Gobius filamentosus* Sauvage, 1883, both originally described from New Caledonia)

*Bryaninops loki* Larson, 1985

*Bryaninops natans* Larson, 1985

*Bryaninops yongei* (Davis & Cohen, 1968)

*Cabillus tongarevae* (Fowler, 1927)

*Callogobius hasseltii* (Bleeker, 1851)

*Callogobius maculipinnis* (Fowler, 1918) (Synonyms: *Callogobius irrasus*; *Intosagobius kuderi* Herre, 1943)

*Callogobius sclateri* (Steindachner, 1880)

*Cryptocentrus fasciatus* (Playfair & Günther, 1867)

*Cryptocentrus leptcephalus* Bleeker, 1876 (Reported from New Caledonia by several authors including Laboute & Grandperrin [2000: 409] under the name *Cryptocentrus lutheri*; new record from New Caledonia, also based on SMNS 21552 from southern Grande Terre)

*Cryptocentrus strigilliceps* (Jordan & Seale, 1906)

*Ctenogobiops aurocingulus* (Herre, 1935)

*Ctenogobiops feroculus* Lubbock & Polunin, 1977

*Ctenogobiops pomastictus* Lubbock & Polunin, 1977

- Ctenotrypauchen microcephalus* (Bleeker, 1860)
- Eviota afelei* Jordan & Seale, 1906
- Eviota albolineata* Jewett & Lachner, 1983
- Eviota cometa* Jewett & Lachner, 1983
- Eviota distigma* Jordan & Seale, 1906 (Reported by Whitley [1961: 64] under the name *Eviota abax*)
- Eviota fasciola* Karnella & Lachner, 1981
- Eviota herrei* Jordan & Seale, 1906 (New record from New Caledonia based on SMNS 18327  
from southeastern Grande Terre)
- Eviota hoessei* Gill & Jewett, 2004
- Eviota latifasciata* Jewett & Lachner, 1983
- Eviota melasma* Lachner & Karnella, 1980
- Eviota monostigma* Fourmanoir, 1971
- Eviota nebulosa* Smith, 1958
- Eviota nigriventris* Giltay, 1933
- Eviota pellucida* Larson, 1976
- Eviota prasina* (Klunzinger, 1871)
- Eviota prasites* Jordan & Seale, 1906
- Eviota pseudostigma* Lachner & Karnella, 1980
- Eviota punctulata* Jewett & Lachner, 1983 (New record from New Caledonia based on SMNS  
material from southern Grande Terre and Maré/Loyalty Islands)
- Eviota queenslandica* Whitley, 1932 (New record from New Caledonia based on SMNS material  
from northern Grande Terre and Lifou/Loyalty Islands)
- Eviota sebreei* Jordan & Seale, 1906
- Eviota smaragdus* Jordan & Seale, 1906
- Eviota sparsa* Jewett & Lachner, 1983
- Eviota zebrina* Lachner & Karnella, 1978
- Eviota zonura* Jordan & Seale, 1906
- Exyrias belissimus* (Smith, 1959) (Reported from New Caledonia by Laboute & Grandperrin  
[2000: 417] under the name 'Gobiidae indéterminé 1')
- Exyrias puntang* (Bleeker, 1851) (Synonym: *Gobius canalaee* Sauvage, 1881)
- Favonigobius reichei* (Bleeker, 1853)
- Fusigobius duospilus* Hoesse & Reader, 1985
- Fusigobius gracilis* (Randall, 2001)
- Fusigobius humeralis* (Randall, 2001)
- Fusigobius longispinus* Goren, 1978 (New record from New Caledonia based on SMNS 22888  
from Ile des Pins)
- Fusigobius maximus* (Randall, 2001)
- Fusigobius neophytus* (Günther, 1877)
- Fusigobius pallidus* (Randall, 2001)
- Fusigobius signipinnis* Hoesse & Obika, 1988
- Glossogobius* sp. (An undescribed species which was reported by several authors under the name  
*Glossogobius celebius*)
- Glossogobius biocellatus* (Valenciennes in Cuvier & Valenciennes, 1837)
- Glossogobius giuris* (Hamilton Buchanan, 1822)
- Gnatholepis anjerensis* (Bleeker, 1851)
- Gnatholepis cauerensis cauerensis* (Bleeker, 1853) (Synonym: *Gnatholepis scapulostigma*)
- Gobiodon acicularis* Harold & Winterbottom, 1995 (New record from New Caledonia based on  
SMNS 23914 from Lifou, Loyalty Islands)
- Gobiodon axillaris* DeVis, 1884 (New record from New Caledonia based on SMNS material from  
Grande Terre and Lifou, Loyalty Islands)
- Gobiodon brochus* Harold & Winterbottom, 1999
- Gobiodon citrinus* (Rüppell, 1838)
- Gobiodon multilineatus* Wu, 1979
- Gobiodon okinawae* Sawada, Arai & Abe, 1972
- Gobiodon quinquestrigatus* (Valenciennes in Cuvier & Valenciennes, 1837)

- Gobiodon rivulatus* (Rüppell, 1830)
- Gobiopsis malekulae* (Herre, 1935) (**New record** from New Caledonia based on SMNS 23692 from Lifou, Loyalty Islands)
- Hetereleotris* sp. (Undescribed species; record based on SMNS material from Lifou, Loyalty Islands)
- Istigobius decoratus* (Herre, 1927) (Reported from New Caledonia by Laboute & Grandperrin [2000: 417] under the name 'Gobiidae indéterminé 6')
- Istigobius nigroocellatus* (Günther, 1873) (Reported from New Caledonia by Laboute & Grandperrin [2000: 412] under the name *Istigobius* sp.)
- Istigobius ornatus* (Rüppell, 1830)
- Istigobius rigilius* (Herre, 1953)
- Istigobius spence* (Smith, 1947) (Synonym: *Acentrogobius aestuarius* Smith, 1959)
- Koumansetta rainfordi* (Whitley, 1940)
- Lubricogobius exiguus* Tanaka, 1915
- Lubricogobius ornatus* Fourmanoir, 1966
- Macrodontogobius wilburi* Herre, 1936
- Mahidolia mystacina* (Valenciennes in Cuvier & Valenciennes, 1837) (Synonym: *Mahidolia duque* Smith, 1947)
- Mugilogobius mertoni* (Weber, 1911)
- Mugilogobius notospilus* (Günther, 1877) (Synonym: *Mugilogobius duospilus*)
- Oplopomus caninoides* (Bleeker, 1852)
- Oplopomus oplopomus* (Valenciennes in Cuvier & Valenciennes, 1837)
- Oxyurichthys papuensis* (Valenciennes in Cuvier & Valenciennes, 1837) (Reported from New Caledonia by Laboute & Grandperrin [2000: 413] under the name *Oxyurichthys* sp.)
- Oxyurichthys tentacularis* (Valenciennes in Cuvier & Valenciennes, 1837)
- Paragobiodon echinocephalus* (Rüppell, 1828)
- Paragobiodon lacunicolus* (Kendall & Goldsborough, 1911)
- Paragobiodon melanosomus* (Bleeker, 1852)
- Paragobiodon modestus* (Regan, 1908) (**New record** from New Caledonia based on SMNS material from northern and southern Grande Terre, and Maré and Lifou, Loyalty Islands)
- Paragobiodon xanthosomus* (Bleeker, 1852)
- Periophthalmus argenteolineatus* Valenciennes in Cuvier & Valenciennes, 1837 (Synonym: *Periophthalmus vulgaris* Eggert, 1935)
- Pleurosicya bilobata* (Koumans, 1941) (Synonym: *Pleurosicya taisnei* Plessis & Fourmanoir, 1966)
- Pleurosicya coerulea* Larson, 1990
- Pleurosicya micheli* Fourmanoir, 1971
- Pleurosicya mossambica* Smith, 1959
- Priolepis cincta* (Regan, 1908) (Synonym: *Priolepis naraharae*; reported from New Caledonia by some authors including Plessis & Fourmanoir [1966: 139] under the name *Quisquilius eugenius*)
- Priolepis compita* Winterbottom, 1985
- Priolepis fallacincta* Winterbottom & Burridge, 1992
- Priolepis inhaca* (Smith, 1949) (**New record** from New Caledonia based on SMNS 21677 from Lifou, Loyalty Islands)
- Priolepis kappa* Winterbottom & Burridge, 1992
- Priolepis semidoliata* (Valenciennes in Cuvier & Valenciennes, 1837)
- Priolepis squamogena* Winterbottom & Burridge, 1989 (**New record** from New Caledonia based on SMNS 22934 from Lifou, Loyalty Islands)
- Stonogobiops yasha* Yoshino & Shimada, 2001 (Reported by Laboute & Grandperrin [2000: 414] under the name *Stonogobiops* sp.)
- Sueviota lachneri* Winterbottom & Hoese, 1988
- Tomiyamichthys* sp. (Reported from New Caledonia by Laboute & Grandperrin [2000: 417] under the name 'Gobiidae indéterminé 5')
- Trimma* sp. (Undescribed species; previously reported by several authors under the name *Trimma naudei*)
- Trimma benjamini* Winterbottom, 1996
- Trimma caesiura* Jordan & Seale, 1906
- Trimma emeryi* Winterbottom, 1985 (**New record** from New Caledonia based on SMNS material

from Ile des Pins and Chesterfield Islands)  
*Trimma milta* Winterbottom, 2002 (**New record** from New Caledonia based on SMNS material  
from Chesterfield Islands and Lifou, Loyalty Islands)  
*Trimma okinawae* (Aoyagi, 1949)  
*Trimma striata* (Herre, 1945) (Synonym: *Zonogobius capostriatus* Goren, 1981)  
*Trimma taylori* Lobel, 1979  
*Trimma tevegae* Cohen & Davis, 1969  
*Trimma unisquamis* (Gosline, 1959)  
*Trimmatom eviotops* (Schultz, 1943) (Synonym: *Eviota corneliae* Fricke, 1998)  
*Trimmatom nanus* Winterbottom & Emery, 1981  
*Valenciennea decora* Hoese & Larson, 1994  
*Valenciennea helsdingenii* (Bleeker, 1858)  
*Valenciennea limicola* Hoese & Larson, 1994  
*Valenciennea longipinnis* (Lay & Bennett, 1839)  
*Valenciennea parva* Hoese & Larson, 1994  
*Valenciennea puellaris* (Tomiyama, 1955)  
*Valenciennea randalli* Hoese & Larson, 1994  
*Valenciennea strigata* (Broussonet, 1782)  
*Valenciennea wardii* (Playfair, 1866)  
*Vanderhorstia* sp. 1 (Undescribed species; Blackblotched shrimp goby; reported by several authors  
under the name *Vanderhorstia ornatissima*)  
*Vanderhorstia* sp. 2 (Undescribed species; Halfstreaked shrimp goby; reported from New  
Caledonia by Laboute & Grandperrin [2000: 418] under the name 'Gobiidae indéterminé 10')  
*Vanderhorstia ambanoro* (Fourmanoir, 1957)  
*Yongeichthys criniger* (Valenciennes in Cuvier & Valenciennes, 1837) (Reported from New  
Caledonia by Laboute & Grandperrin [2000: 418] under the name 'Gobiidae indéterminé 12')  
*Yongeichthys nebulosus* (Forsskål in Niebuhr, 1775)  
*Yongeichthys pavidus* (Smith, 1959)

#### MICRODESMIDAE

*Gunnellichthys curiosus* Dawson, 1968  
*Gunnellichthys monostigma* Smith, 1958  
*Gunnellichthys pleurotaenia* Bleeker, 1858 (**New record** from New Caledonia based on SMNS  
21546 from southern Grande Terre)  
*Gunnellichthys viridescens* Dawson, 1968

#### PTERELEOTRIDAE

*Nemateleotris decora* Randall & Allen, 1973  
*Nemateleotris helfrichi* Randall & Allen, 1973  
*Nemateleotris magnifica* Fowler, 1938  
*Parioglossus dotui* Tomiyama, 1958  
*Parioglossus neocaldonicus* Dingerkus & Sérét, 1992  
*Parioglossus philippinus* (Here, 1945) (**New record** from New Caledonia based on SMNS material  
from southern Grande Terre and Lifou/Loyalty Islands)  
*Parioglossus nudus* Rennis & Hoese, 1985 (**New record** from New Caledonia based on SMNS  
21674 from Lifou/Loyalty Islands)  
*Parioglossus rainfordi* McCulloch, 1921  
*Ptereoleotris evides* (Jordan & Hubbs, 1925)  
*Ptereoleotris hanae* (Jordan & Snyder, 1901)  
*Ptereoleotris heteroptera* (Bleeker, 1855)  
*Ptereoleotris microlepis* (Bleeker, 1856)  
*Ptereoleotris monoptera* Randall & Hoese, 1985  
*Ptereoleotris zebra* Fowler, 1938

#### EPHIPPIDAE

*Platax orbicularis* (Forsskål in Niebuhr, 1775)  
*Platax pinnatus* (Linnaeus, 1758)  
*Platax teira* (Forsskål in Niebuhr, 1775)

## SCATOPHAGIDAE

- Scatophagus argus* (Linné, 1766)  
*Scatophagus tetricanthus* (Lacepède, 1801)  
*Selenotoca multifasciata* (Richardson, 1846)

## SIGANIDAE

- Siganus argenteus* (Quoy & Gaimard, 1825)  
*Siganus corallinus* (Valenciennes in Cuvier & Valenciennes, 1835)  
*Siganus doliatus* Guérin-Méneville, 1829-1838  
*Siganus fuscescens* (Houttuyn, 1782) (Synonym: *Siganus nebulosus*; reported from New Caledonia by several authors under the names *Siganus oramin* or *Siganus canaliculatus*, and by Laboute & Grandperrin [2000: 443] as *Siganus* sp.)  
*Siganus lineatus* (Valenciennes in Cuvier & Valenciennes, 1835)  
*Siganus puillus* (Schlegel, 1852)  
*Siganus punctatus* (Forster in Bloch & Schneider, 1801)  
*Siganus spinus* (Linnaeus, 1758)  
*Siganus uspi* Gawel & Woodland, 1974  
*Siganus vermiculatus* (Valenciennes in Cuvier & Valenciennes, 1835)  
*Siganus vulpinus* (Schlegel & Müller, 1844)  
*Siganus woodlandi* Randall & Kulbicki, 2005 (This species has been misidentified by authors as *Siganus argenteus* [non Quoy & Gaimard, 1825])

## ZANCLIDAE

- Zanclus cornutus* (Linnaeus, 1758)

## ACANTHURIDAE

- Acanthurus achilles* Shaw, 1803  
*Acanthurus albopectoralis* Allen & Ayling, 1987  
*Acanthurus blochii* Valenciennes in Cuvier & Valenciennes, 1835  
*Acanthurus dussumieri* Valenciennes in Cuvier & Valenciennes, 1835  
*Acanthurus grammoptilus* Richardson 1843 (New record from New Caledonia based on SMNS 22986 from southern Grande Terre)  
*Acanthurus guttatus* Bloch & Schneider, 1801  
*Acanthurus leucopareius* (Jenkins, 1903)  
*Acanthurus lineatus* (Linnaeus, 1758)  
*Acanthurus mata* (Cuvier in Cuvier & Valenciennes, 1829) (Synonym: *Acanthurus bleekeri*)  
*Acanthurus nigricans* (Linnaeus, 1758) (Synonym: *Acanthurus aliala* Lesson, 1831; reported from New Caledonia by several authors under the name *Acanthurus gahm*).  
*Acanthurus nigrocauda* Duncker & Mohr, 1929  
*Acanthurus nigrofucus* (Forsskål in Niebuhr, 1775)  
*Acanthurus nigroris* (Valenciennes in Cuvier & Valenciennes, 1835) (New record from New Caledonia based on SMNS 19367 from the barrier reef off southern Grande Terre)  
*Acanthurus nubilus* (Fowler & Bean, 1929)  
*Acanthurus olivaceus* Forster in Bloch & Schneider, 1801  
*Acanthurus pyroferus* Kittlitz, 1834 (Synonym: *Acanthurus celebicus* Bleeker, 1852)  
*Acanthurus thompsoni* (Fowler, 1923)  
*Acanthurus triostegus* *triostegus* (Linnaeus, 1758)  
*Acanthurus xanthopterus* Valenciennes in Cuvier & Valenciennes, 1835 (Synonym: *Acanthurus fuliginosus* Lesson, 1831)  
*Ctenochaetus binotatus* Randall, 1955  
*Ctenochaetus cyanocheilus* Randall & Clements, 2001 (Reported by various authors under the name *Ctenochaetus strigosus*)  
*Ctenochaetus striatus* (Quoy & Gaimard, 1825)  
*Naso annulatus* (Quoy & Gaimard, 1824)  
*Naso brachycentron* (Valenciennes in Cuvier & Valenciennes, 1835)  
*Naso brevirostris* (Cuvier, 1829) rnfish  
*Naso caesius* Randall & Bell, 1992  
*Naso fageni* Morrow, 1954 (doubtful record from New Caledonia by Randall [2002: 81])

*Naso hexacanthus* (Bleeker, 1855) (Synonyms: *Naso vomer*; *Naso thorpei*)  
*Naso lituratus* (Forster in Bloch & Schneider, 1801)  
*Naso lopezi* Herre, 1927  
*Naso maculatus* Randall & Struhsaker, 1981  
*Naso tonganus* (Valenciennes in Cuvier & Valenciennes, 1835) (Reported by various authors under the name *Naso tuberosus*)  
*Naso unicornis* (Forsskål in Niebuhr, 1775)  
*Naso vlamingii* (Valenciennes in Cuvier & Valenciennes, 1835)  
*Paracanthurus hepatus* (Linné, 1766)  
*Prionurus maculatus* Ogilby, 1887  
*Zebrasoma scopas* (Cuvier, 1829)  
*Zebrasoma veliferum* (Bloch, 1797) (Synonym: *Zebrasoma veliferum novacaledoniae* Borodin, 1932; reported by Fourmanoir & Laboute [1976: 210] and other authors under the name *Zebrasoma desjardini*)

#### **SPHYRAENIDAE**

A record of *Sphyraena waitii* Ogilby, 1908 by several authors (under the name *S. waittei*) needs verification.  
*Sphyraena barracuda* (Walbaum, 1792)  
*Sphyraena flavicauda* Rüppell, 1838  
*Sphyraena forsteri* (Cuvier in Cuvier & Valenciennes, 1829) (Synonym: *Sphyraena toxema* Fowler, 1904; reported from New Caledonia by authors under the name *Sphyraena japonica*, and by Laboute & Grandperrin [2000: 337] as *Sphyraena* sp. 2)  
*Sphyraena helleri* Jenkins, 1901 (Reported from New Caledonia by authors under the name *Sphyraena acutipinnis* and *Sphyraena novae-hollandiae*)  
*Sphyraena jello* Cuvier in Cuvier & Valenciennes, 1829 (Reported from New Caledonia by Laboute & Grandperrin [2000: 337] under the name *Sphyraena* sp. 1)  
*Sphyraena obtusata* Cuvier in Cuvier & Valenciennes, 1829 (Reported from New Caledonia by Fourmanoir & Laboute [1976: 185] under the name *Sphyraena chrysotaenia*)  
*Sphyraena putnamiae* Jordan & Seale, 1905 (Synonym: *Sphyraena bleekeri* Williams, 1959)  
*Sphyraena qenie* Klunzinger, 1870

#### **GEMPYLIDAE**

*Gempylus serpens* Cuvier, 1829  
*Prometichthys prometheus* (Cuvier in Cuvier & Valenciennes, 1831)  
*Ruvettus pretiosus* Cocco, 1829

#### **TRICHIURIDAE**

*Trichiurus lepturus* Linnaeus, 1758 (Synonym: *Trichiurus haumela*)

#### **SCOMBRIDAE**

*Acanthocybium solandri* (Cuvier in Cuvier & Valenciennes, 1831)  
*Auxis thazard* (Lacepède, 1801)  
*Euthynnus affinis* (Cantor, 1849)  
*Grammatocynus bicarinatus* (Quoy & Gaimard, 1824)  
*Grammatocynus bilineatus* (Rüppell, 1836)  
*Gymnosarda unicolor* (Rüppell, 1838)  
*Katsuwonus pelamis* (Linnaeus, 1758)  
*Rastrelliger kanagurta* (Cuvier, 1829)  
*Scomberomorus commerson* (Lacepède, 1802)  
*Thunnus alalunga* (Bonnaterre, 1788)  
*Thunnus albacares* (Bonnaterre, 1788)  
*Thunnus obesus* (Lowe, 1839)

#### **NOMEIDAE**

*Psenes cyanophrys* Valenciennes in Cuvier & Valenciennes, 1833

#### **CAPROIDAE**

*Antigonias capros* Lowe, 1843

#### **PARALICHTHIDAE**

*Pseudorhombus arsius* (Hamilton Buchanan, 1822)

## BOTHIDAE

- Arnoglossus dalgleishi* (Bonde, 1922) (New record from New Caledonia, based on SMNS 23571  
from Lifou, Loyalty Islands)  
*Arnoglossus japonicus* Hubbs, 1915  
*Arnoglossus macrolophus* (Alcock, 1899)  
*Arnoglossus polystipus* (Günther, 1880)  
*Arnoglossus tenuis* Günther, 1880  
*Asterorhombus annulatus* (Weber, 1913)  
*Asterorhombus fijiensis* (Norman, 1931)  
*Asterorhombus filifer* Hensley & Randall, 2003  
*Asterorhombus intermedius* (Bleeker, 1866)  
*Bothus mancus* (Broussonet, 1782) (Synonym: *Pleuronectes lunulatus* Jouan, 1861)  
*Bothus pantherinus* (Rüppell, 1830)  
*Engyprosopon bellonaensis* Amaoka, Mihara & Rivaton, 1993  
*Engyprosopon grandisquamum* (Temminck & Schlegel, 1846)  
*Engyprosopon hureau* Quéro & Golani, 1990  
*Engyprosopon longipterum* Amaoka, Mihara & Rivaton, 1993  
*Engyprosopon macrolepis* (Regan, 1908)  
*Engyprosopon maldivensis* (Regan, 1908)  
*Engyprosopon rostratum* Amaoka, Mihara & Rivaton, 1993  
*Engyprosopon septempes* Amaoka, Mihara & Rivaton, 1993 (The species has been reported from  
the Chesterfield Islands under the names *Engyprosopon longipelvis* or *Engyprosopon latifrons*)  
*Grammatobothus pennatus* Ogilby, 1913  
*Grammatobothus polyophthalmus* (Bleeker, 1866)

## SAMARIDAE

- Samaris cristatus* Gray, 1831  
*Samaris macrolepis* Norman, 1927  
*Samaris spinea* Mihara & Amaoka, 2004  
*Samariscus latus* Matsubara & Takamuri, 1951  
*Samariscus triocellatus* Woods in Schultz, Woods & Lachner, 1966

## SOLEIDAE

- Aesopias cornuta* (Kaup, 1858)  
*Aseraggodes magnoculus* Randall, 2005  
*Aseraggodes melanostictus* (Peters, 1877) (New record from New Caledonia based on SMNS  
23790 from Lifou, Loyalty Islands)  
*Aseraggodes ramsayi* (Ogilby, 1889)  
*Aseraggodes whitakeri* Woods in Schultz et al., 1966  
*Pardachirus pavoninus* (Lacepède, 1802)  
*Soleichthys heterorhinos* (Bleeker, 1856)  
*Zebrias japonica* (Bleeker, 1860)

## CYNOGLOSSIDAE

- Cynoglossus interruptus* Günther, 1880

## BALISTIDAE

- A record of *Pseudobalistes flavimarginatus* (Rüppell, 1829) by Rivaton et al. (1989: 26) needs verifica-  
tion.  
*Abalistes stellatus* (Anonymus [ex Commerson, ex Lacepède], 1798)  
*Balistapus undulatus* (Mungo Park, 1797)  
*Balistoides conspicillum* (Bloch & Schneider, 1801)  
*Balistoides viridescens* (Bloch & Schneider, 1801)  
*Canthidermis maculatus* (Bloch, 1786)  
*Melichthys vidua* (Solander, 1844)  
*Odonus niger* (Rüppell, 1836) (Synonym: *Balistes erythrodon* Günther, 1870)  
*Pseudobalistes fuscus* (Bloch & Schneider, 1801)  
*Rhinecanthus aculeatus* (Linnaeus, 1758)  
*Rhinecanthus lunula* Randall & Steene, 1983

*Rhinecanthus rectangulus* (Bloch & Schneider, 1801)  
*Rhinecanthus verrucosus* (Linnaeus, 1758)  
*Sufflamen bursa* (Bloch & Schneider, 1801)  
*Sufflamen chrysopterum* (Bloch & Schneider, 1801)  
*Sufflamen fraenatum* (Latreille, 1804) (Synonym: *Sufflamen capistratus*)  
*Xanthichthys auromarginatus* (Bennett, 1831) (Reported by several authors under the name  
    *Xanthichthys ringens* or *Balistes ringens*)

#### MONACANTHIDAE

*Acreichthys radiatus* (Popta, 1900)  
*Acreichthys tomentosus* (Linnaeus, 1758) (Reported from New Caledonia by Laboute &  
    Grandperrin [2000: 461] under the name 'Monacanthidae indéterminé 1')  
*Aluteres monoceros* (Linnaeus, 1758)  
*Aluteres scriptus* (Osbeck, 1765)  
*Amanses scopas* (Cuvier, 1829)  
*Brachaluteres jacksonianus* (Quoy & Gaimard, 1824)  
*Brachaluteres taylori* Woods in Schultz, Woods & Lachner, 1966  
*Cantherhines dumerilii* (Hollard, 1854)  
*Cantherhines fronticinctus* (Günther in Playfair & Günther, 1867)  
*Cantherhines pardalis* (Rüppell, 1835)  
*Kentrocapros flavofasciatus* (Kamohara, 1938)  
*Oxymonacanthus longirostris* (Bloch & Schneider, 1801)  
*Paraluteres prionurus* (Bleeker, 1851)  
*Paramonacanthus curtorhynchos* (Bleeker, 1855) (Reported by several authors under the name  
    *Paramonacanthus japonicus*, and by Laboute & Grandperrin [2000: 461] under the name  
    'Monacanthidae indéterminé 2')  
*Paramonacanthus lowei* Hutchins, 1997  
*Pervagor alternans* (Ogilby, 1889)  
*Pervagor aspricaudus* (Hollard, 1854)  
*Pervagor janthinosoma* (Bleeker, 1854)  
*Pervagor melanocephalus* (Bleeker, 1853)  
*Pseudalutarius nasicornis* (Temminck & Schlegel, 1846)  
*Thamnaconus modestoides* (Barnard, 1927)

#### OSTRACIIDAE

*Lactoria cornuta* (Linnaeus, 1758)  
*Lactoria diaphana* (Bloch & Schneider, 1801)  
*Lactoria fornasini* (Bianconi, 1846)  
*Ostracion cubicus* Linnaeus, 1758 (Synonym: *Ostracion tuberculatus* Linnaeus, 1758)  
*Ostracion meleagris* Shaw, 1796  
*Tetrosomus concatenatus* (Bloch & Schneider, 1801)  
*Tetrosomus gibbosus* (Linnaeus, 1758)

#### TRIODONTIDAE

*Triodon macropterus* Lesson in Cuvier & Valenciennes, 1829

#### TETRAODONTIDAE

*Arothron alboreticulatus* (Tanaka, 1908)  
*Arothron caeruleopunctatus* Matsura, 1994  
*Arothron hispidus* (Linnaeus, 1758)  
*Arothron immaculatus* (Bloch & Schneider, 1801)  
*Arothron manilensis* (Marion de Procé, 1822)  
*Arothron mappa* (Lesson, 1830)  
*Arothron meleagris* (Lacepède, 1798)  
*Arothron nigropunctatus* (Bloch & Schneider, 1801)  
*Arothron stellatus* (Bloch & Schneider, 1801)  
*Canthigaster amboinensis* (Bleeker, 1865) (New record from New Caledonia based on SMNS  
    23610 from Lifou, Loyalty Islands)  
*Canthigaster bennetti* (Bleeker, 1854)

*Canthigaster compressa* (Marion de Procé, 1822)  
*Canthigaster coronata* (Vaillant & Sauvage, 1875)  
*Canthigaster epilampra* (Jenkins, 1903)  
*Canthigaster janthinoptera* (Bleeker, 1855)  
*Canthigaster ocellicincta* Allen & Randall, 1977  
*Canthigaster papua* (Bleeker, 1848)  
*Canthigaster rivulata* (Temminck & Schlegel, 1850)  
*Canthigaster solandri* (Richardson, 1844)  
*Canthigaster valentini* (Bleeker, 1853)  
*Chelonodon patoca* (Hamilton Buchanan, 1822) (New record from New Caledonia based on AMS  
IB.7080 from southern Grande Terre; specimen identified by G. Hardy)  
*Lagocephalus sceleratus* (Gmelin, 1789)  
*Sphoeroides pachygaster* (Müller & Troschel, 1848)  
*Torquigener brevipinnis* (Regan, 1902)  
*Torquigener hypselogeneion* (Bleeker, 1852)  
*Torquigener pallimaculatus* Hardy, 1983  
*Torquigener tuberculiferus* (Ogilby, 1912)  
*Tylerinus spinosissimus* (Regan, 1908)

#### DIODONTIDAE

*Cyclichthys spilostylus* (Leis & Randall, 1982)  
*Cyclichthys orbicularis* (Bloch, 1785)  
*Diodon holocanthus* Linnaeus, 1758  
*Diodon hystrix* (Linnaeus, 1758)  
*Diodon liturosus* Shaw, 1804  
*Lophodiodon calori* (Bianconi, 1855)

#### MOLIDAE

*Masturus lanceolatus* (Liénard, 1841)  
*Mola mola* (Linnaeus, 1758)  
*Mola ramsayi* (Giglioli, 1883)  
*Ranzania laevis* (Pennant, 1776)



© IRD/Pierre Laboute

*Acanthurus olivaceus*

© IRD/Pierre Laboute

*Caesio cuning*

© IRD/Pierre Laboute

*Chaetodon baronessa*

© IRD/Pierre Laboute

*Oxycheilinus bimaculatus*

© IRD/Pierre Laboute

*Gymnothorax nudivomer*

© IRD/Georges Bertolaat

*Lethrinus miniatus*

© IRD/Pierre Laboute

*Lutjanus adetii*

© IRD/Pierre Laboute

*Manta birostris*



© IRD/Pierre Laboute

*Mulloidichthys vanicolensis*

© IRD/Pierre Laboute

*Oxymonacanthus longirostris*

© IRD/Pierre Laboute

*Paracirrhites forsteri*

© IRD/Pierre Laboute

*Parupeneus indicus*

© IRD/Pierre Laboute

*Pomacanthus semicirculatus*

© IRD/Pierre Laboute

*Sargocentron rubrum*

© IRD/Pierre Laboute

*Scarus rubroviolaceus*

© IRD/Pierre Laboute

*Stegostoma fasciatum*

**DOCUMENTS  
SCIENTIFIQUES  
et TECHNIQUES**

**II7**

**Volume spécial**

# **COMPENDIUM of marine species from New Caledonia**



Edited by  
**CLAUDE E. PAYRI**  
Bertrand RICHER DE FORGES

**IRD**

Institut de recherche  
pour le développement

CENTRE DE NOUMÉA

# **DOCUMENTS SCIENTIFIQUES et TECHNIQUES**

---



Publication éditée par:  
Centre IRD de Nouméa  
BP A5, 98848 Nouméa CEDEX  
Nouvelle-Calédonie  
Téléphone : (687) 26 10 00  
Fax : (687) 26 43 26

L'IRD propose des programmes regroupés en 5 départements pluridisciplinaires :

- I DME Département milieux et environnement
- II DRV Département ressources vivantes
- III DSS Département sociétés et santé
- IV DEV Département expertise et valorisation
- V DSF Département du soutien et de la formation des communautés scientifiques du Sud

**Modèle de référence bibliographique à cette revue :**

Adjeroud M. *et al.*, 2000. Premiers résultats concernant le benthos et les poissons au cours des missions TYPATOLL.  
*Doc. Sci. Tech.* II 3, 125 p.

Première de couverture : Récif corallien (Côte Ouest, NC) © IRD/C.Geoffray  
Vignettes : voir les planches photographiques

Quatrième de couverture : *Platygyra sinensis* © IRD/C.Geoffray

Matériel de plongée sous-marine  © IRD/C.Geoffray	L'Aldric, moyen naviguant de l'IRD  © IRD/J.-M. Boré
Récoltes et photographies sous-marines en scaphandre autonome  © IRD/J.-L. Menou	Traitement des récoltes en laboratoire  © IRD/L. Mattio

CONCEPTION/MAQUETTE/MISE EN PAGE  
MAQUETTE DE COUVERTURE  
PLANCHES PHOTOGRAPHIQUES  
TRAITEMENT DES PHOTOGRAPHIES

JEAN PIERRE MERMOUD  
CATHY GEOFFRAY/ MINA VILAYLECK  
CATHY GEOFFRAY/ JEAN-LOUIS MENOU/ GEORGES BARGIBANT  
NOËL GALAUD

La traduction en anglais des textes d'introduction, des Ascidies et des Echinodermes a été assurée par EMMA ROCHELLE-NEWALL, la préface par MINA VILAYLECK.

Ce document a été produit par le Service ISC, imprimé par le Service de Reprographie du Centre IRD de Nouméa et relié avec l'aimable autorisation de la CPS, financé par le Ministère de la Recherche et de la Technologie.

# **COMPENDIUM OF MARINE SPECIES FROM NEW CALEDONIA**

Edited by

CLAUDE E. PAYRI, BERTRAND RICHER DE FORGES

---