

Geology of the New Caledonia region and its implications for the study of the New Caledonian biodiversity

Bernard PELLETIER

*Institut de Recherche pour le Développement (IRD) Nouméa,
UMR Géosciences Azur, BP A5 Nouméa, Nouvelle-Calédonie
Bernard.Pelletier@noumea.ird.nc*

Introduction

The New Caledonian exclusive economic zone is located in the Southwest Pacific between Australia and the Vanuatu archipelago (formerly New Hebrides). It extends over 1200 km from north to south and 1800 km from west to east, from the Chesterfield Islands up to the Matthew and Hunter Islands at the southern tip of the Vanuatu archipelago (Figure 1, page 8). The main islands of New Caledonia are located on two parallel NW-SE trending ridges. The largest island (the 400 km-long and 50 km-wide Grande Terre) and subordinate islands including the Belep islands in the north and the Isle of Pines in the south are supported by the New Caledonia Ridge. The Loyalty Islands are supported by the Loyalty Ridge.

Geologically, the area is schematically composed of a series of NW-SE trending ridges and basins formed by i) stretching and spreading along the eastern margin of Australia during Late Cretaceous-Paleocene times and ii) Eocene convergence which was responsible for the emplacement of one of the largest ophiolitic complexes in the world: the New Caledonian ophiolitic nappe (a slice of oceanic lithospheric mantle) from which weathering formed one of the world's largest reservoirs of nickel. The ridges and basins are nowadays supported by the Australia plate that subducts beneath the Vanuatu active volcanic arc. The southernmost segment of the arc supports the active volcanic islands of Matthew and Hunter. These latter islands are thus on a plate (a micro-plate) that differs from the one supporting the main islands of New Caledonia territory.

After reviewing the different morphostructural units and discussing the geological history of the New Caledonia area, I will point out some facts that may be relevant in the study and understanding of New Caledonia biodiversity.

Geological of the morphostructural units of the New Caledonia region

From west to east, the main geological units between Australia and Vanuatu include the Tasman Sea Basin, the Lord Howe Rise with subordinate basins, ridges and chains, the New Caledonia Basin, the New Caledonia Ridge, the South Loyalty Basin, the Loyalty Ridge, the North Loyalty Basin and the Vanuatu Ridge (Missègue *et al.*, 1991; Smith and Sandwell, 1997; ZoNéCo 1998) (Fig. 1 plate 1/1 and Fig. 2 plate 2/1).

The Tasman Sea Basin

To the southeast of Australia the Tasman Sea Basin is a wide (up to 2000 km) and deep (4000 m) basin floored with oceanic crust formed by spreading from the Cretaceous (85 Ma: Santonian to Early Campanian) to Earliest Eocene (52 Ma) (Hayes et Ringis, 1973; Weissel and Hayes 1977; Gaina *et al.*, 1998). In the central part of the basin, the N-S trending Tasmanid volcanic chain is an Oligocene to Late Miocene hot spot chain showing the northward drift of the Australia plate (Vogt and Conolly, 1971; Mc Dougall and Duncan, 1988).

The Lord Howe Rise

The Lord Howe Rise is a main bathymetric feature which averages 400 km in width and extends over 1600 km from the Challenger Plateau off New Zealand to the Chesterfield area. Water depths on the crest are 1200 to 750 m. In the north, at the latitude of New Caledonia, the Lord Howe Rise includes several basins, ridges and chains. They are from west to east: the Dampier Ridge, the Middleton basin, the Chesterfield/Bellona Plateau, the Faust basin, the Lord Howe Rise crest, the Fairway Basin and the Fairway Ridge. The Lord Howe Rise is interpreted as a thinned fragment of continental crust split from Gondwana by Cretaceous rifting and subsequent spreading in Tasman Sea Basin, and it is

probably composed of Paleozoic basement overlain by rift basins with up to 4 km of Mesozoic and Cenozoic sediments (Willcox *et al.*, 2001; Van de Beuque *et al.*, 2003; Exon *et al.*, 2004). Stratigraphy also indicates an emergence and major unconformity from the Late Eocene to Early Oligocene (Burns *et al.*, 1973), which correlates with described compressive structures of that age (Lafoy *et al.*, 1994; Symonds *et al.*, 1999; Auzende *et al.*, 2000).

The Lord Howe rise has Cenozoic seamount chains as exemplified by N-S trending volcanic edifices on its crest (Van de Beuque *et al.*, 1998; Exon *et al.*, 2004) and the Oligocene-Miocene Lord Howe seamount chain on its western side. The Chesterfield/Bellona plateau is supported by five guyots that constitute the northern and oldest volcanoes (Late Oligocene ?) along the Lord Howe hotspot chain (Missegeue et Collot, 1987).

The 800 km long, 130 km wide and NW-SE to NNW-SSE trending and southward deepening (1000 to 3000 m) Fairway Basin, firstly interpreted as oceanic in nature (Ravenne *et al.*, 1977; Mignot, 1984; Eade, 1988; Uruski and Wood, 1991; Van de Beuque, 1999), is now considered to be floored by stretched-thinned continental crust based on the presence of salt diapirs derived from Cretaceous series (Auzende *et al.*, 2000), gravity modeling (Vially *et al.*, 2003) and extensional-type horst and graben structures of the crust (Lafoy *et al.*, 2005). It is proposed that the formation of the Fairway Basin took place during the Late Cretaceous (95-65 Ma) by continental stretching, at the same time as the stretching in the Middleton Basin (Lafoy *et al.*, 2005).

The 600 km-long and NW-SE trending Fairway Ridge, culminating in its northernmost part at the Lansdowne bank, thins and deepens southward. The origin of the ridge is still controversial. Previously interpreted as a ridge of oceanic nature (Ravenne *et al.*, 1977; Mignot, 1984) and as an oceanic piece of the New Caledonia basin crust overthrust along the Lord Howe Rise (Lafoy *et al.*, 1994; van de Beuque, 1999; Auzende *et al.*, 2000), it is now considered as thinned continental crust (Vially *et al.*, 2003; Lafoy *et al.*, 2005).

The New Caledonia Basin

The New Caledonia Basin extends from west of Northern New Zealand to west of New Caledonia, parallel to the Lord Howe Rise. The deepest (3600-3700 m) northern part (north of 22°30'S) strikes NW-SE while the central part (3000 m deep) strikes NNW-SSE. Origin of the basin is controversial from oceanic type (Shor *et al.*, 1971; Dubois *et al.*, 1974; Weissel et Hayes, 1977; Willcox *et al.*, 1980; Kroenke, 1984; Mignot, 1984 ; Sutherland, 1999 ; Auzende *et al.*, 2000) to thinned continental type (Etheridge *et al.*, 1989 ; Uruski et Wood, 1991 ; Sdrolias *et al.*, 2003 ; Vially *et al.*, 2003 ; Lafoy *et al.*, 2005). The horst and graben structure of the crust with westward tilted blocks and a sedimentary section up to 8 km thick suggests a thinned continental crust for the northern NW-SE segment. On this segment the crust dips to the east, toward the western margin of the New Caledonia ridge. At the base of this margin, buried deformation features interpreted as the result of compression have been recognized (Rigolot and Pelletier, 1988). In contrast to the northern segment, the central segment of the basin, with magnetic lineations and an axial ridge buried by a 4 km-thick sedimentary sequence, is interpreted as a segment floored with oceanic crust that formed during the Paleocene (possibly from 62 to 56 Ma) after Late Cretaceous–Earliest Paleocene stretching (Lafoy *et al.*, 2005).

The New Caledonia Ridge

The New Caledonia Ridge is the NW-SE trending northern segment of the 70-100 km-wide Norfolk Ridge which extends over 1500 km from the d'Entrecasteaux Reef to the northern tip of New Zealand. As for the Lord Howe Rise, the Norfolk ridge is interpreted to be a continental ribbon detached from Gondwana. Geology of the Grande-Terre of New Caledonia (Lillie and Brothers, 1970; Paris, 1981; Picard, 1999) has great similarities with that of New Zealand. Main geologic features of the island have been tentatively extended along the submerged northward (Collot *et al.*, 1988) and southward (Rigolot, 1988) segments of the ridge.

The island is composed of a series of various terranes assembled during two tectonic events : a Late Jurassic to Early Cretaceous tectonic collage (Paris, 1981 ; Meffre 1995 ; Aitchison *et al.* 1998) and a Late Eocene subduction/collision resulting in the emplacement at the Latest Eocene (38-34 Ma) of a large ophiolitic nappe (Avias 1967; Paris, 1981 ; Collot *et al.*, 1987 ; Aitchison *et al.*, 1995 ; Cluzel *et al.*, 1994, 2001).

The pre-Cretaceous terranes, mainly located in the central chain, are unconformably overlain by Upper Cenomanian (Late Cretaceous) to Upper Eocene sediments, and include disrupted Late Carboniferous ophiolite (Meffre *et al.*, 1996 ; Aitchison *et al.*, 1998), mid-Triassic to late Jurassic volcano-sedimentary arc terrane (Meffre *et al.*, 1996), Mid Permian to Late Jurassic volcano-sedimentary arc terrane (Campbell *et al.*, 1985), post-Liassic unit composed of oceanic crust and volcano-sedimentary distal deposits (Cluzel, 1996) and affected by a Late Jurassic high pressure metamorphism (150 Ma : Blake *et al.*, 1977).

The post-Early Cretaceous terranes also involved in the Late Eocene major event include:

- an unmetamorphosed Upper Cretaceous to Upper Eocene sedimentary pile well exposed along the western side of the Grande-Terre. This pile is composed of two sequences separated by an unconformity (Paris, 1981; Cluzel *et al.*, 2001). The lower sequence includes an Upper Cretaceous fining-upward clastic series of conglomerates, sandstones, coaly siltstones and volcanic rocks, overlain by Paleocene to Middle Eocene pelagic limestones and cherts, indicative of a deepening of the deposit environment. The upper sequence is a Upper Eocene (Upper Bartonian to Priabonian) flysch formation (Nouméa-Bourail and Népoui flyschs), deposited after a short period of deformation and erosion and showing a coarsening upward sequence with (a) basal neritic limestones, (b) a «lower flysch» member of marls and calcareous sandy marls, (c) an «upper flysch» member composed of fine-grained calcareous turbidites interbedded with mafic breccias, and (d) a «wildflysch» member that incorporates blocks and olistoliths of siliceous shales, limestones, basalts and flysch, topped by an olistostrome recording the Late Eocene tectonic paroxysm.

- a mafic unit of oceanic basalts interbedded with argillite and cherts (named basalt nappe or Poya unit) of Late Cretaceous (Campanian) to Late Paleocene-Earliest Eocene age (85-55 Ma), with back-arc or fore-arc affinities (Routhier, 1953; Espirat, 1963; Eissen *et al.*, 1998; Cluzel *et al.*, 1997, 2001). This unit, severely sheared and folded, is mainly exposed along the northern half part of the west coast but also outcrops along the east coast; it always underlies the ultramafic nappe and tectonically overlies the Upper Eocene sedimentary rocks.

- mafic high pressure-low temperature metamorphic units (Pouebo and Diahot units) located in the northeastern part of the Grande Terre (Brothers, 1974; Paris, 1981; Yokoyama *et al.*, 1986; Maurizot *et al.*, 1989; Black *et al.*, 1993; Clarke *et al.*, 1997; Cluzel *et al.*, 1995; Baldwin *et al.*, 1999; Carson *et al.*, 1999; Rawling and Lister, 2002; Fitzherbert *et al.*, 2004. Spandler *et al.*, 2005).

The units and especially the Pouebo unit are interpreted to be the equivalent of the basalt nappe metamorphosed under blueschist to eclogitic facies conditions, the latter (20 kbar, 650°C) indicating an underthrusting equivalent to a depth of 60-70 km. Radiometric dating of the metamorphism peak is 44 Ma (Spandler *et al.*, 2005) while those from the cooling ages range from 40 to 34 Ma (Baldwin *et al.*, 1999), indicating a rapid unroofing and exhumation of the metamorphic units in the north and a synchronism with the final emplacement of the ophiolitic nappe (38-34 Ma) in the west and south of the island.

- the ophiolitic nappe mainly composed of peridotites and well exposed in the southern part of the island (3 km thick in the southern massif), Belep and Pines islands and as a series of klippen along the western northern half of the island (Avias, 1967; Guillon et Routhier 1971; Guillon, 1975; Prinzoffer *et al.*, 1980). If the age of its emplacement is Late Eocene (Paris *et al.*, 1979), the age of these mantellic rocks is not accurate and considered to be Late Cretaceous or older from radiometric datings of associated mafic and felsic dikes yielded Late Cretaceous (100-80 Ma) and Eocene (42-52 Ma) ages (Paris, 1981; Prinzoffer, 1981). The peridotites are mainly harburgites and are the mother rocks for the nickel of New Caledonia.

The Oligocene is characterised by a lack of marine sediments, and post-obduction granodiorite intrusions (St Louis and Koum) radiometrically dated from 32 to 24 Ma (Guillon, 1975) and interpreted as the result of short-lived convergence episode along the western margin of the New Caledonia ridge (Cluzel *et al.*, 2005). The peridotites have been extensively weathered under aerial conditions since Oligocene, leading to the development of thick Ni-rich lateritic mantles (Trescases, 1973, 1975; Latham, 1986) and relict lateritic landsurfaces (Chevillotte *et al.*, 2006). Brittle extensional deformation plays an important role in the post-obduction morphotectonic evolution of the island (Leguere, 1976; Lagabrielle *et al.*, 2005; Chardon and Chevillotte, 2006). This extension which may initiated in the Oligocene is expressed in the Neogene by the disruption of land surface formed during Oligocene planation and in the outcrop at Nepoui of Lower-Middle Miocene fluvial conglomerates with shallow water marine limestones (Coudray, 1976). Neogene ridge-normal then ridge-parallel to oblique extensional tectonics are also responsible for the shape and subsidence of the New Caledonia ridge margins (Daniel *et al.*, 1976; Dugas and Debenay, 1978; Bitoun and Récy, 1982; Rigolot, 1989; Chardon and Chevillotte, 2005; Flamand 2006). The Grande Terre displays one of the largest barrier-reefs in the world, isolating a locally-wide lagoon. The barrier reef settled in the Early Pleistocene (Coudray, 1976). Vertical motions from the 125 Ka reef indicate different tectonic blocks and a general slow subsidence (0.03 to 0.16 mm/year) of the coast except in the southeastern part of the Grande-Terre and île des Pins where coasts are uplifted (Launay and Récy, 1972; Launay, 1985; Cabioch, 1988, Cabioch *et al.*, 1996) and where quaternary faults have been observed (Lafoy *et al.*, 2000; Lagabrielle *et al.*, 2005 ; Flamand, 2006) and seismicity occurs (Régnier *et al.*, 1999; Pillet and Pelletier, 2004).

The South Loyalty Basin

Parallel to the Norfolk/New Caledonia Ridge, the South Loyalty Basin (called also the West Loyalty Basin) is a 1300 km-long, narrow (45-65 km wide), and northward deepening (from 2000 to 3800 m) basin with oceanic crust dipping northwestward and filled with thick (up to 8 km) sediments which are considered to be mainly post Eocene in age (Bitoun and Recy, 1982 ; Pontoise *et al.*, 1982 ; Collot, *et al.*, 1987). Geophysical data suggests that the oceanic basement is the continuity of the ophiolitic nappe of the Grande Terre. The age of the crust is unknown but considered as pre Late Cretaceous (Collot *et al.*, 1987) or Late Cretaceous to Paleocene (Cluzel *et al.*, 2001).

The Loyalty Ridge

The loyalty ridge is a narrow ridge parallel to the South Loyalty Basin and Norfolk Ridge and more or less continuous from the Cook Fracture zone in the South to the d'Entrecasteaux zone in the north. It is composed of a series of seamounts and guyots and supports the Loyalty islands.

The geology of the ridge is poorly known and its origin and nature are unknown. Parallelism with other ridges bordering the Australian margin suggests an old and continental origin (Monzier, 1993). However, taking into account its possible link with the d'Entrecasteaux zone (an Eocene subduction zone) and for convenience in the understanding the geology of New Caledonia, the Loyalty Ridge is considered as an Eocene island arc in most of the reconstructions, (Maillet *et al.*, 1983; Kroenke, 1984; Eissen *et al.*, 1998 ; Cluzel *et al.*, 1994, 2001 ; Crawford *et al.*, 2003 ; Sdrolias *et al.*, 2003 ; Schellart *et al.*, 2006). Middle to Upper Oligocene non orogenic volcanism has been also proposed for the origin of the ridge (Rigolot, 1989; Monzier, 1993). Only few volcanic rocks have been recovered in two areas. Upper Miocene (9-11 Ma) alkalic basalts (Baubron *et al.*, 1976) outcrop on Mare island. Submersible dives off Mare along the eastern flank of the ridge (Monzier *et al.*, 1989) recovered volcanic breccias. Middle Oligocene (32 Ma) alkaline rhyolites, Middle Oligocene tuffaceous sandstones, Middle Upper Oligocene chinks, Upper Oligocene (27 Ma) alkalic basalts, Lower Miocene (20 Ma) back-arc basalts, and algae and reefal limestones with reworked Eocene-Oligocene and Mio-Pliocene fauna (Monzier, 1993). The ridge appears to be composed, at least partly, of non orogenic alkaline volcanics of Middle to Late Oligocene and Late Miocene age, the youngest being likely a part of a N-S trending hot spot track (Rigolot *et al.*, 1988). None of the recovered rocks argues for a volcanic arc origin, although this hypothesis, possible and attractive, is widely accepted in the literature.

The Loyalty islands are mainly composed of Late Miocene to Pleistocene uplifted reef formations covering the basement (Chevalier, 1968 ; Marshall et Launay, 1978 ; Bourrouilh, 1996; Carrière, 1987; Guyomard *et al.*, 1996). The varying altitude of the islands shows the bulge of the Australia plate in front of its subduction eastward beneath the Vanuatu arc (Dubois *et al.*, 1974, 1977, 1988). The islands diachronously emerged during the Pleistocene (possibly in the Latest Pliocene for Mare) and are still emerging and uplifting (as exemplified by Ouvea) except the ones which have passed the top of the bulge and are thus subsiding.

The North Loyalty Basin

The North Loyalty Basin (named also East Loyalty Basin) is a deep (3000 to 5000 m) basin floored with oceanic crust dated of pre Middle Eocene age in its northernmost part (Andrews *et al.*, 1975). Bounded northward by the d'Entrecasteaux zone, it is the remaining part of a larger basin which disappears eastward in the active Vanuatu subduction zone. Initially interpreted to have formed by spreading in Early Eocene and regarded as the old part of the South Fiji Basin (Lapouille, 1982; Weissel *et al.*, 1982), it is now considered to have formed in Late Eocene (44 to 35 Ma; Sdrolias *et al.*, 2003) as a back arc basin of the Loyalty arc (Maillet *et al.*, 1983; Cluzel *et al.*, 2001; Schellart *et al.*, 2006).

The Vanuatu Trench and Ridge, the North Fiji Basin and the Vitiav Trench Lineament

The Vanuatu Ridge is a 1500 km-long active volcanic arc related to the subduction of the Australia plate since the Late Miocene (12-10 Ma). Oldest known arc volcanic rocks from the Vanuatu ridge are however Early Miocene in age (Mitchell and Warden, 1971 ; Carney and MacFarlane, 1982) and related to the fossil west-dipping Vitiav subduction zone along which Pacific plate subducted, Late Miocene subduction reversal and initiation of east-dipping Vanuatu subduction being due to collision of the Ontong Java Plateau and the Melanesian border plateau (Packham, 1973; Kroenke, 1984; Brocher, 1985; Pelletier and Auzende, 1996). The Vanuatu arc rotated clockwise leading to the formation of the active north Fiji back arc complex basin (Chase *et al.*, 1971 ; Falvey, 1975 ; Auzende *et al.*, 1988, 1995).

The Vanuatu Trench is a segment of the present-day Australia-Pacific converging plate boundary along which Australia plate dips eastward and is consumed. Relative motion of convergence at trench is ENE-WSW and rate of motion varies along the trench and is about 12 cm/year at the latitude of the Loyalty islands (Dubois *et al.*, 1977; Louat et Pelletier 1989 ; Pelletier *et al.*, 1998 ; Calmant *et al.*, 1995, 2003). Near 22°S, the Loyalty Ridge enters the trench and subducts/collides with the Vanuatu arc since 300 ka (Monzier *et al.*, 1989), reducing the convergence motion south of the impact point and forming sinistral E-W strike-slip motion across the arc and isolating a micro plate (Louat and Pelletier 1989 ; Calmant *et al.*, 2003) on which Matthew and Hunter islands are active volcanoes (Maillet *et al.*, 1986). This incipient collision is also supposed to have tectonic effects on the Loyalty Ridge (Lafoy *et al.*, 1996).

Tectonic evolution of the New Caledonia domain

Numerous Cretaceous to Cenozoic reconstructions of the Southwest Pacific and New Caledonia region have been proposed in the past years (Kroenke 1984 ; Yan and Kroenke, 1993; Veevers, 2000 ; Muller *et al.*, 2000 ; Cluzel *et al.*, 2001 ; Sutherland *et al.*, 2001 ; Hall, 2002 ; Crawford *et al.*, 2003 ; Sdrolias *et al.*, 2003 ; Schellart *et al.*, 2006). Tectonic evolution concerning the New Caledonia domain can be divided into 5 stages (Fig. 3, plate 2/2).

The pre-Late Cretaceous period

The Paleozoic to Early Cretaceous period is thought to be marked by a subduction zone along the eastern margin of Gondwana. It ended with a Late Jurassic to Early Cretaceous tectonic orogeny (correlated with the Rangitata orogeny in New Zealand) that resulted in a collage of different units. The various Late Carboniferous to Late Jurassic units found in New Caledonia assembled at that time to form the old core of New Caledonia (Paris, 1981 ; Meffre, 1995 ; Cluzel *et al.*, 2001).

The Early Late Cretaceous (120-100 Ma) to Earliest Eocene (55-50 Ma) period: marginal rifting and spreading along the east Gondwana margin

This extensional tectonics period is marked by dislocation of the east Gondwana margin by Late Cretaceous rifting and subsequent Latest Cretaceous to Latest Paleocene/Earliest Eocene spreading. This results in a series of at least two thinned fragments of continental crust (the Lord Howe Rise, the Norfolk Ridge and possibly the Loyalty Ridge ? and/or a ridge further to the east ?) and at least three main basins floored with oceanic crust (the Tasman Sea Basin (100-85 Ma stretching, 85/80-52 Ma spreading), the New Caledonia Basin (95-62 ? Ma stretching, 62-56 Ma ? spreading) and the South Loyalty Basin (100/80 ? or 85/80 -55 Ma ? spreading). Ages of the crust of the two latter basins still derive from interpretations.

The major problem in the reconstruction for this period is the nature of the plate boundary east of the area. Reconstructions have suggested this boundary is an east dipping subduction zone, a west-dipping subduction zone, a strike-slip boundary, or no boundary at all. The recent reconstructions (Cluzel *et al.*, 2001 ; Crawford *et al.*, 2003; Shellart *et al.*, 2006) propose a continuous west-dipping subduction of the Pacific plate that rolled back eastward to accommodate the basins opening. However, this attractive model does not fit well with the « classic » formation of successive back-arc basins in which the closest basin to the trench is the youngest, because the basins are thought to be more or less similar in age (except if an older age -Early Cretaceous- for the South Loyalty Basin is chosen). Also, the associated volcanic arc accompanying this long term subduction is still largely undocumented, even if one may consider that it disappeared by erosion or it is hidden beneath younger volcanic arcs or sediments.

The Early Eocene (55-50 Ma) to Latest Eocene/Early Oligocene period : lithospheric shortening and the New Caledonian orogen

This period is marked by convergence inside the previously dismembered east Gondwana margin, by the partial closure of North Loyalty Basin and finally by the emplacement of the ophiolitic nappe in New Caledonia. Recent reconstructions (Cluzel *et al.*, 2001 ; Schellart *et al.*, 2006) propose a relatively long-lived intra-oceanic east-dipping subduction zone inside the South Loyalty Basin during Early to Middle Eocene. Subduction was locked in the Late Eocene (38-34 Ma) by the underthrusting of the Norfolk ridge, resulting in the thrusting over the New Caledonia block of mafic and ultramafic units with coeval (60-70 km) exhumation of metamorphic rocks by buoyancy-driven uplift. In the final stage, the convergence motion jumped westward along the west margin of the New Caledonia ridge, as suggested by compressive features at the toe of the western margin, the abandoned slab beneath southern New Caledonia (Régnier, 1988) and the post-orogenic Lower Oligocene intrusives. Compression also affected the Fairway Ridge and Lord How Rise.

In the above-mentioned model, a large portion of the oceanic lithosphere of the wide South Loyalty Basin would have been absorbed in subduction, and the subduction would be responsible for the development of the Loyalty Ridge arc and the North Loyalty back-arc basin. However, the fore arc of this subduction zone does not resemble classic fore arc domain, and volcanic rocks with arc affinity are still unknown in the Loyalty ridge, and the direction of spreading in the North Loyalty Basin, deduced from E-W trending magnetic anomaly lineations, is parallel -instead of normal- to the arc and the subduction zone. An alternative model proposed here is to consider a narrow original North Loyalty basin from which a small panel of lithosphere forming the foot of the New Caledonia Ridge's margin has been obducted, with no requirement of Eocene subduction to create the Loyalty Ridge.

The Oligocene to Late Miocene period : planation, extensional tectonics and subsidence, volcanism
This poorly-documented post orogenic period is marked by (low to moderate ?) uplift, erosion and planation of the New Caledonia ridge. Isostatic uplift is likely accompanied by ridge-normal extensional tectonics and subsidence of the margins, and could be due to the buoyancy force of the orogenic root thickened during collision, to loading by the overthrust oceanic lithosphere and to erosion. Ni-rich lateritic mantles likely mainly developed during this period of extensive alteration of the peridotites.

This period is also marked by alkaline volcanism related to several hot spots that affect the Australia plate, as exemplified by the Tasmanid chain in the Tasman Sea Basin, the Lord Howe volcanic chain and the volcanic edifices along the Norfolk Ridge/Loyalty Ridge.

The Late Miocene (10 Ma) to Present period : extension, initiation of subduction, bulge-related deformations and arc-ridge subduction/collision

The Latest Miocene is marked, to the east of the New Caledonia domain, by the initiation of the east-dipping Vanuatu subduction zone. This zone of convergence, behind which the active North Fiji Basin opened and along which part of the North Loyalty Basin (and other basins and features ?) was consumed, appears to play a significant role in the Late Neogene evolution of New Caledonia and Loyalty Ridges, as shown by (1) Late Neogene ridge-parallel to oblique extensional tectonics that affected the ridges, (2) Quaternary faults, bulge-related Quaternary vertical motions of the Loyalty islands and southern part of the New Caledonia Ridge, (3) Loyalty Ridge-Vanuatu arc interaction-related deformation and (4) shallow seismicity of relatively low intensity in the Grande Terre (intense seismicity, however, occurs in the easternmost part of the Loyalty Ridge close to the active Vanuatu plate boundary).

Some geological aspects for the understanding of the New Caledonia biodiversity and endemism

The problem of lands for refuge of the Gondwanian flora and fauna

The geology of the Grande Terre of New Caledonia indicates that after the Cretaceous the Norfolk Ridge/New Caledonia Ridge was below sea level up to the Late Eocene (or possibly up to the Middle Eocene), thus for a period of about 20 Ma (from 65 to 45 Ma). Indeed, the Paleocene-Lower Eocene pelagic limestones and cherts following the Upper Cretaceous fining-upward clastic sequence indicate relatively deep water deposits. Presence of an island on the New Caledonia Ridge is established since the Late Eocene, the size of which was larger than today since the d'Entrecasteaux Reef area, the wide northern lagoon, the wide southern lagoon, as well as parts of the upper slope of the present-day margins were probably above sea level. Other islands likely existed in the Late Eocene-Early Oligocene, on the Lord Howe Rise, Fairway Ridge and Loyalty Ridge. Islands also existed in Late Oligocene on the Chesterfield/Bellona zone, and in Late Miocene on the Loyalty Ridge. Thus, one may infer that the Gondwanian fauna and flora were introduced in New Caledonia during or after the Middle to Late Eocene.

If New Caledonia was not a refuge for the Gondwanian biodiversity, did other land -except Australia- exist in the region before the Middle-Late Eocene ? This in turn poses the major question concerning the nature of the plate boundary east of Australia before 45-50 Ma. If we consider (as proposed by Veevers *et al.*, 2000; Cluzel *et al.*, 2001; Crawford *et al.*, 2003; Schellart *et al.*, 2006) that a west dipping subduction continuously proceeded since that time during rifting and subsequent basin spreading, therefore a volcanic arc must have been always active. This provides the possibility for the building of an alignment of islands likely restricted in size east of the Norfolk-New Caledonia Ridge, such as the present-day islands of the active Tonga or Vanuatu arcs.

The influence of the New Caledonian ophiolite

It is well known that the nature of the ultra basic ophiolitic nappe of New Caledonia plays an important role in the floral endemism of the island. The peculiar speciation and diversification are driven by the unusual type of ultra basic rock-derived soils rich in metallic elements and depleted in mineral salts. Most of the flora living on lateritic soils is endemic. Today ultrabasic rocks and associated soils only cover about one third of the island. The ophiolitic nappe constituting the top of the tectonic pile of the Late Eocene orogen, it is likely that when it emerged the Grande-Terre island was fully covered by the ophiolite, leading to species selection. However, the present-day ophiolitic nappe lacks a complete crustal sequence (sheeted dike complex and mafic rocks) and we do not know if the mafic crustal sequence existed on the top of the ultramafic rocks or if it has been removed by erosion.

An other point about the possible role of the ophiolitic nappe for the biodiversity is alkaline hydrothermalism resulting from reactions between water and upper mantle rocks (serpentinization). Recently (Pelletier *et al.*, 2006) an extensive active alkaline hydrothermal field has been revealed by 2004 and 2005 swath mapping and scuba diving in the floor of Prony Bay which developed in the large peridotitic massif at the southern tip of the Grande Terre, the “Prony aiguille” chimney was previously known (Launay and Fontes, 1985). Uncommon characteristics of this alkaline hydrothermal system are similar (same high pH, same deposits of CaCo₃ and Mg hydroxyde and same peridotitic substratum) to those of the Lost City site recently discovered on the Mid-Atlantic Ridge (Kelley *et al.*, 2001) and which produces methane- and hydrogen-rich fluids serving as energy sources for archaeal and eubacterial communities (Kelley *et al.*, 2005). Such alkaline systems which derive from hydration of the outcropping ultramafic rocks is interesting because it may share several characteristics with hydrothermal environments at the beginning of life on Earth. The Prony Bay site is accessible by scuba diving and thus may be of wide interest. It is possible that such other active or recent hydrothermal sites exist in the present-day submerged portions of the remaining ophiolitic nappe (parts of southeastern and eastern lagoons of the Grande Terre). Probably, such hydrothermal fields already existed in the past since the Late Eocene emplacement of the ophiolitic nappe, and may have played a first-order role in the evolution of the New Caledonian biodiversity.

Pleistocene sea level variation

The sea level has 100 Ka cyclic high and low stands since the last 0.9 Ma (Shackelton, 1987, 2000). Low stands were 120-130 m below present-day sea level and high stands were sometimes 5 to 10 m above present-day sea level. Such high amplitude and rapid variations may have significant effects on the distribution of the New Caledonia marine biodiversity. Taking into account the slow subsidence of the margins, the wide lagoon around the Grande-Terre with depths much lower than 120 m has been largely emerged and submerged several times during the last million years. Detailed bathymetry of the lagoon however suggests that lakes with brackish water existed during sea falls in the deepest parts of the lagoon, the depths of the passes being generally shallower than that the lagoon itself (Dugas *et al.*, 1980; Chevillotte *et al.*, 2005). Lakes were likely in the relatively deep (60 m) and closed northern lagoon (Collot *et al.*, 1988), as well as in the southern lagoon as suggested by 25-35 m deep closed basins in front of the Pirogue river and east of the Mato Pass. The production of maps at different times during the last cycles, especially during the last glaciation/deglaciation period (last 125 ka), taking into account subsidence, sedimentation and detailed bathymetry of the lagoon, barrier reef and passes may be useful to locate possible refuges through Late Pleistocene for the species of the lagoon.

Acknowledgments

I am thankful to Guy Cabioch and Dominique Chardon, both at IRD Nouméa, for fruitful discussions and reviews of the content of the paper.

REFERENCES

- AITCHISON J., CLARKE G., MEFFRE S. & CLUZEL D., 1995. Eocene arc-continent collision in New Caledonia and implications for regional southwest Pacific tectonic evolution. *Geology*, 23(2): 161-164.
- AITCHISON J.C., IRELAND T.R., CLARKE G.L., CLUZEL D., & MEFFRE S., 1998. U/Pb SHRIMP age constraints on the tectonic evolution of New Caledonia and regional implications. *Tectonophysics* 299: 333-343.
- ANDREWS J.E., PACKHAM G. *et al.*, 1975. Sites 285 and 286, in *Initial Reports of the Deep Sea Drilling Project*, Washington, D.C., U.S. Government Printing Office, 30: 27-131.
- AUZENDE J.M., LAFOY Y. & MARSSET B., 1988. Recent geodynamic evolution of the North Fiji Basin (SW Pacific), *Geology*, 16: 925-929.
- AUZENDE J.-M., PELLETIER B. & EISSEN J.-P., 1995. The North Fiji Basin, geology, structure and geodynamic evolution. In: Brian Taylor (Ed), *Backarc Basins: Tectonics and Magmatism*, New York, Plenum Press, 139-175.
- AUZENDE J.M., VAN DE BEUQUE S., REGNIER M., LAFOY Y. & SYMONDS P., 2000. Origin of the New Caledonian ophiolites based on a French-Australian Seismic Transect. *Marine Geology*, 162: 225-236.
- AUZENDE J. M., VAN DE BEUQUE S., DICKENS G., FRANÇOIS C., LAFOY Y., VOUTAY O. & EXON N., 2000, Deep sea diapirs and bottom simulating reflector in Fairway Basin (SW Pacific), *Mar. Geophys. Res.* 21: 579-587.
- AVIAS J., 1967. Overthrust structure of the main ultrabasic New Caledonia massives. *Tectonophysics*, 4(4-6): 531-541.
- BALDWIN S.L., RAWLINGS T., & FITZGERALD P.G., 1999. Thermochronology of the northern high P/T terrane of New Caledonia: implications for mid-Tertiary plate boundary processes in the SW Pacific. In : Baldwin, L., Lister, G.S. (Eds.), Penrose Conference. *Mid-Cretaceous to recent plate boundary processes in the Southwest Pacific*, Abstr Vol., p. 13.
- BAUDRON J.C., GUILLON J.H. & RÉCY J., 1976. Géochronologie par la méthode K/Ar du substrat volcanique de l'île de Maré, archipel des Loyauté (Sud-Ouest Pacifique). *Bulletin BRGM*, 3 série 2: 165-176.
- BITOUN G., RÉCY J., 1982. Origine et évolution du bassin des Loyauté et de ses bordures après la mise en place de la série ophiolitique de Nouvelle-Calédonie, In : Equipe de Géologie-Géophysique ORSTOM Nouméa (Ed), Contribution à l'étude Géodynamique du Sud-Ouest Pacifique, *Trav. Doc. ORSTOM*, 147: 505 - 539.
- BLACK P.M., MAURIZOT P., GHENT E.D., & STOUT M.Z., 1993. Mg-Fe carpholites from aluminous schists in the Diahot region and implications for preservation of high-pressure low-temperature schist, northern New Caledonia. *J. Metamorph. Geol.* 11, 455- 460.
- BLAKE M.C.J., BROTHERS R.N. & LANPHERE M.A., 1977. Radiometric ages of blueschists in New Caledonia. In: Technip (Ed). *International Symposium on Geodynamics in the South-West Pacific, Nouméa 1976*, Paris, 279-282.
- BOURROUILH-LE JAN F., 1996. Plats-formes carbonatées et atolls du centre et sud Pacifique : Stratigraphie, sédimentologie, minéralogie et géochimie. Diagenèse et émergences : Aragonite, calcite, dolomite, bauxite et phosphate. *Document du BRGM*, 249, 365 pp.
- BROCHER T.M., 1985. On the formation of the Vitiav Trench lineament and North Fiji Basin. In: T.M. Brocher, ed., «Investigations of the Northern Melanesian Borderland», *Circum-Pacific Council for Energy and Mineral Resources, Earth Science Series*, 3, Houston, Texas, 13-34.
- BROTHERS R.N., 1974. High-pressure schists in northern New Caledonia. *Contrib. Mineral. Petrol.* 46 (2): 109-127.
- BURNS R.E., ANDREWS J.E. *et al.*, 1973. *Initial Reports of the Deep Sea Drilling Project*, Washington, D.C., U.S. Government Printing Office, 21, 931 pp.
- CABIOCH G., 1988 - *Récifs frangeants de Nouvelle-Calédonie (Pacifique sud-ouest). Structure interne et influences de l'eustatisme et de la néotectonique*. - Thèse Doct. Univ. Provence, Publ. Univ. Aix-MarseilleI, 291p.
- CABIOCH G., RÉCY J., JOUANNIC C. & TURPIN L., 1996. Contrôle climatique et tectonique de l'édification récifale en Nouvelle-Calédonie au cours du Quaternaire terminal. *Bull. Soc. géol. France*, 167(6): 729 - 742.
- CALMANT S., LEBELLEGARD P., TAYLOR F.W., BEVIS M., MAILLARD D., RECY J. & BONNEAU J., 1995. Geodetic measurements of convergence across the New Hebrides subduction zone, *Geophys. Res. Lett.*, 22: 2573-2576.
- CALMANT S., PELLETIER B., BEVIS M., TAYLOR F., LEBELLEGARD P., & PHILLIPS D., 2003 - New insight on the tectonics of the New Hebrides subduction zone based on GPS results. *J. Geophys. Res.*, 108, B6, 2316.
- CAMPBELL H.J., GRANT-MACKIE J.A., & PARIS J.P., 1985. Geology of the Moindou-Téremba area, New Caledonia. Stratigraphy and structure of the Téremba Group (Permian-Lower Triassic) and Baie de St. Vincent Group (Upper Triassic -Lower Jurassic). *Géologie de la France Paris, Bureau des Recherches Géologiques et Minières* 1: 19-36.
- CARNEY J.N. & MACFARLANE A., 1982. Geological evidence bearing on the Miocene to Recent structural evolution of the New Hebrides arc, *Tectonophysics*, 87: 147-175.
- CARRIÈRE D., 1987. *Sédimentation, diagenèse et cadre géodynamique de l'atoll soulevé de Maré, Nouvelle-Calédonie*, Thèse Dr Sci. nat. : Univ. Paris Sud-Orsay. 369 pp.
- CARSON C.J., CLARKE G.L., & POWELL R., 2000. Hydration of eclogite, Pam Peninsula, New Caledonia. *J. Metamorph. Geol.* 18: 79-90.
- CHASE C.G., 1971. Tectonic history of the Fiji plateau, *Geol. Soc. Am. Bull.*, 82: 3087-3110.

- CHARDON D. & CHEVILLOTTE V., 2006. Morphotectonic evolution of the New Caledonia ridge (Pacific Southwest) from post obduction tectonosedimentary record. *Tectonophysics*, 420: 473-491.
- CHEVALIER J.P., 1968. Géomorphologie de l'île de Maré. Les récifs de l'île de Maré. In: Fond. Singer-Polignac (Ed), *Expédition française sur les récifs carbonatés de la Nouvelle-Calédonie*, Paris, 3: 1-158.
- CHEVILLOTTE V., CHARDON D., BEAUVAIS A., MAURIZOT P. & COLIN F., 2006. Long term tropical morphogenesis of New Caledonia (Southwest Pacific): importance for epeirogeny and climate change. *Geomorphology*, 81, 361-375.
- CHEVILLOTTE V., DOUILLET P., CABIOCH G., LAFOY Y., LAGABRIELLE Y. & MAURIZOT P., 2005. Evolution géomorphologique de l'avant-pays du Sud-Ouest de la Nouvelle-Calédonie durant les derniers cycles glaciaires. *Comptes Rendus Geosciences*, 337(7): 695 - 701
- CLARKE G., AITCHISON J.C., & CLUZEL D., 1997. Eclogites and blueschists of the Pam Peninsula NE New Caledonia: a reappraisal. *J. Metamorph. Petrol.* 38 (7): 843- 876.
- CLUZEL D., AITCHISON J., CLARKE G., MEFFRE S. & PICARD C., 1994. Point de vue sur l'évolution tectonique et géodynamique de la Nouvelle-Calédonie (Pacifique. France). *C. R. Acad. Sci. Paris*, 319: 683 - 690.
- CLUZEL D., CLARKE G., & AITCHISON J.C., 1995. Northern New Caledonia high-pressure metamorphic core complex. From continental subduction to extensional exhumation. *Pacrim'95 Congress, Auckland 11/95, Proceedings* Vol., 129- 134.
- CLUZEL D., 1996. Affinités intra-océaniques des métavolcanites de l'Unité de la Boghen (ex-"anté-Permien" de Nouvelle-Calédonie. Pacifique SW). Conséquences paléogéographiques. *C.R. Acad. Sci. Paris*, 323: 657- 664.
- CLUZEL D., PICARD C., AITCHISON J.C., LAPORTE C., MEFFRE S. & PARAT F., 1997. La Nappe de Poya (ex-formation des basaltes) de Nouvelle-Calédonie (Pacifique Sud-Ouest), un plateau océanique Campanien-Paléocène supérieur obducté à l'Eocène supérieur. *C. R. Acad. Sci. Paris*, 324: 443 - 451.
- CLUZEL D., AITCHISON J.C. & PICARD C., 2001. Tectonic accretion and underplating of mafic terranes in the Late Eocene intraoceanic fore-arc of New Caledonia (Southwest Pacific): geodynamic implications. *Tectonophysics*, 340: 23-59.
- CLUZEL D. & MEFFRE S., 2002. L'unité de la Boghen (Nouvelle-Calédonie, Pacifique sud-ouest) : un complexe d'accrétion jurassique. Données radiochronologiques préliminaires U-Pb sur les zircons détritiques. *Comptes Rendus Geosciences*, 334: 867-874.
- CLUZEL D., BOSCH D., PAQUETTE J.L., LEMENNICIER Y., MONTJOIE P. & MÉNOT R.P., 2005. Late Oligocene post-obduction granitoids of New Caledonia: A case for reactivated subduction and slab break-off. *Island Arc*, 14: 254 - 271.
- COLLOT J.Y., MALAHOFF A., RECY J., LATHAM G. & MISSÈGUE F., 1987. Overthrust emplacement of New Caledonia ophiolite: geophysical evidence. *Tectonics*, 6(3): 215 - 232.
- COLLOT J.Y., RIGOLOTT P. & MISSEGUE F., 1988. Geologic structure of the Northern New Caledonia ridges, as inferred from magnetic and gravity anomalies, *Tectonics*, 7 (5): 991-1013.
- COLLOT J.Y., MISSEGUE F., EISSEN J.P., & RIGOLOTT P., 1988 - Carte bathymétrique de la région du Grand Lagon Nord de la Nouvelle-Calédonie et des récifs d'Entrecasteaux. Carte au 1/500000, édition ORSTOM 1988.
- COUDRAY J., 1976 - Recherches sur le Néogène et le Quaternaire marins de la Nouvelle-Calédonie. Contribution de l'étude sédimentologique à la connaissance de l'histoire géologique post-éocène de la Nelle-Calédonie. - Thèse doc. d'Etat, Université de Montpellier In: *Expéd. française sur les récifs coralliens de Nelle-Calédonie*, Fond. Singer-Polignac, Paris, 8: 1-276.
- CRAWFORD A.J., MEFFRE S. & SYMONDS P.A., 2003. 120 to 0 Ma tectonic evolution of the southwest Pacific and analogous geological evolution of the 600 to 220 Ma Tasman Fold Belt System. *Geological Society of Australia Special Publication*, 22: 377 - 397.
- DANIEL J., DUGAS F., DUPONT J., JOUANNIC C., LAUNAY J., MONZIER M. & RÉCY J., 1976. La zone charnière Nouvelle-Calédonie - Ride de Norfolk (S.W. Pacifique) - Résultats de dragages et interprétations. *Cah. ORSTOM, sér. Géol.*, VIII(1): 95-105.
- DUBOIS J., RAVENNE C., AUBERTIN A., LOUIS J., GUILLAUME R., LAUNAY J. & MONTADERT L., 1974. Continental margin near New Caledonia. In: C.A. Burck and C.L. Drake (Ed), *The geology of Continental margins*, New York, Springer-Verlag. 521-535.
- DUBOIS J., LAUNAY J. & RÉCY J., 1974. Uplift movements in New Caledonia-Loyalty Islands area and their plate tectonics interpretation. *Tectonophysics*, 24: 133-150.
- DUBOIS J., LAUNAY J., RÉCY J. & MARSHALL J.F., 1977. New Hebrides trench: subduction rate from associated lithospheric bulge. *Can. J. Earth. Sci.*, 14: 250 - 255.
- DUBOIS J., DEPLU C., DIAMENT M., DANIEL J., & COLLOT J. Y., 1988 - Subduction of the Bougainville Seamount (Vanuatu): mechanical and geodynamic implications, *Tectonophysics*, 149 (1-2): 111-119.
- DUGAS F. & DEBENAY, J.-P., 1978. Interférences des failles-flexures littorales et de l'érosion karstique sur les constructions coralliennes : le lagon de Nouvelle-Calédonie. *C.R. Acad. Sci. Paris, série D*, 290: 963-966.
- DUGAS F., VILLE P., & COUDRAY J., 1980. Etude sismique du lagon sud-ouest de la Nouvelle-Calédonie (Sud-Ouest Pacifique). Paléomorphologies successives et comportement au Quaternaire supérieur du littoral de l'île. *C.R. Acad. Sc. Paris, série D*, 290: 963-966.

- EADE J., 1988. The Norfolk Ridge system and its margins. In: A.E.M. Nairn, F.G. Stehli and S. Uyeda (Ed), *The Ocean Basins and Margins: The Pacific Ocean*, New York, London, Plenum Press, 7B, 303-324.
- EISSEN J.-P., CRAWFORD A.J., COTTEN J., MEFFRE S., BELLON H. & DELAUNE M., 1998. Geochemistry and tectonic significance of basalts in the Poya Terrane, New Caledonia. *Tectonophysics*, 284(3-4): 203 - 219.
- ESPIRAT J.-J., 1963. Etude géologique de régions de la Nouvelle-Calédonie septentrionale (extrémité nord et versant est). Thèse doct. Etat, Toulouse, 217 pp.
- ETHERIDGE M. A., SYMONDS P. A. & LISTER G. S., 1989, Application of the detachment model to reconstruction of conjugate passive margins. In: TANKARD A.J. & BALKWILL H.R. (eds) *Extension Tectonics and Stratigraphy of the North Atlantic Margins*. American Association of Petroleum Geologists Memoirs, 46: 23-40.
- EXON N., HILL P., LAFOY Y., FELLOWS M., PERRY K., MITTS P., HOWE R., CHAPRONIERE G., DICKENS G., USSLER B. & PAULL C., 2004, Geology of the Fairway and New Caledonia Basins in the Tasman Sea from: sediment, pore water, diapirs and bottom simulating reflectors (Franklin Cruise FR9/ 01 and Geoscience Australia Survey 232), *Geoscience Australia Record 2004/26*, 112.
- EXON N., QUILTY P.J., LAFOY Y., & AUZENDE J.M., 2004 – Miocene volcanic seamounts on northern Lord Howe Rise : lithology, age, ferromanganese crusts and origin. *Australian Journal of Earth Sciences*, 51: 291-300.
- FALVEY D.A., 1975. Arc reversals, and a tectonic model for the North Fiji Basin. *Austr. Soc. of Explor. Geophys. Bull.*, 6: 47-49.
- FITZERBERT J.A., CLARKE G.L., BARMO B., & POWELL R., 2004 – The origin and P-T evolution of peridotites and serpentinites of the NE New Caledonia : prograde interaction between continental margin and the mantle wedge. *Journal of Metamorphic Geology*, 22: 327-344.
- FLAMAND B., 2006 – *Les pentes externes du récif-barrière de la Grande-Terre de Nouvelle-Calédonie : morphologie, lithologie, contrôle de la tectonique et de l'eustatisme*. Thèse Université de Bretagne Occidentale, 240 pp + cdrom.
- GAINA C., MULLER R. D., ROYER J.-Y., STOCK J., HARDEBECK J. & SYMONDS P., 1998, The tectonic evolution of the Tasman Sea: A tectonic puzzle with thirteen pieces, *J. Geophys. Res.* 103: 12413–12433.
- GUILLON J.H., 1975. Les massifs péridotitiques de Nouvelle-Calédonie: type d'appareil ultrabasique stratiforme de chaîne récente. *Mém. ORSTOM*, 76: 1-120.
- GUILLON J.H. & ROUTHIER P., 1971. Les stades d'évolution et de mise en place des massifs ultramafiques de Nouvelle-Calédonie. *Cah. O.R.S.T.O.M., sér. Géol.*, IV(2): 5-38.
- GUYOMARD T.S., AISSAOUI D.M., & McNEILL D.F., 1996 – Magnetostratigraphic dating of the uplifted atoll of Mare : geodynamics of the Loyalty Ridge, SW Pacific. *Journal Geophys. Res.* 101: 601-612.
- HALL R., 2002 – Cenozoic geological and plate tectonic evolution of the SE Asia and the SW Pacific ; computer-based reconstructions, model and animation. *Journal of Asian Earth Sciences*, 20: 353-431.
- HAYES D.E. & RINGIS J., 1973. Seafloor spreading in the Tasman Sea, *Nature*, 243: 454-458.
- KELLEY D. *et al.*, 2001 – An off-axis hydrothermal vent field near the Mid-Atlantic Ridge at 30°N. *Nature* 412: 145-149.
- KELLEY D. *et al.*, 2005 – A serpentinite-hosted ecosystem, the Lost City hydrothermal field. *Science*, 307: 1428-1434.
- KROENKE L.W., 1984. Cenozoic tectonic development of the Southwest Pacific. *Committee for co-ordination of joint prospecting for mineral resources in South Pacific offshore areas (CCOP/SOPAC) Tech. Bull.*, New Zealand, 122 pp.
- LAFOY Y., PELLETIER B., AUZENDE J.M., MISSEGUE F. & MOLLARD L., 1994. Tectonique compressive cénozoïque sur les rides de Fairway et Lord Howe, entre Nouvelle-Calédonie et Australie. *C. R. Acad. Sci. Paris*, 319, série IIA: 1063-1069.
- LAFOY Y., MISSEGUE F., CLUZEL D., & LE SUAVE R., 1996 – The Loyalty-New Hebrides arc collision : effects on the Loyalty Ridge and basin system. Southwest Pacific (first results of the ZoNéCo Programme). *Marine Geophys. Res.* 18: 337-356.
- LAFOY Y., AUZENDE J.-M., SMITH R. & LABAILS C., 2000. Evolution géologique post-Pleistocene moyen du domaine lagunaire Néo-Calédonien méridional. *C. R. Acad. Sci. - Series IIA - Earth and Planetary Sciences*, 330(4): 265-272.
- LAFOY Y., BRODIEN R.V. & EXON N.F., 2005. Structure of the basin and ridge system west of New Caledonia (Southwest Pacific): A synthesis. *Marine Geophysical Researches*, 13 pp.
- LAGABRIELLE Y., MAURIZOT P., LAFOY Y., CABIOCH G., PELLETIER B., REGNIER M., WABETE I. & CALMANT S., 2005. Post-Eocene extensional tectonics in Southern New Caledonia (SW Pacific): Insights from onshore fault analysis and offshore seismic data. *Tectonophysics*, 403(1-4): 1 - 28.
- LAPOUILLE A., 1982 – Etude des bassins marginaux fossiles du Sud-Ouest Pacifique : bassin Nord-d'Entrecasteaux, Bassin Nord-Loyauté, Bassin Nord-Fidjien. In : Contribution à l'étude géodynamique du Sud-Ouest Pacifique : *Travaux et Documents de l'ORSTOM*, 147, 409-438.
- LATHAM G., 1986. Altération et pédogenèse sur roches ultrabasiques en Nouvelle-Calédonie. Etudes et Thèses, ORSTOM: 331 pp.
- LAUNAY J., RECY J., 1972 - Variations relatives du niveau de la mer et néotectonique en Nouvelle-Calédonie au Pléistocène supérieur et à l'Holocène. - *Rev. Géogr. phys. Géol. dyn.*, 14 (1): 47-65.
- LAUNAY J., 1985. Paléoniveaux marins et néotectonique à l'île des Pins (Nouvelle-Calédonie). *Bull. Soc. Géol. France*, 1: 77-81.

- LAUNAY J. & FONTES J.-C., 1985 – Les sources thermales de Prony (Nouvelle-Calédonie) et leurs précipités chimiques. Exemple de formation de brucite primaire. *Géologie de la France*, 1, 83-100.
- LEGUÉRÉ J., 1976. Des corrélations entre la tectonique cassante et l'altération supergène des péridotites de Nouvelle-Calédonie. Ph D Thesis, Université des Sciences et Techniques du Languedoc, Montpellier, 80 p.
- LILLIE A.R. & BROTHERS R.N., 1970 – The geology of New Caledonia. *N.Z. Journal of Geology and Geophysics*, 13, 145-184.
- LOUAT R. & PELLETIER B., 1989 - Seismotectonics and present-day relative plate motions in the New Hebrides - North Fiji basin region. *Tectonophysics*, 167: 41-55.
- MAILLET P., MONZIER M., SELO M. & STORZER D., 1983. The d'Entrecasteaux zone (SW Pacific). A petrological and geochronological reappraisal. *Marine Geology*, 53: 179 - 197.
- MAILLET P., MONZIER M., & LEFEVRE C., 1986. Petrology of Matthew and Hunter volcanoes, south New Hebrides island arc (southwest Pacific). *J. Volcanol. Geotherm. Res.*, 30, 1-27.
- MARSHALL J.F. & LAUNAY J., 1978 – Uplift rates of the Loyalty islands as determined by Th/U dating of raised coral terraces. *Quaternary Research*, 9: 186-192.
- MAURIZOT P., EBERLE J.M., HABAUT C. & TESSAROLO C., 1989. Carte géol. Territoires d'Outre-Mer, Nouvelle-Calédonie (1/50000), feuille Pam-Ouégoa, 2e édition, B.R.G.M., Notice explicative par Maurizot P., Eberlé J.M., Habaut C., Tessarolo C., 81 pp.
- McDOUGALL, I. & DUNCAN, R.A., 1988. Age progressive volcanism in the Tasmanid Seamounts. *Earth Planet. Sci. Lett.*, 89: 207-220.
- MEFFRE S., 1995. The development of arc-related ophiolites and sedimentary sequences in New Caledonia, Ph. D. Thesis, University of Sydney, 236 pp.
- MEFFRE S., AITCHISON J.C. & CRAWFORD A.J., 1996. Geochemical stratigraphy of boninites and tholeiites from the Permo-Triassic Koh Ophiolite, New Caledonia. *Tectonics*, 15: 67-83.
- MIGNOT A., 1984. Sismo-stratigraphie de la terminaison Nord de la ride de Lord Howe, Evolution géodynamique du Sud-Ouest Pacifique entre l'Australie et la Nouvelle-Calédonie, Thèse, Université Pierre et Marie Curie, 203 pp.
- MISSÈGUE F. & COLLOT J.-Y., 1987. Etude géophysique du plateau des Chesterfield (Pacifique sud-ouest). Résultats préliminaires de la campagne ZOE 200 du N/O Coriolis. *C. R. Acad. Sci. Paris*, 304, série II, 279-283.
- MISSÈGUE F., *et al.*, 1991 – Carte bathymétrique de la zone économique exclusive de la Nouvelle-Calédonie au 1/1100000. In: MISSÈGUE F., DUPONT J. & DANIEL J., 1992. Carte bathymétrique de synthèse de la zone économique de Nouvelle-Calédonie. projet ZOE500. Rapport de fin d'exécution des travaux, convention ORSTOM / MEDETOM n°2487-00C: 44 pp.
- MITCHELL A.H.G. & WARDEN A.J., 1971. Geological evolution of the New Hebrides island arc. *J. Geol. Soc. London*, 127, 501-529.
- MONZIER M. *et al.*, 1989. Premiers résultats des plongées Nautile de la campagne SUPSOI sur la zone de collision ride des Loyauté arc des Nouvelles Hébrides (sud-ouest Pacifique). *C.R. Acad. Sci. Paris*, 309, II, 2069-2076.
- MONZIER M., DANIEL J. & MAILLET P., 1990. La collision « ride des Loyauté/arc des Nouvelles Hébrides » (Pacifique Sud-Ouest). *Oceanol. Acta*, 10, 43-56.
- MONZIER M., 1993. *Un modèle de collision arc insulaire - ride océanique. Evolution sismo-tectonique et pétrologie des volcanites de la zone d'affrontement arc des Nouvelles-Hébrides - ride des Loyauté.* ORSTOM, Thèse Doct., Université Française du Pacifique. 322 pp.
- MULLER R.D. *et al.* - 2000 Mesozoic/Cenozoic tectonic events around Australia. In : Richards M.A., Gordon R.G., van der Hilst R.D. (Eds), History and dynamics of global Plate motions. *Geophysical Monograph. American Geophysical Union*, 161-188.
- PACKHAM G.H., 1973. A speculative Phanerozoic history of the South-west Pacific. In : P.J. Coleman, ed., *The Western Pacific: Island Arcs, Marginal Seas and Geochemistry*, Nedlands, Western Australia Press, 369-388.
- PARIS J.P., ANDREIEFF P. & COUDRAY J., 1979. Sur l'âge Eocène supérieur de la mise en place de la nappe ophiolitique de Nouvelle-Calédonie, une unité de charriage océanique périaustralien, déduit d'observations nouvelles sur la série de Népoui. *C. R. Acad. Sci. Paris*, 288, série D: 1659 - 1661.
- PARIS J.P., 1981. Géologie de la Nouvelle-Calédonie. *Mém. BRGM*, 133: 1-278.
- PELLETIER B. & AUZENDE J.M., 1996. Geometry and structure of the Vitiaz trench lineament (SW Pacific). *Marine Geophys. Res.*, 18: 305-335.
- PELLETIER B., CALMANT S. & PILLET R., 1998 - Current tectonics of the Tonga-New Hebrides region. *Earth Planet. Sci. Lett.* 164. 263-276.
- PELLETIER B., CHEVILLON C., MENOUE J.-L., BUTSCHER J., FOLCHER E., GEOFFRAY C., BORE J.-M., PANCHÉ J.-Y. & PERRIER J., 2006 - Plongées, forage et cartographie Baie du Prony et Banc Gail, lagon Sud de Nouvelle-Calédonie. N.O. ALIS, 13-17 juin 200 et cartographie Baie du Prony N.O. ALIS, 25-26 septembre 2005. *Rapports de missions, Sci. Terre, Geol-Geophys.* 70, Nouméa IRD septembre 2006, 44 pp.

- PICARD M., 1999 – L'archipel néo-calédonien. 300 millions d'années pour assembler les pièces d'un puzzle géologique. *CTRDP*, 93 pp.
- PILLET R. & PELLETIER B., 2004. Tectonique active, tsunamis et sismicité en Nouvelle-Calédonie. *Notes Techniques, Sciences de la Terre, Géologie-Géophysique*, Centre IRD de Nouméa, 28: 24 pp.
- PONTOISE B., COLLOT J.-Y., MISSÈGUE F. & LATHAM G. V., 1982. Sismique réfraction dans le bassin des Loyauté ; résultats et discussion. In : Contribution à l'étude géodynamique du Sud-Ouest Pacifique : *Travaux et Documents de l'ORSTOM*, 147: 541-548.
- PRINZHOFER A., NICOLAS A., CASSARD D., MOUTTE J., LEBLANC M., PARIS J.P. & RABINOVITCH M., 1980. Structures in the New Caledonia peridotites-gabbros: implications for oceanic mantle and crust. *Tectonophysics*, 69, 85-112.
- PRINZHOFER A., 1981. Structure et pétrologie d'un cortège ophiolitique: le massif du Sud (Nouvelle-Calédonie), *Thèse d'Ing. Doct., ENSM*, Paris, 185 p.
- RAVENNE C., DE BROIN C.E., DUPONT J., LAPOUILLE J. & LAUNAY J., 1977. New Caledonia Basin - Fairway Ridge: structural and sedimentary study. *International Symposium on geodynamics in South West Pacific*, Nouméa (New Caledonia), Editions TECHNIP, Paris: 145-154.
- RAWLING T. J. & LISTER G.S., 2002. Large-scale structure of the eclogite-blueschist belt of New Caledonia. *Journal of Structural Geology*, 24(8): 1239-1258.
- RÉGNIER M., 1988. Lateral variation of upper mantle structure beneath New Caledonia determined from P-wave receiver function: evidence for a fossil subduction zone. *Geophysical Journal of the Royal Astronomical Society*, 95: 561-577.
- RÉGNIER M., VAN DE BEUQUE S., BALDASSARI C. & TRIBOT LASPIÈRE G., 1999. La sismicité du Sud de la Nouvelle-Calédonie : implications structurales. *C. R. Acad. Sci. Paris*, 329: 143-148.
- RIGOLOT P., 1988. Prolongement méridional des grandes structures de Nouvelle-Calédonie et découvertes de monts sous-marins interprétés comme un jalon dans un alignement de hot spot. *C. R. Acad. Sci. Paris*, 307, série II. 965-972.
- RIGOLOT P. & PELLETIER B., 1988. Tectonique compressive récente le long de la marge ouest de la Nouvelle-Calédonie : Résultats de la campagne ZOE 400 du N/O Vauban (mars 1987). *C. R. Acad. Sci. Paris*, 307: 179-184.
- RIGOLOT P., 1989. *Origine et évolution du «système» ride de Nouvelle-Calédonie/Norfolk (Sud-Ouest Pacifique) : Synthèse des données de géologie et de géophysique marine, étude des marges et bassins associés*. Thèse Doct., Université de Bretagne Occidentale. 319 pp.
- ROUTHIER P., 1953. Versant occidental de la Nouvelle-Calédonie entre le col de Boghen et la pointe d'Arama. *Mém. Soc. Géol. Fr.*, 67, 271 pp.
- SCHELLART W.P., LISTER G.S. & TOY V.G., 2006 – A Late Cretaceous and Cenozoic reconstruction of the southwest Pacific region: tectonic controlled by subduction and slab rollback process. *Earth Science Reviews*, 76(3-4): 191-233.
- SDRIOLAS M., MÜLLER R.D. & GAINA C., 2002. Tectonic evolution of the southwest Pacific using constraints from backarc basins. *Geological Society of Australia Special Publication*, 22: 343-359.
- SHACKELTON N.J., 1987 – Oxygen isotopes, ice volumes and sea level. *Quaternary Science Reviews*, 6: 183-190.
- SHACKELTON N.J., 2000 – The 100,000-year ice-age cycle identified and found from lag temperature, carbon dioxide and orbital excentricity. *Science*, 289, 1897-1902.
- SMITH W.H.F., SANDWELL D.T., 1997 – Global sea floor topography from satellite altimetry and ship depth soundings. *Science*, 277, 1956-1962.
- SPANDLER C., RUBATTO D., HERMANN J., 2005 – Late Cretaceous-Tertiary tectonics of the Southwest Pacific: insights from U-Pb sensitive, high-resolution ion microprobe (SHRIMP) dating of eclogite facies rocks from New Caledonia. *Tectonics*, 24(3), TC3003, doi:10.129/2004TCOO1709.
- SUTHERLAND R., 1999. Basement geology and tectonic development of the greater New Zealand region: an interpretation from regional magnetic data. *Tectonophysics* 308: 341-362.
- SUTHERLAND R., KING P., WOOD R., 2001 – tectonic evolution of the Cretaceous rift basins in south-eastern Australia and New Zealand : implications for exploration risk assesment. In Hill K. C., Bernecker T. (Eds.), *Eastern Australasian Basins Symposium 2001*. Petroleum Exploration Society of Australia Special Publication. The Australian Institute of Mining and Metallurgy, Carlton, 3-14.
- SHOR G.G.J., KIRK H.K. & MENARD H.W., 1971. Crustal structure of the Melanesian area, *J. Geophys. Res.*, 76: 2562-2586.
- SYMONDS P., STAGG H., BORISSOVAI., 1999. The transition from rifting and break-up to convergence in the Lord Howe Rise-Norfolk Ridge region. In: Baldwin, L., Lister, G.S. (Eds.), *Penrose Conference, Mid-Cretaceous to recent plate boundary processes in the Southwest Pacific*, Abstr. Vol., 94- 96.
- TRESCASES J. J., 1973. Weathering and geochemical behaviour of the elements of ultramafic rocks in New Caledonia. Metallogenic provinces and mineral deposits in the southwestern Pacific. *Bulletin - Australia. Bureau of Mineral Resources, Geology and Geophysics*, 141: 149-161.
- TRESCASES J.J., 1975 – L'évolution géologique supergène des roches ultrabasiques en zone tropicale. Formation des gisements nickelifères de Nouvelle-Calédonie. *Mém. ORSTOM*, 78: 1-259.

- URUSKI C. & WOOD R., 1991. A new look at the New Caledonia Basin, an extension of the Taranaki Basin, offshore North Island, New Zealand, *Mar. Pet. Geol.*, 8: 379-391.
- VAN DE BEUQUE S., AUZENDE J.-M., LAFOY Y. & MISSEGUE F., 1998. Tectonique et volcanisme tertiaire sur la ride de Lord Howe (Sud-Ouest Pacifique), *C. R. Acad. Sci. Paris*, 326, série II, 663-669.
- VAN DE BEUQUE S., 1999. *Evolution géologique du domaine péri-calédonien (Sud-Ouest Pacifique)*, Thèse Doct., Université de Bretagne Occidentale, Brest, 270 pp.
- VAN DE BEUQUE S., STAGG H.M.J., SAYERS J., WILLCOX J.B. & SYMONDS P.A., 2003 – Geological framework of the northern Lord Howe Rise and adjacent areas. *Geoscience Australia Record* 2003/01, 108 pp.
- VEEVERS J.J., 2000 – Change of tectono-stratigraphic regimes in the Australian plate during the 99 Ma (mid-Cretaceous) and the 43 Ma (mid-Eocene) swerves of the Pacific. *Geology*, 28: 47-50.
- VIALLY R., LAFOY Y., AUZENDE J.M. & FRANCE R., 2003. Petroleum potential of New Caledonia and its Offshore Basins, *American Association of Petroleum Geologists International Conference*, Barcelona, Spain, pp. 1-6.
- VOGT P.R. & CONNOLLY J.R., 1971. Tasmanid guyots, the age of the Tasman basin and motion between the Australia plate and the mantle, *Geol. Soc. Am. Bull.*, 82(9): 2577-2583.
- WEISSEL J.K. & HAYES D.E., 1977. Evolution of the Tasman sea reappraised, *Earth Planet. Sci. Lett.*, 36: 77-84.
- WEISSEL J.K., WATTS A.B. & LAPOUILLE A., 1982. Evidence for late paleocene to late eocene seafloor in the Southern New Hebrides basin, *Tectonophysics*, 87: 243-251.
- WILLCOX J.B., SAYERS J., STAGG H.M.J. & VAN DE BEUQUE S., 2001 – Geological framework of the Lord Howe Rise and adjacent ocean basins. In : Hill K.C., Bernecker T. (Eds.) *Eastern Australasian Basins Symposium. A refocused energy perspective for the future*. Petroleum Exploration Society of Australia, Special Publication, 211-225.
- YAN A. & KROENKE L.W., 1993. A plate tectonic reconstruction of the Southwest Pacific 0-100 Ma. In Berger T., L.W. Kroenke, L. Mayer *et al.*, *Proc. ODP, Sci. Results*, 130, College Station, TX, 697-709.
- YOKOYAMA. K., BROTHERS, R.N. & BLACK, P.M., 1986. Regional facies in the high-pressure metamorphic belt of New Caledonia. *Geol. Soc. Am. Mem.* 184: 407– 423.





Figure 1. Bathymetry of the New Caledonia Exclusive Economic Zone (from : ZoNéCo Programm, 1998)

(a)



(b)




Fig. 2. (a) Topography and bathymetry of the Southwest Pacific region (from Smith and Sandwell (1997)) and (b) regional tectonic setting of (a). Cfz, Cook fracture zone; ChRfsz, Chatham Rise fossil subduction zone; d'ER, d'Entrecasteaux Ridge; EP, East Papua; ER, Efate Re-entrant; LPI, Louisiade Plateau; LoR, Loyalty Ridge; LTr, Louisiade Trough; MaB, Manus Basin; MeR, Melish Rise; NB, New Britain; NBT, New Britain Trench; NCfsz, New Caledonia fossil subduction zone; Nd'EB, North d'Entrecasteaux Basin; NHT, New Hebrides Trench; NLoB, North Loyalty Basin; NST, North Solomon Trough; QT, Queensland Trough; ReB, Reinga Basin, ReR, Reinga Ridge; SCB, Santa Cruz Basin; SCT, San Cristobal Trench; SER, South Efate Re-entrant; SLoB, South Loyalty Basin; SoS, Solomon Sea; SReT, South Rennell Trough; TaB, Taranaki Basin; TKR, Three Kings Ridge; ToT, Townsville Trough; TrT, Trobriand Trough; VMfz, Vening Meinesz fracture zone; WoB, Woodlark Basin; WTP, West Torres Plateau. 1, normal fault; 2, strike-slip fault; 3, subduction zone; 4, spreading ridge (double line) and transform faults (single lines); 5, land; 6-8, sea, with 6, continental or arc crust; 7, oceanic plateau; and 8, basin/ocean floor. Structures in light grey indicate that they are inactive. Thick continuous east-west line at latitude 20° S in panel (a) shows location of cross-section plotted in Fig. 3 h. From Schellart et al., 2006




Figure 3 : East-west cross-sections illustrating the evolution of the Southwest Pacific region since the Late Cretaceous. Line of section is shown on Fig 2a. (modified from Schellart et al, 2006)

DOCUMENTS
SCIENTIFIQUES
et TECHNIQUES

II7

Volume spécial
Deuxième édition

COMPENDIUM
of marine species from
New Caledonia



Edited by
Claude E. PAYRI
Bertrand RICHER DE FORGES



Institut de recherche
pour le développement

CENTRE DE NOUMEA

**DOCUMENTS
SCIENTIFIQUES
et TECHNIQUES**



Institut de recherche
pour le développement

Publication éditée par :
Centre IRD de Nouméa
BP A5, 98848 Nouméa CEDEX
Nouvelle-Calédonie
Téléphone : (687) 26 10 00
Fax : (687) 26 43 26

L'IRD propose des programmes regroupés en 5 départements pluridisciplinaires :

- I DME Département milieux et environnement
- II DRV Département ressources vivantes
- III DSS Département sociétés et santé
- IV DEV Département expertise et valorisation
- V DSF Département du soutien et de la formation des communautés scientifiques du Sud

Modèle de référence bibliographique à cette revue :

Adjeroud M. *et al.*, 2000. Premiers résultats concernant le benthos et les poissons au cours des missions TYPATOLL.
Doc. Sci. Tech. II 3, 125 p.

ISSN 1297-9635. Seconde édition

Numéro II 7 - Octobre 2007

© IRD 2007

Distribué pour le Pacifique par le Centre de Nouméa.



Première de couverture : Récif corallien (Côte Ouest, NC) © IRD.C. Geoffray

Vignettes : voir les planches photographiques

Quatrième de couverture : *Platygyra sinensis* © IRD.C. Geoffray

Matériel de plongée sous-marine © IRD.C. Geoffray	L'Aldric, moyen naviguant de l'IRD © IRD/J.-M. Boré
Récoltes et photographies sous-marines en scaphandre autonome © IRD/J.-L. Menou	Traitement des récoltes en laboratoire © IRD/L. Mattio

CONCEPTION/MAQUETTE/MISE EN PAGE

MAQUETTE DE COUVERTURE

PLANCHES PHOTOGRAPHIQUES

TRAITEMENT DES PHOTOGRAPHIES

JEAN PIERRE MERMOUD/ISABELLE RITZENTHALER

CATHY GEOFFRAY/MINA VILAYLECK

CATHY GEOFFRAY/JEAN-LOUIS MENOÙ/GEORGES BARGIBANT

NOËL GALAUD

La traduction en anglais des textes d'introduction, des Ascidies et des Echinodermes a été assurée par EMMA ROCHELLE-NEWALL.
la préface par MINA VILAYLECK.

Ce document est publié par le Service ISC du Centre IRD de Nouméa et financé par le Ministère de la Recherche et de la Technologie.

ISSN 1297-9635. Seconde édition

Numéro 117 - Octobre 2007

© IRD 2007

**COMPENDIUM OF MARINE SPECIES FROM
NEW CALEDONIA**

SECOND EDITION

Edited by
CLAUDE E. PAYRI, BERTRAND RICHER DE FORGES
