

## New Caledonian seabirds

Jérôme SPAGGIARI<sup>1,2</sup>, Nicolas BARRÉ<sup>1</sup>, Julien BAUDAT-FRANCESCHI<sup>1,3</sup>, Philippe BORSA<sup>1,4</sup>

<sup>1</sup> Société calédonienne d'ornithologie, BP 3135, 98846 Nouméa cedex, New Caledonia

<sup>2</sup> Conservation International, BP 3135, 98846 Nouméa cedex, New Caledonia  
jerome.spaggiari@lagoon.nc

<sup>3</sup> World Wildlife Fund New Caledonia, BP 692, 98845 Nouméa, New Caledonia

<sup>4</sup> Institut de recherche pour le développement, UR 128 Biocomplexité des écosystèmes coralliens, BP A5,  
98848 Nouméa cedex, New Caledonia  
borsa@noumea.ird.nc

### Abstract

Fifty-nine seabird species (among which 28 are confirmed breeders) from 26 genera and 11 families were listed for different regions of New Caledonia (Chesterfield and Bampton atolls, d'Entrecasteaux reefs, Loyalty islands, Grande-Terre / île des Pins, Northern lagoon, Southern lagoon, Walpole island, Matthew island, and Hunter island). A detailed account of systematics, taxonomy, distribution, and conservation status was given for the four breeding species represented by subspecies considered to be endemic to New Caledonia. These include two Procellariidae (New Caledonian Petrel, *Pterodroma leucoptera caledonica*, and the New Caledonian subspecies of Tahiti Petrel, *Pseudobulweria rostrata trouessarti*) and two Laridae (New Caledonian Silver Gull, *Larus novaehollandiae forsteri*, and New Caledonian Fairy Tern, *Sterna nereis exsul*). Two seabird species that breed in New Caledonia are considered by the World Conservation Union as vulnerable (Gould's Petrel, *Pterodroma leucoptera* and White-throated Storm Petrel, *Nesofregetta fuliginosa*) and two others, as near-threatened (Tahiti Petrel and Beach Thick-knee, *Esacus magnirostris*). Several species (Masked Booby, *Sula dactylatra*, Brown Booby, *S. leucogaster*, Fairy Tern, Silver Gull and possibly Herald Petrel, *Pterodroma heraldica*) have been extirpated from regions of New Caledonia where they bred until recently. Drastic conservation measures are urgently needed for the New Caledonian Fairy Tern.

### Résumé

**Oiseaux marins de Nouvelle-Calédonie.** – Cinquante-neuf espèces d'oiseaux marins (dont 28 nicheuses confirmées), appartenant à 26 genres et 11 familles, ont été inventoriées pour différentes régions de la Nouvelle-Calédonie (atolls des Chesterfield et Bampton, récifs d'Entrecasteaux, îles Loyauté, Grande-Terre / Ile des Pins, lagon Nord, lagon Sud, île Walpole, îlot Matthew et îlot Hunter). Une présentation approfondie est faite de la systématique, de la taxinomie, de la distribution, et de l'état de conservation des quatre espèces représentées en Nouvelle-Calédonie par des sous-espèces considérées comme endémiques. Il s'agit de deux Procellariidae (le Pétrel de Nouvelle-Calédonie, *Pterodroma leucoptera caledonica*, et la sous-espèce néo-calédonienne du Pétrel de Tahiti, *Pseudobulweria rostrata trouessarti*), ainsi que deux Laridae (la sous-espèce néo-calédonienne de la Mouette argentée, *Larus novaehollandiae forsteri*, et celle de la Sterne nereis, *Sterna nereis exsul*). Parmi les espèces d'oiseaux marins qui nichent en Nouvelle-Calédonie, deux sont considérées par l'Union mondiale pour la nature comme étant vulnérables (le Pétrel de Gould, *Pterodroma leucoptera* et le Pétrel-tempête à gorge blanche, *Nesofregetta fuliginosa*) et deux autres comme étant quasi-menacées (le Pétrel de Tahiti et l'Oedycnème des récifs, *Esacus magnirostris*). Plusieurs autres espèces (Fou masqué, *Sula dactylatra*, Fou brun, *S. leucogaster*, Sterne nereis, Mouette argentée, ainsi que, peut-être, le Pétrel de Herald, *Pterodroma heraldica*) ont disparu de régions de Nouvelle-Calédonie où elles étaient, jusqu'à récemment, nicheuses. Des mesures drastiques de conservation s'imposent pour la Sterne nereis en Nouvelle-Calédonie.

## Introduction

Visiting naturalists to New Caledonia and surrounding islands have mentioned seabirds as early as 1858–1860, (Bourne *et al.* 2005). Layard and Layard (1882) provided a first list of New Caledonian “waterbirds”, including four Procellariidae [*Oestrelata rostrata* (now *Pseudobulweria rostrata*), *O. mollis* (*Pterodroma mollis*, apparently a misidentification of *P. leucoptera caledonica*), *Adamastor cinereus* (*Procellaria cinerea*), *Puffinus brevicaudus* (*P. tenuirostris*)], one Hydrobatidae [*Oceanites wilsoni* (*O. oceanicus*)], two Phaethontidae [*Phaethon candidus* (*P. lepturus*), *P. rubricauda*], two Sulidae [*Sula piscator* (*S. sula*) and *Dysporus sula* (*S. leucogaster* ?)], one Phalacrocoracidae [*Phalacrocorax melanoleucus* (*P. melanoleucus*)], one Fregatidae [*Tachypetes aquilus* (*Fregata minor*?)], one Burhinidae (*Esacus magnirostris*) and six Laridae [*Larus novaehollandiae*, *Sterna bergii*, *S. melanuchen* (*S. sumatrana*), *S. gracilis* (*S. dougallii*), *Sternula placens* (*Sterna nereis*), *Haliplana fuliginosa* (*S. fuscata*)]. However, it was only recently that the census of populations was completed for the Southern lagoon of New Caledonia’s Grande Terre (Pandolfi-Benoît and Bretagnolle 2002), and initiated in the Northern lagoon, excluding the Belep and Daos archipelagos (Baudat-Franceschi 2006). Our knowledge of seabird populations in the other regions of New Caledonia is still partial.

The objective of this paper was to present an updated list of New Caledonian seabirds by region, with emphasis on endemic taxa. For the latter, we provided details on current distribution, habitats, population sizes, and conservation status.

## Methods

New Caledonia is located in the southwest Pacific Ocean, approximately 1,200 km east of Australia and 1,500 km northwest of New Zealand. The New Caledonian administrative territory extends from approximately 16°52’S to 23°10’S and 157°51’E to 173°57’E (Dubois 1981). The geopolitical definition of New Caledonia thus encompasses a main island, Grande Terre, the Loyalty chain of islands (Beaumtemps–Beaupré, Ouvéa, Lifou, Tiga, Maré, Walpole, all calcareous platforms arisen by tectonic movements) and about a hundred islands and islets scattered from the middle of the Coral sea (the Chesterfield archipelago, comprising low coral sand islets and cays) to the southern New Hebrides ark (Matthew and Hunter islands, both andesitic stratovolcanoes). Grande Terre is by far the largest island, at 350 km in length and between 50 to 70 km wide. A mountain range peaking over 1,600 m runs the length of the island. Surrounding Grande Terre, a barrier reef delimitates a large lagoon, geographically divided into two main lagoons, at the northern and southern extremities of Grande Terre. We compiled all the information that was accessible to us from the literature on the occurrence, taxonomy and distribution of New Caledonian seabirds. This included articles accessible through ISI Web of Science (Institute for Scientific Information, Philadelphia; <http://portal.isiknowledge.com>), the articles and documents listed in the bibliographic catalogue on the marine environment of New Caledonia compiled by Fromaget and Richer de Forges (1992), the articles and other reports listed in Barré and Dutson’s (2000) commented list of New Caledonian birds, unpublished reports by researchers from ORSTOM / IRD archived at the IRD library, Nouméa, unpublished reports by researchers from CIRAD / IAC, Port-Laguerre, and the collection of articles, books, and unpublished reports archived at Société calédonienne d’ornithologie (SCO), Nouméa, which currently is the only birdwatcher association of New Caledonia and BirdLife International affiliate since 2000. Many of the reports presently archived at SCO were collected during the BirdLife Int. project on important bird areas or IBAs in New Caledonia (Spaggiari *et al.* 2006). Additional, unpublished reports were provided by Direction des ressources naturelles of the province Sud government, Nouméa.

The term ‘seabirds’ as it is used here follows Harrison (1995), and includes all species of the families Diomedeidae, Procellariidae, Hydrobatidae, Phaethontidae, Pelecanidae, Sulidae, Phalacrocoracidae, Fregatidae, and Laridae. We added to this list the Osprey (Accipitridae) and the Beach Thick-knee (Burhinidae) as these two species rely mainly, if not exclusively, on marine

resources and nest on the shore or on islets in the lagoon. The taxonomy used in the present list followed Brooke (2004) for Diomedeidae, Procellariidae and Hydrobatidae, and Marchant and Higgins (1990, 1993) and Higgins and Davies (1996) for the other seabird families.

## Results and Discussion

The present paper compiled the data on seabirds from 42 articles in scientific journals, 7 books and 38 other reports, to which a few unpublished observations were added. The updated list of New Caledonian seabirds, by region, is presented in Table 1. Although we may have missed some valuable contributions, we believe that the list presented here is likely to be exhaustive regarding the seabird species currently known to breed in New Caledonia. Fifty-nine seabird species, belonging to 26 genera, from 11 families were thus listed in total. Among them, 28 were confirmed breeders in New Caledonia. We did not include the Grey Petrel, *Procellaria cinerea* in our list of 59, as the location where it was collected was not precise enough ("off the coast, between Noumea and Australia"; Layard and Layard 1882). Neither did we include Beck's Petrel, *Pseudobulweria becki* in that list. Beck's Petrel, which was unrecorded since 1929 and considered as possibly extinct, was recently sighted in the Coral sea (BirdLife International 2006). This observation took place near Cato island, in Australian waters (R. Baxter, in litt.) and not strictly within New Caledonian waters. However, since Beck's Petrel's colonies historically were, and presumably still are located in the Solomon islands, it is likely from R. Baxter's sighting that its zone of foraging includes a wide part of the northern Coral sea and is therefore likely to extend to New Caledonian waters.

Among the species that breed in New Caledonia, two (Gould's Petrel, *Pterodroma leucoptera*, White-throated Storm Petrel *Nesofregetta fuliginosa*) are considered by the World conservation union (IUCN) as vulnerable, and two other species (Tahiti Petrel, *Pseudobulweria rostrata*, Beach Thick-knee, *Esacus magnirostris*), as near-threatened (IUCN 2006). Some other species have been extirpated from regions where they bred until recently: *Sula dactylatra* and *S. leucogaster* signalled as breeders in the southern lagoon (de Naurois and Rancurel 1978) no longer nest there. Idem, *Sterna nereis exsul* on Grande Terre (Layard and Layard 1882). We consider the latter to be the most endangered of all New Caledonian seabirds (see below). The Herald Petrel, *Pterodroma heraldica*, has not been observed on the Chesterfield islands since the collection of two presumed breeding adults, presumably on those islands in 1858-1860 (Bourne *et al.* 2005). The seabird fauna, and the fragile vegetation and soil of the Chesterfield islands were subsequently devastated by whalers and by guano-extraction activities (Bourne *et al.* 2005). This may have extirpated Herald Petrels from the Chesterfield islands. Last, the Silver Gull, *Larus novaehollandiae forsteri*, was said to breed in large numbers on the Huon islands, d'Entrecasteaux reef, at the end of the 19<sup>th</sup> century (Layard and Layard 1882) but subsequent reports have not mentioned it there.

Four of the breeding species are represented by subspecies considered to be endemic, including two Procellariidae (New Caledonian Petrel, *Pterodroma leucoptera caledonica* and Tahiti Petrel, *Pseudobulweria rostrata trouessarti*), and two Laridae (Silver Gull, *Larus novaehollandiae forsteri* and Fairy Tern, *Sterna nereis exsul*). We restricted the following species account to those 4 endemics. This account includes details on their taxonomy and systematics, their distribution and their current conservation status.

Table 2 is a provisional list of the voucher specimens available for a number of species, including two of the four endemics.

### New Caledonian Petrel, *Pterodroma leucoptera caledonica* de Naurois, 1978

Gould's Petrel (*Pterodroma leucoptera*) breeds only in Australia and New Caledonia. The Australian subspecies *P. l. leucoptera* is restricted to two closely distant breeding locations – Cabbage Tree island and Boondelbah island at the entrance to Port Stephens, New South Wales (Marchant and Higgins 1990; Priddel and Carlile 1997; Brooke 2004).

A form of Gould's Petrel breeding in the mountains of New Caledonia was discovered by de Naurois (1978) during his studies of Petrels in New Caledonia. This author first recognized New Caledonian Gould's Petrel as a distinct subspecies on the basis of larger bill and paler plumage on the back, wings and sides of chest (Imber and Jenkins 1981) but eventually changed his mind because he believed that the old Australian *P. leucoptera* specimens he had compared his New Caledonian specimens to were too few to ascertain the distinction, hence the validity of the new subspecies (Imber and Jenkins 1981; Palma and Tennyson 2005). Morphometrics subsequently allowed Imber and Jenkins (1981) to assign specimens of Gould's Petrels washed up on the shores of New Zealand's North island between 1942 and 1980 to the New Caledonian subspecies. The New Caledonian Petrel is clearly distinct from its close Australian relative, hence it was considered a valid subspecies (Imber and Jenkins 1981). These authors considered de Naurois (1978) as the authority for the new subspecies, as "he provided an available name, a breeding locality and some valid characters" although he did not formally describe the subspecies nor designate a type specimen. Imber and Jenkins (1981) nevertheless understood that the specimens studied by de Naurois (1978), now deposited at Museum National d'Histoire Naturelle, Paris, were the types of *P. leucoptera caledonica*, but Palma and Tennyson (2005) recently considered otherwise and designated a specimen from New Caledonia preserved at the American Museum of Natural History as lectotype (Table 2). Palma and Tennyson (2005) also proposed that the authorship of the subspecies be ascribed to Imber and Jenkins (1981), instead of de Naurois (1978) as it is currently cited in major ornithological publications (e.g. Marchant and Higgins 1990; del Hoyo 1992; Dickinson 2003).

This subspecies breeds in New Caledonia and Vanuatu (Tana island: V. Bretagnolle, in Brooke 2004). New Caledonian Petrel breeding sites in New Caledonia are the steep, vegetated slopes of the central chain of mountains at 400–650 m above sea level (de Naurois 1978; V. Bretagnolle, in Brooke 2004). The population size of New Caledonian Petrel is estimated to be on the order of 1,000 – 10,000 pairs (V. Bretagnolle, in Brooke 2004).

#### Tahiti Petrel, *Pseudobulweria rostrata trouessarti* Brasil, 1917

The Tahiti Petrel is widespread in the South West Pacific. Its breeding locations include the Society islands, the Gambier archipelago, the Marquesas islands, Fiji, and New Caledonia (Villard *et al.* 2006, and references therein).

Two subspecies are currently recognized for Tahiti Petrel, namely *Pseudobulweria r. rostrata* and *P. r. trouessarti*. Brasil's (1917) recognition of the New Caledonian form of Tahiti Petrel as a distinct subspecies was later challenged by Murphy and Pennoyer (1952). However, a morphometric analysis of 14 individuals from New Caledonia and 13 from Tahiti confirmed the distinctness of the New Caledonian Tahiti Petrels, on the basis of its heavier bill and longer tarsus (de Naurois and Erard 1979), hence validating Brasil's (1917) taxonomy. Further biometric measurements have confirmed the distinction between *P. r. trouessarti* and *P. r. rostrata* (the other subspecies) from Polynesia (Villard *et al.* 2006). Based on mitochondrial-DNA (cytochrome *b* gene) sequences, Bretagnolle *et al.* (1998) reported that *P. r. trouessarti* differed from Polynesian *P. r. rostrata* by 0.6% nucleotide divergence, which was less than the divergences estimated between nominal species in *Pseudobulweria* and in *Pterodroma*. Additional biometric and genetic data for those and geographically intermediate samples are nevertheless needed to confirm whether *P. r. trouessarti* is a valid subspecies, and whether it eventually would deserve specific rank.

In New Caledonia, the Tahiti Petrel breeds in small scattered colonies on the mountain slopes of Grande-Terre up to 1000 m or more, and on coral and rocky islets in the lagoon (Spaggiari *et al.* 2006; Villard *et al.* 2006, and references therein). Tahiti Petrels are frequently observed from the outer reef to further offshore (Baudat-Franceschi 2006; BirdLife International 2006; Borsa 2006). Individuals are regularly encountered along roads or near villages, up to 45 km from the sea in northern Grande-Terre (Baudat-Franceschi 2006).

The Tahiti Petrel, which digs its burrow under arbustive or forested cover, is threatened by introduced predators (feral pigs and cats, and dogs) and habitat destruction (grazing ungulates, fires and open-cast mining). As those threats occur in many of the islands of the tropical southern Pacific, that is overall almost its whole breeding range, the survival of Tahiti Petrels depends on the long-term preservation of forested terrestrial island ecosystems in the Pacific. The current population size of Tahiti Petrel in New Caledonia is unknown. An estimate of population size has only been provided for the Southern lagoon (ca. 100 pairs; Pandolfi-Benoît and Bretagnolle 2002).

#### New Caledonian Silver Gull, *Larus novaehollandiae forsteri* Mathews, 1912

Three subspecies of Silver Gull have been recognized following the review of Johnstone (1982) : *L. n. novaehollandiae*, which is distributed along the shores of Australia and Tasmania, *L. n. scopulinus*, which occurs in New Zealand and in the Chatham, Snares, Auckland and Campbell islands, and *L. n. forsteri* in New Caledonia (Higgins and Davies 1996). Silver Gulls are occasional visitors to Lord Howe and Norfolk islands, and southern New Guinea. They have also been sighted in Vanuatu (Higgins and Davies 1996).

Silver Gulls from New Caledonia have distinct wing patterns to birds from Queensland (Johnstone 1982), with which they were earlier thought to be of the same subspecies, while recognized as different from those of southern Australia and Tasmania (Dwight 1925, in Higgins and Davies 1996). Although abundant morphometric data are currently available for *L. n. novaehollandiae* and *L. n. scopulinus* (Higgins and Davies 1996), no data was mentioned by these authors for *L. n. forsteri*. More study is therefore needed to clarify the patterns of geographical variation in Silver Gulls and, in particular, to characterize the New Caledonian subspecies. We believe this should be based on morphometrics, coupled with molecular population genetics. Novel microsatellite markers for Silver Gull (Given *et al.* 2002) should prove adequate tools to investigate population structure and gene flow patterns between populations.

New Caledonian Silver Gulls forage on the sandy, coralline or rocky beaches and nearby reefs, beach-rock platforms, mudflats, and sand banks. They also catch small fish swimming at the surface. On land, Silver Gulls are generally attracted to sites of human waste, such as fish landing docks and dumps. They breed on both low vegetated islets and rocky islets of the lagoon around Grande Terre (Pandolfi-Benoît and Bretagnolle 2002; Baudat-Franceschi 2006). As its population size in the Southern lagoon is < 1,500 pairs (Pandolfi-Benoît and Bretagnolle 2002), and that of the Northern lagoon is < 500 pairs (Baudat-Franceschi 2006), the total population size of New Caledonian Silver Gull, non-breeders included, is likely to be no more than within the few thousands.

#### New Caledonian Fairy Tern, *Sterna nereis exsul* Mathews 1912

Three subspecies are currently recognized that differ mainly in size, one each for Australia and Tasmania (*S. n. nereis*), New Zealand (*S. n. davisae*), and New Caledonia (*S. n. exsul*) (Higgins and Davies 1996). The Australian-Tasmanian population of Fairy Tern is the largest, with ca. 3,000-9,000 individuals, and that of New Zealand is the smallest (8 breeding pairs), with intermediate population size for New Caledonia (with, perhaps, ca. 100 breeding pairs) (Brunton and Baling 2005; Baling *et al.* 2006b, and references therein).

Brunton and Baling (2005) produced mitochondrial DNA nucleotide sequences (1,041 base pairs of the ND2 gene) for samples of Fairy Tern from western Australia, southeastern Australia, New Zealand and New Caledonia. A salient feature of the parsimony network of Fairy Tern haplotypes (Fig. 2 of Brunton and Baling 2005) was the long branch (0.5% nucleotide divergence) separating the unique New Caledonian haplotype from that of New Zealand, the latter being in turn separated from a cluster of Australian haplotypes by ca. 0.3% nucleotide divergence. Conversely, populations as far apart as those from western and southeastern Australia shared the same, major haplotype. In other terms, on the basis of the available data, the three Fairy Tern subspecies are geographically isolated

from each other, reaching relatively strong levels of genetic divergence and forming reciprocally monophyletic lineages. To our opinion, specific rank might well be justified for the current subspecies of Fairy Tern. Further genetic studies on Fairy Tern populations including, if possible, the analysis of samples from an intermediate location for which no data are currently available (the Chesterfield and Bampton atolls) may help address that question.

Until recently, the New Caledonian Fairy Tern was known to breed only on small coral islets of the Southern lagoon, including the islets immediately south to Nouméa (Layard and Layard 1878), Kouaré islet (Rancurel 1976) and Redika islet (P.B., unpubl. obs. in 1993: 12 pairs). The breeding population was estimated to be <10 pairs in the mid-1990s (Pandolfi-Benoît and Bretagnolle 2002). More recently, the population size in the Southern lagoon was estimated to be <20 pairs, breeding on 3 islets including Kaé islet and Atiré islet, with a very low reproduction success as checked at the fledgling stage (Brunton and Baling 2005; Baling *et al.* 2006b). Fairy Terns were reported to also breed in the Chesterfield islands (Rancurel 1976; de Naurois et Rancurel 1978). The islets off the northwestern coast of Grande Terre have recently been identified as a key area for the New Caledonian Fairy Tern (N. Baillon, N.B., J.B-F. unpubl.). Overall, the population is considered to be declining, owing to human disturbance at nesting sites (Brunton and Baling 2005; Baling *et al.* 2006b). As the status of Fairy Tern in New Caledonia is now highly critical, drastic steps for its conservation should be taken urgently. Considerable effort has been devoted in New Zealand to achieve similar objectives, not without some success (Ferreira *et al.* 2005).

## REFERENCES

- ANONYMOUS 1996. Un plus pour la faune aviaire du territoire de Nouvelle-Calédonie. *SCO Infos* 9: 3-5.
- BALING M., BRUNTON D.H., JEFFRIES D. (2006a) Marine bird survey in islands of the Southern Lagoon, New Caledonia 4-21 September 2004. *Notornis*, subm.
- BALING M., BRUNTON D.H., JEFFRIES D., BARRÉ N. (2006b) Fairy Tern (*Sterna nereis*) breeding success in southern New Caledonia 2004. *Emu*, subm.
- BARRÉ N., BACHY P. (2003) Complément à la liste commentée des oiseaux de Nouvelle-Calédonie. *Alauda* 71: 31-39.
- BARRÉ N., BAUDAT-FRANCESCHI J., SPAGGIARI J., CHARTENDRAULT V., BACHY P., DESMOULINS F., GUHRING J. (2006a) Second complément à la liste commentée des oiseaux de Nouvelle-Calédonie. *Alauda* 75: 129-144.
- BARRÉ N., DUTSON G. (2000) Oiseaux de Nouvelle Calédonie - Liste commentée. *Alauda* 68: 1-48.
- BARRÉ N., GÉRAUX H. (2004) Great cormorant (*Phalacrocorax carbo*) breeds in New Caledonia. *Notornis* 51: 113-114
- BARRÉ N., VILLARD P., MANCEAU N., MONIMEAU L., MÉNARD C. (2006b) Les oiseaux de l'archipel des Loyauté (Nouvelle-Calédonie) inventaire et éléments d'écologie et de biogéographie. *Rev. Ecol. (Terre Vie)* 61: 45-64.
- BARRITT M.K. (1975) A visit to Hunter and Matthew islands, two little-known islands in the Hunter island ridge, south-east of the New Hebrides chain, by HMS Hydra, surveying ship. *Sea Swallow* 25: 13-15.
- BAUDAT-FRANCESCHI J. (2006) Oiseaux marins côtiers et nicheurs en province Nord. Société calédonienne d'ornithologie, Nouméa. 105 pp.
- BEDIN J.-P. (1996) Compte-rendu de croisière à l'archipel des îles Chesterfield du 16 décembre 1995 au 26 décembre 1995. Association pour la sauvegarde de la nature néo-calédonienne. Nouméa, 3 pp.
- BELL M. (1998) Seabird island restoration: New Caledonia. Eradication of rats. October 1998. Report prepared for Service des parcs et réserves terrestres. Direction des Ressources Naturelles. Nouméa, New Caledonia. Wildlife Management International, Wellington. 12 pp.
- BEUGNET F., COSTA R., FERRÉ O., MARCHAL V. (1993) Statut sanitaire et inventaire de l'avifaune des îlots français du Pacifique Sud - Etude de l'île Surprise. *Rev. Méd. Vét.* 144: 607-613.
- BIRDLIFE INTERNATIONAL. (2000) Threatened birds of the world. Lynx Edicions, Barcelona, 852 pp.
- BIRDLIFE INTERNATIONAL. (2006) Beck's is back! Downloaded from <http://www.birlife.org> on 19/9/2006.
- BORSA P. (2004) Mission ornithologique sur l'îlot Matthew, 10-13 août 2004. Institut de recherche pour le développement, Nouméa, 4 pp.
- BORSA P. (2006) Mission ornithologique aux îles Chesterfield, 12-16 décembre 2005. Institut de recherche pour le développement, Nouméa, 8 pp.
- BORSA P., Boiteux N. (2007) Recensement des oiseaux marins de l'île Longue (atoll des Chesterfield), 18-22 juin 2007. Institut de recherche pour le Développement. Nouméa, 14 pp.

- BOURNE W.R.P. (1966) Observation of seabirds. *Sea Swallow* 18: 9-36.
- BOURNE W.R.P. (1967) Observation of seabirds and review of literature. *Sea Swallow* 19: 51-76.
- BOURNE W.R.P. (1970) Observation of seabirds. *Sea Swallow* 20: 47-52.
- BOURNE W.R.P. (1984) Reports of seabirds received 1973-1977 - Part I. *Sea Swallow* 19
- BOURNE W.R.P., DAVID A.C.F., McALLAN I.A.W. (2005) The birds of the southern Coral sea including observations by HMS Herald in 1858-60. *Atoll Res. Bull.* 541: 239-263.
- BRETAGNOLLE V. (2001) Le pétrel de la chaîne *Pterodroma (leucoptera) caledonica*: statut et menaces. Centre d'études biologiques de Chizé, Beauvoir-sur-Niort, 33 pp.
- BRETAGNOLLE V., ATTIE C., PASQUET E. (1998) Cytochrome b evidence for the validity and phylogenetic relationships of *Pseudobulweria* and *Bulweria* (Procellariidae). *Auk* 115: 188-195
- BRETAGNOLLE V., PANDOLFI-BENOÎT M. (1997) Distribution, status and notes on the breeding biology of the bridled tern *Sterna anaethetus* in New Caledonia. *Emu* 97: 310-315.
- BRETAGNOLLE V., PANDOLFI M., LECOQ V., BROUDISSOU J. (2001) Le balbuzard pêcheur *Pandion haliaetus* en Nouvelle-Calédonie: effectif, répartition et menaces. *Alauda* 69: 491-501.
- BROOKE M. (2004) Albatrosses and petrels across the world. Oxford University Press, New York, 499 pp.
- BRUCE M.D. (1978) L'avifaune de Lifou (archipel des Loyautés). *Alauda* 46: 295-308.
- BRUCE M.D. (1985) (Oiseaux de Nouvelle-Calédonie et des Loyautés) New Caledonian birds. *Emu* 85: 275-276.
- BRUNTON D.H., BALING M. (2005) Conservation genetics of the New Zealand fairy tern (*Sterna nereis daviseae*). Auckland UniServices Limited, Auckland, 32 pp.
- CHAPMAN S.E. (1983) Note on seabird reports received 1981-1982. *Sea Swallow* 32: 12-21.
- COHIC F. (1959) Report on a visit to the Chesterfield Islands, September 1957. *Atoll Res. Bull.* 63: 1-11.
- CONDAMIN M. (1977) Compte-rendu de mission aux îles Chesterfield du 29.IX au 7.X.1977. Office de la recherche scientifique et technique outre-mer, Nouméa, 8 pp.
- CONDAMIN M. (1978) Compte-rendu de mission aux îles Walpole, Hunter et Matthew (6 au 8-XII-1977 : 4-I-1978). Office de la recherche scientifique et technique outre-mer, Nouméa, 8 pp.
- CONDAMIN M. (1978) Redécouverte d'*Esacus magnirostris* (Burhinidae) en Nouvelle-Calédonie. *Oiseau Rev. Fr. Ornithol.* 4: 381-382.
- CONDAMIN M. (1979) Quelques observations sur les Procellariiformes en Nouvelle-Calédonie. *Oiseau Rev. Fr. Ornithol.* 49: 51.
- CONDAMIN M., DE NAUROIS R. (1987) Sur deux espèces de Sterninae observées aux îles Walpole, Hunter et Matthew (Pacifique Sud). *Alauda* 55: 140-146.
- COSTA R., THÉVENON J. (1987) Surveillance sanitaire des oiseaux migrateurs des îles Chesterfield. Rapport de Mission. Direction du développement et de l'économie rurale, Nouméa, 12 pp.
- DELACOUR J. (1966) Guide des oiseaux de la Nouvelle-Calédonie. Delachaux & Nicstlé, Neuchâtel, 173 pp.
- DEL HOYO J., ELLIOTT A., SARGATAL J. (eds.) (1992) Handbook of the birds of the world, vol. 1: Ostrich to Ducks. Lynx Edicions, Barcelone, 696 pp.
- de NAUROIS R. (1978) Procellariidae reproducteurs en Nouvelle-Calédonie pendant l'été austral. *C. R. Acad. Sci. Paris, D* 287: 269-271.
- de NAUROIS R. (1985) Field notes on the New Caledonian raptors with particular reference to *Haliastur sphenurus*. *Bonn. Zool. Beiträge* 36: 65-68.
- de NAUROIS R., ERARD C. (1979) L'identité subspecifique des populations néo-calédoniennes de *Pterodroma rostrata* Peale 1848. *Oiseau Rev. Fr. Ornithol.* 49: 235-239.
- de NAUROIS R., RANCUREL P. (1978) Données nouvelles sur la répartition et l'écologie de quelques espèces aquatiques et d'oiseaux de mer de Nouvelle-Calédonie. *C. R. Acad. Sci. Paris, D* 287: 627-629.
- de NAUROIS R., RANCUREL P. (1978) Observations nouvelles sur les Laridae reproducteurs en Nouvelle-Calédonie. *C. R. Acad. Sci. Paris, D* 287: 495-498.
- DESMOULINS F., BARRÉ N. (2006) L'avifaune des forêts sèches et des milieux avoisinants de Gouaro Déva. Institut agronomique néo-calédonien. Port Laguerre, 7 pp.
- DICKINSON E.C. (ed.) (2003) The Howard and Moore complete checklist of the birds of the world, 3<sup>rd</sup> edn. Christopher Helm, London, 1040 pp.
- DUBOIS J.-P. (1981) Le cadre géopolitique. In: Sautter G. (ed.) Atlas de la Nouvelle-Calédonie et dépendances. Office de la recherche scientifique et technique outre-mer, Paris, pl. 1.
- FERREIRA S.M., HANSEN K.M., PARRISH G.R., PIERCE R.J., PULHAM G.A., TAYLOR S. (2005) Conservation of the endangered New Zealand fairy tern. *Biol. Conserv.* 125: 345-354.
- FROMAGET M., RICHER de FORGES B. (1992) Catalogue bibliographique indexé du milieu marin de Nouvelle Calédonie. 2<sup>e</sup> édition. Office de la recherche scientifique et technique outre-mer. Nouméa, 274 pp.
- GIBSON J.D. (1960) Seabird log - Sydney to Cape Town and Panama to Sydney. *Emu* 60: 11-19.

- GIVEN A.D., MILLS J.A., BAKER A.J. (2002) Isolation of polymorphic microsatellite loci from the red-billed gull (*Larus novaehollandiae scopulinus*) and amplification in related species. *Mol. Ecol. Notes* 2: 416-418;
- GODARD P. (1982) L'île la plus proche du paradis. Editions d'art calédoniennes, Nouméa, 238 pp.
- GRAY G.R. (1859) Catalogue of the birds of the tropical islands of the Pacific ocean in the collection of the British Museum. Taylor and Francis, London.
- HAMEL P. (1993) Campagne d'observations et de marquage des tortues marines aux îles Surprise, Le Leizour et Huon (récif d'Entrecasteaux). Office de la recherche scientifique et technique outre-mer, Nouméa, 17 pp.
- HANNECART F. (1988) Les oiseaux menacés de la Nouvelle-Calédonie et des îles proches. In: Thibault J.-C., Guyot I. (eds.) Livre rouge des oiseaux menacés des régions françaises d'outre-mer. Monogr. 5. Conseil international pour la protection des oiseaux, Cambridge, pp. 43-165.
- HANNECART F., LAPLAGNE M. (1969) Les Chesterfield - Surprise (27 septembre - 8 octobre 1969). Société calédonienne d'ornithologie, Nouméa, 7 pp..
- HANNECART F., LÉTOCART Y. (1980) Oiseaux de l'Île-Calédonie et des Loyautés - tome 1. Cardinalis, Nouméa, 150 pp.
- HANNECART F., LÉTOCART Y. (1983) Oiseaux de l'Île-Calédonie et des Loyautés - tome 2. Cardinalis, Nouméa, 134 pp
- HARRISON P. (1995) Oiseaux de mer. Broquet, Ottawa, 448 pp.
- HIGGINS P.J., DAVIES S.J.J.F. (eds.) (1996) Handbook of Australian, New Zealand and Antarctic Birds, vol. 3, Snipes to Pigeons. Oxford University Press, Melbourne, 1028 pp.
- IMBER M.J., JENKINS J.A.F. (1981) The New Caledonian petrel. *Notornis* 28: 149-160.
- IUCN (2006) 2006 IUCN red list of threatened species. Downloaded from <http://www.iucnredlist.org> on 26th September 2006.
- JESPERSEN P. (1933) Observations of oceanic birds of the Pacific and the adjacent waters. *Videnskabelige Meddelelser fra Dansk Naturhistorisk Forening i København* 94: 187-221.
- JOHNSTONE R.E. (1982) Distribution, status and variation of the silver gull *Larus novaehollandiae* Stephens, with notes on the *Larus cirrocephalus* species-group. *Rec. West. Aus. Mus.* 10: 133-165.
- KUSSER J. (1986) L'archipel des Chesterfield. Direction du développement de l'économie rurale, Nouméa, 12 pp.
- KUSSER J., SUPRIN B. (1990) Rapport de mission aux Chesterfield. Direction du développement de l'économie rurale, Nouméa, 15 pp.
- LAMBERT C. (1987) Surveillance sanitaire des oiseaux migrateurs des îles Chesterfield. Direction du développement de l'économie rurale, Nouméa, 12 pp.
- LAYARD E.L., LAYARD E.L.C. (1878a) Notes on the avifauna of New Caledonia. *Ibis* 20: 250-267.
- LAYARD E.L., LAYARDE L. C. (1880) Notes on the avifauna of the Loyalty islands. *Ibis* 4: 220-234.
- LAYARD E.L., LAYARD E.L.C. (1882) Notes on the avifauna of the New Caledonia. A catalogue of the birds of the island known to E.L. & E.L.C. Layard. With remarks by the Reverend Canon Tristram. *Ibis* 6, 493-546.
- MacDONALD J.D., LAWFORD P.A. (1954) Sight records of birds in the Pacific: compiled from the bird log during the recent cruises of HMS Challenger. *Emu* 54, 7-28.
- MacMILLAN L. (1938) Notes sur les oiseaux des îles Loyautés. *Bull. Soc. Etudes Mélanésiennes* 1: 22-26.
- MANCEAU N. & BARRÉ N. (2001) Inventaire et répartition des oiseaux de Lifou. Principales zones d'intérêt patrimonial. Institut agronomique néo-calédonien, Port Laguerre, 30 pp.
- MARCHANT S., HIGGINS P.J. 1990 (Eds). Handbook of Australian, New Zealand and Antarctic birds. vol. 1, Ratites to Ducks. Oxford University Press, Melbourne, 1400 pp.
- MARCHANT S., HIGGINS P.J. 1993 (Eds). Handbook of Australian, New Zealand and Antarctic birds, vol. 2, Raptors to Lapwings. Oxford University Press, Melbourne, 984 pp.
- MAYR E. (1945) Birds of the southwest Pacific. MacMillan. New York, 316 pp.
- MEETH P., MEETH K. (1983) Seabird observations from six Pacific ocean crossings. *Sea Swallow* 32. 56-65.
- MURPHY R.C., PENNOYER J.M. (1952) Larger petrels of the genus *Pterodroma*. *Am. Mus. Novitates* 1580: 1-43.
- PALMA R.L., TENNYSON A.J. (2005) Designation of a lectotype and clarification of authorship and date of publication for the New Caledonian petrel (*Pterodroma leucoptera caledonica*). *Notornis* 52: 247-248.
- PANDOLFI-BENOÎT M. (1993a) Mission ornithologique à l'île Loop (février 1993). Direction du développement rural, Province Sud, Nouméa, 16 pp.
- PANDOLFI-BENOÎT M. (1993b) Mission ornithologique aux îles Walpole et Matthew (juillet 1993). Direction du développement rural, Province Sud, Nouméa, 16 pp.
- PANDOLFI-BENOÎT M. (1993c) Rapport de la mission ornithologique. Direction du Développement Rural, Province Sud, Nouméa, 28 pp.
- PANDOLFI-BENOÎT M., BRETAGNOLLE V. (2002) Seabirds of the southern lagoon of New-Caledonia: distribution, abundance and threats. *Waterbirds* 25: 202-213.
- PRIDDEL D., CARLILE N. (1997) Boondelbah island confirmed as a second breeding locality for Gould's petrel *Pterodroma leucoptera leucoptera*. *Emu* 97: 245-248.

- RANCUREL P. (1973a) Compte-rendu de mission aux îles Chesterfield du 21 au 28 juin 1973. ORSTOM, Nouméa, 8 pp.
- RANCUREL P. (1973b) Compte-rendu d'une visite aux îles Hunter-Matthew-Walpole du 16 au 22 décembre 1973. ORSTOM, Nouméa, 14 pp.
- RANCUREL P. (1974) Compte-rendu d'une visite à l'île Surprise le 31 janvier 1974. ORSTOM, Nouméa, 6 pp.
- RANCUREL P. (1976) Liste préliminaire des oiseaux de mer des îles et îlots voisins de la Nouvelle-Calédonie. *Cah. O.R.S.T.O.M., Océanogr.* 14: 163-168.
- ROBINET O., CRAIG J.L., & CHARDONNET L. (1998) Impact of rat species in Ouvéa and Lifou (Loyalty islands) and their consequences for conserving the endangered Ouvea parakeet. *Biol. Conserv.* 86: 223-232.
- ROBINET O., SIRGOUANT S., & BRETAGNOLLE V. (1997) Marine birds of d'Entrecasteaux reefs (New Caledonia, Southwestern Pacific): diversity, abundance, trends and threats. *Colon. Waterbirds* 20: 282-290.
- SALVIN O. (1888) Critical notes on the Procellariidae. *Ibis* 30: 411-415.
- SCO (1996) Rapport de mission aux récifs d'Entrecasteaux du 17 au 30 novembre 1996. Société calédonienne d'ornithologie, Nouméa, 13 pp.
- SIRGOUANT S. (1994) Mission ornithologique sur l'île de Walpole (du 28 novembre 1993 au 4 décembre 1993). Société Calédonienne d'Ornithologie, Nouméa, 8 pp.
- SPAGGIARI J., & BARRÉ N. (2003) Dénombrement des puffins du Pacifique (*Puffinus pacificus chlororhynchus*) nichant dans la colonie de la presqu'île de Pindai. Société calédonienne d'ornithologie, Nouméa, 23 pp.
- SPAGGIARI J., & BARRÉ N. (2004) Inventaire complémentaire des sites de nidification du pétrel de Tahiti *Pseudobulweria rostrata trouessarti* sur le massif du Koniambo. Société calédonienne d'ornithologie, Nouméa, 28 pp.
- SPAGGIARI J., & BARRÉ N. (2005) Dénombrement des puffins fouquets (*Puffinus pacificus chlororhynchus*) nichant dans la colonie de la presqu'île de Pindai. Société calédonienne d'ornithologie, Nouméa, 19 pp.
- SPAGGIARI J., CHARTENDRAULT V., & BARRÉ N. (2006) Zones importantes pour la conservation des oiseaux de Nouvelle-Calédonie. Société calédonienne d'ornithologie, Nouméa, 204 pp.
- SULLIVAN C. R. (1928) Bird notes from Walpole Island. *Eruu* 28: 136-137.
- VILLARD P. (2001) Inventaire et répartition des oiseaux de Tiga. Institut agronomique néo-calédonien, Port Laguerre, 14 pp.
- VILLARD P. (2002a) Inventaire et répartition des oiseaux de Maré. Principales zones d'intérêt patrimonial. Institut agronomique néo-calédonien, Port Laguerre, 25 pp.
- VILLARD P. (2002b) Inventaire et répartition des oiseaux d'Ouvéa. Principales zones d'intérêt patrimonial. Institut agronomique néo-calédonien, Port Laguerre, 27 pp.
- VILLARD P. (2003) Rapport intermédiaire d'étude pour le compte de la Province Sud, Nouvelle-Calédonie. Centre d'études biologiques de Chizé, Beauvoir-sur-Niort, 35 pp.
- VILLARD P., & BARRÉ N. (2002) Inventaire et statut des oiseaux dans la zone d'emprise du projet Koniambo (Province Nord, Nouvelle-Calédonie) pp. 56. Institut agronomique néo-calédonien, Port Laguerre, 56 pp.
- VILLARD P., DANO S., & BRETAGNOLLE V. (2006) Morphometrics and the breeding biology of the Tahiti Petrel *Pseudobulweria rostrata*. *Ibis* 148: 285-291
- WARNER D.W. (1947) The ornithology of New Caledonia and the Loyalty Islands. Cornell University Press, Ithaca, 232 pp.

**Table 1** Updated list of New Caledonian seabirds, by region. *Chest.* Chesterfield- Bampton islands; *d'Entr.* d'Entrecasteaux reef; *Loyalty* Loyalty islands; *Gde Terre* Grande Terre and Isle of Pines; *N. Lag.* Northern lagoon; *S. Lag.* Southern lagoon; *Walpole* Walpole island; *Matthew* Matthew island; *Hunter* Hunter island; *o* observed; breeding status: *b* confirmed breeder; *l* likely breeder. *At sea* at-sea records: *o* records for which references were provided (for all non-breeding and for some rarely observed species); + many records, for which references were not provided. References: 1. Bourne (1967); 2. Delacour (1966); 3. Hannecart and Létocart (1983); 4. MacDonald and Lawford (1954); 5. Warner (1947); 6. J.S., unpubl. obs. in 2006; 7. Anonymous (1996); 8. Barré *et al.* (2006a); 9. Chapman (1983); 10. Rancurel (1976); 11. P.B., unpubl. obs. in 2002; 12. Layard and Layard (1882); 13. Barré and Dutson (2000); 14. Barré *et al.* (2006b); 15. Baudat-Franceschi (2006); 16. Bedin (1996); 17. Bell (1998); 18. Beugnet *et al.* (1993); 19. Bourne *et al.* (2005); 20. Condamin (1977); 21. Condamin (1978); 22. Condamin (1979); 23. de Naurois and Rancurel (1978); 24. Desmoulin and Barré (2006); 25. Kusser and Suprin (1990); 26. Lambert (1987); 27. Pandolfi-Benoît (1993a); 28. Pandolfi-Benoît (1993c); 29. Pandolfi-Benoît and Bretagnolle (2002); 30. Rancurel (1973a); 31. Rancurel (1973b); 32. Robinet, Craig and Chardonnet (1998); 33. Robinet *et al.* (1997); 34. SCO (1996); 35. Spaggiari and Barré (2003); 36. Spaggiari and Barré (2005); 37. Villard (2002a); 38. Villard (2002b); 39. Villard (2003); 40. T. Walker and F. Savage, in Bourne *et al.* (2005); 41. Bourne (1984); 42. Meeth and Meeth (1983); 43. Layard and Layard (1878a); 44. Gray (1859); 45. Mayr (1945); 46. Bourne (1966); 47. Jespersen (1933); 48. Borsa (2004); 49. de Naurois (1978); 50. de Naurois and Erard (1979); 51. Hannecart (1988); 52. Spaggiari and Barré (2004); 53. Villard and Barré (2002); 54. Villard *et al.* (2006); 55. obs. by F. Hannecart in 1981 (pers comm.); 56. Salvin (1888); 57. Spaggiari *et al.* (2006); 58. Gibson (1960); 59. P.B., unpubl. obs. in 2004; 60. Bretagnolle (2001); 61. Imber and Jenkins (1981); 62. J.S., unpubl. obs. in 2005; 63 Bourne (1970); 64. Barritt (1975); 65. de Naurois and Rancurel (1978); 66. Sullivan (1928); 67. Condamin and de Naurois (1987); 68. Godard (1982); 69. Layard and Layard (1880); 70. Pandolfi-Benoît (1993b); 71. Sirgouant (1994); 72. Villard (2001); 73. Borsa (2006); 74. Cohic (1959); 75. Costa and Thévenon (1987); 76. Hamel (1993); 77. Hannecart and Laplagne (1969); 78. Kusser (1986); 79. Rancurel (1974); 80. Barré and Géraux (2004); 81. Hannecart and Létocart (1980); 82. Manceau and Barré (2001); 83. Bretagnolle *et al.* (2001); 84. Bruce (1978); 85. Bruce (1985); 86. Condamin (1978); 87. obs. by G. Dutson in 2003 (pers comm.); 88. Barré and Bachy (2003); 89. Baling *et al.* (2006b); 90. Bretagnolle and Pandolfi-Benoît (1997); 91. de Naurois (1985); 92. MacMillan (1938); 93. J.B.-F., unpubl. obs. in 2006; 94. N.B.. unpubl. obs. in 2005; Borsa and Boiteux (2007).

Table 1

Family, Species	Vernacular name	Region of New Caledonia								At sea	References
		Chest..	d'Entr.	Loyalty	Gde Terre	N. Lag	S. Lag	Walpole	Matthew	Hunter	
<b>Diomedeidae</b>											
<i>Diomedea exulans</i>	Wandering Albatross									o	1, 2, 3, 4, 5
<i>Diomedea antipodensis</i>	Antipodean Albatross										6
<i>Diomedea epomophora</i>	Southern Royal Albatross										7
<i>Thalassarche melanophris</i>	Black-browed Albatross									o	2, 3, 5
<b>Procellariidae</b>											
<i>Macronectes giganteus</i>	Southern Giant Petrel										3
<i>Macronectes halli</i>	Northern Giant Petrel										8
<i>Daption capense</i>	Cape Petrel									o	9, 10, 11
<i>Procellaria cinerea</i>	Grey Petrel									o	12
<i>Calonectris leucomelas</i>	Streaked Shearwater									o	13
<i>Puffinus pacificus chlororhynchos</i> <sup>1</sup>	Wedge-tailed Shearwater	b	b	b	b	b	o	o	b	+	10, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40
<i>Puffinus bulleri</i>	New Zealand Shearwater									o	95
<i>Puffinus carneipes</i>	Flesh-footed Shearwater									o	11, 41, 95
<i>Puffinus griseus</i>	Sooty Shearwater									o	4, 13, 42
<i>Puffinus tenuirostris</i>	Short-tailed Shearwater						o			o	5, 13, 15, 42, 43, 93
<i>Puffinus gavia</i>	Fluttering Shearwater									o	2, 4, 43, 44, 45
<i>Puffinus lherminieri gunnisi</i>	Audubon's Shearwater									o	2, 8, 45
<i>Puffinus assimilis</i>	Little Shearwater									o	46, 47
<i>Pseudobulweria rostrata troussartii</i>	Tahiti Petrel		o	b	b	b		o	o	+	2, 12, 14, 15, 17, 29, 48, 49, 50, 51, 52, 53, 54, 55, 95
<i>Pterodroma neglecta</i>	Kermadec Petrel									o	59
<i>Pterodroma heraldica</i> <sup>2</sup>	Herald Petrel				1				b	o	1, 8, 15, 19, 45, 56, 57
<i>Pterodroma solandri</i>	Providence Petrel				o					o	58, 59
<i>Pterodroma inexpectata</i>	Mottled Petrel									o	1, 13
<i>Pterodroma brevipes</i>	Collared Petrel									o	8
<i>Pterodroma leptoptera caledonica</i>	New Caledonian Petrel		b		o	o	1			+	10, 45, 49, 51, 60, 61
<i>Pterodroma cookii</i>	Cook's Petrel									o	11, 42
<i>Pterodroma cervicalis</i>	White-necked Petrel									o	42
<i>Pterodroma nigripennis</i>	Black-winged Petrel		o		1	b		o	1	+	15, 21, 22, 29, 42, 51, 55, 61

**Table 1** (continued)

Family, Species	Vernacular name	Region of New Caledonia								At sea	References
		Chest.	d'Entr.	Loyalty	Gde Terre	N. Lag.	S. Lag.	Walpole	Matthew		
<b>Hydrobatidae</b>											
<i>Oceanites oceanicus</i>	Wilson's Storm Petrel					o				o	3, 12, 15, 43, 45, 62, 93
<i>Pterodroma grallaria</i>	White-bellied Storm Petrel									o	4, 63
<i>Nesofregata fuliginosa</i> <sup>3</sup>	White-throated Storm Petrel									o	13, 64
<b>Phaethontidae</b>											
<i>Phaethon lepturus</i>	White-tailed Tropicbird			l				b	o	o	14, 21, 37, 51, 57, 65, 66
<i>Phaethon rubricauda</i>	Red-tailed Tropicbird	b	b	b		o	b	b	b	o	10, 12, 14, 21, 31, 32, 33, 34, 37, 48, 51, 57, 59, 65, 66, 67, 68, 69, 70, 71, 72
<b>Pelecanidae</b>											
<i>Pelecanus conspicillatus</i>	Australian Pelican				o						3
<b>Sulidae</b>											
<i>Sula serrator</i>	Australasian Gannet					o					13
<i>Sula dactylatra personata</i> <sup>4</sup>	Masked Booby	b	b	b		b	b	b	b	+	10, 14, 15, 16, 19, 21, 25, 26, 27, 28, 30, 33, 34, 40, 48, 51, 57, 65, 72, 73, 74, 75, 76, 95
<i>Sula sula rubripes</i> <sup>5</sup>	Red-footed Booby	b	b	o			b	b	b	+	10, 14, 17, 19, 20, 21, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 37, 39, 40, 57, 65, 70, 71, 73, 74, 76, 77, 78, 79, 95
<i>Sula leucogaster plotis</i> <sup>6</sup>	Brown Booby	b	b	b		b	b	b	b	+	10, 14, 15, 16, 19, 20, 21, 25, 26, 27, 28, 30, 31, 33, 34, 37, 40, 51, 57, 65, 67, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 95
<b>Phalacrocoracidae</b>											
<i>Phalacrocorax carbo</i>	Great Cormorant				b						13, 80
<i>Phalacrocorax sulcirostris</i>	Little Black Cormorant	o									13, 20, 68, 80, 81
<i>Phalacrocorax melanoleucus melanoleucus</i>	Little Pied Cormorant		o		b					+	13, 28, 68, 80, 81
<b>Fregatidae</b>											
<i>Fregata ariel ariel</i>	Lesser Frigatebird	b	b	o		o		b	b	+	10, 12, 14, 15, 19, 21, 30, 31, 32, 33, 34, 37, 38, 40, 51, 65, 69, 71, 72, 74, 76, 82, 93
<i>Fregata minor</i>	Great Frigatebird	b	b	o		o		b	o	+	10, 14, 15, 19, 20, 21, 28, 30, 31, 32, 33, 34, 37, 40, 48, 51, 65, 69, 70, 71, 74, 76, 79, 82, 93, 95

Table 1 (continued)

Family, Species	Vernacular name	Region of New Caledonia								At sea	References	
		Chest..	d'Entr.	Loyalty	Gde Terre	N. Lag.	S. Lag.	Walpole	Matthew	Hunter		
<b>Accipitridae</b>												
<i>Pandion haliaetus leucocephalus</i>	Osprey			o	b	b	b			+	10, 15, 45, 81, 83, 84	
<b>Burhinidae</b>												
<i>Esacus magnirostris</i> <sup>7</sup>	Beach Thick-knee					b					3, 12, 15, 85, 86	
<b>Laridae</b>												
<i>Cathartes macromicrurus</i>	South Polar Skua									o	13, 42	
<i>Stercorarius pomarinus</i>	Pomarine Jaeger				o						87	
<i>Stercorarius parasiticus</i>	Arctic Jaeger									o	9, 10, 13	
<i>Larus novaehollandiae forsteri</i> <sup>8</sup>	Silver Gull			o	o	b	b			+	5, 14, 15, 17, 29, 39, 43, 44, 51, 81, 82	
<i>Sterna bergii cristata</i>	Great Crested Tern	b	o	b		b	b			+	5, 10, 12, 14, 15, 19, 23, 29, 30, 37, 38, 39, 40, 43, 51, 57, 68, 72, 82	
<i>Sterna dougallii bangsi</i>	Roseate Tern			o		b	b			+	5, 10, 12, 13, 14, 15, 19, 23, 29, 38, 39, 40, 51	
<i>Sterna sumatrana sumatrana</i>	Black-naped Tern	o	b	o		b	b			+	4, 10, 14, 15, 19, 23, 29, 32, 33, 38, 39, 40, 51	
<i>Sterna albifrons sinensis</i>	Little Tern				o		o			+	88	
<i>Sterna nereis excelsa</i> <sup>9</sup>	Fairy Tern	b				b	b			+	4, 10, 12, 15, 17, 23, 29, 39, 45, 51, 89, 94	
<i>Sterna anaethetus</i>	Bridled Tern		o			b	b			+	10, 15, 17, 23, 29, 39, 57, 90, 91	
<i>Sterna fuscata serrata</i>	Sooty Tern	b	b	b		b	l	o	b	o	+	10, 15, 19, 20, 21, 23, 26, 30, 31, 32, 33, 34, 39, 40, 48, 51, 57, 64, 67, 70, 73, 75, 76, 77, 78, 92, 94
<i>Anous stolidus pileatus</i>	Common Noddy	b	b	o		b	b	b	b	+	5, 10, 14, 15, 16, 17, 19, 20, 21, 23, 25, 26, 28, 29, 30, 31, 32, 33, 34, 39, 40, 51, 67, 71, 73, 76, 78, 95	
<i>Anous minutus minutus</i>	Black Noddy	b	b			l	b	b		+	5, 10, 15, 17, 19, 21, 23, 25, 27, 29, 30, 32, 39, 40, 51, 57, 67, 76, 73, 93, 95	
<i>Cygis alba caudata</i>	White Tern	o						b	o	l	+	10, 51, 59, 67, 71
<i>Procelsterna albivitta</i>	Grey Noddy							b	b	b	+	10, 21, 31, 48, 51, 57, 67

<sup>1</sup> One observation, of a light-phase individual, done in June 2005 at Yaté, Grande Terre (V. Chartendault, pers. comm.); dark phase otherwise (see Plates); <sup>2</sup> Known in the Chesterfield islands from the two specimens presumed to have been collected there by HMS *Herald* in 1858–1860 (Bouin et al. 2005). Not observed in the Chesterfield islands since then; <sup>3</sup> Reported as breeder in New Caledonia (BirdLife International 2000); <sup>4</sup> Historical breeder in the Southern lagoon, until 1974 (de Naurois and Rancurel 1978); <sup>5</sup> Dark and light phases in similar proportions in New Caledonia (Rancurel 1976); <sup>6</sup> Historical breeder in the Southern lagoon, until 1974 (de Naurois and Rancurel 1978); <sup>7</sup> Historical breeder on Huon island, d'Entrecasteaux reef and also reported from the eastern coast of Grande Terre by Layard and Layard (1882); <sup>8</sup> Historical breeder in d'Entrecasteaux reef (Layard and Layard 1882); <sup>9</sup> Historical breeder on Grande Terre (Layard and Layard 1882)

**Table 2.** Voucher specimens for New Caledonian seabirds. *NHM* Natural History Museum, London; *MNHN* Muséum national d'histoire naturelle, Paris; *AMNH* American Museum of Natural History, New York; *MV* Museum Victoria, Melbourne.

Species	Collection (specimen nos.)	Comments	Reference
<i>Pseudobulweria rostrata troussarti</i>	MNHN	Type specimen studied by Brasil (1917) and 12 skins and skeletons collected by de Naurois (1978)	de Naurois and Etard (1979)
	MNHN, AMNH, NHM	Several specimens studied by Villard <i>et al.</i> (2006)	Villard <i>et al.</i> (2006)
<i>Pterodroma heraldica</i>	NHM (nos. 62.6.22.10 and 88.5.18.110)	Two specimens assumed to have been collected by H.M.S. <i>Herald</i> in the Chesterfield in 1858-1860	Bourne <i>et al.</i> (2005)
<i>Pterodroma leucoptera caledonica</i>	AMNH (no. 824271, coll. No. NC20)	Specimen designated lectotype by Palma and Tennyson (2005)	Palma and Tennyson (2005)
	MNHN	Several specimens from Monts Dzumac collected by de Naurois	Imber and Jenkins (1981)
<i>Sterna dougallii</i>	F. Hannecart, private collection	Two stuffed specimens	F. Hannecart, pers. commun.
<i>Anous stolidus</i>	MV	One specimen from Observatory Islet, Middle Bellona Reefs	Bourne <i>et al.</i> (2005)



Adult Gould's Petrel (*Pterodroma leucoptera caledonica*) at the entrance of its burrow, Monts Dzumac, Grande Terre



Juvenile Tahiti Petrel (*Pseudobulweria rostrata trouessarti*), shore of Lac Yaté, Grande Terre



Pair of Wedge-tailed Shearwaters (*Puffinus pacificus*) at Pindai colony, Grande Terre



Juvenile Lesser Frigatebird (*Fregata ariel*) at Surprise island, d'Entrecasteaux reef



Red-tailed Tropicbird (*Phaeton rubricauda*) at Hunter island



Masked Booby (*Sula dactylatra*), adult and chick, Matthew island



Brown Noddy (*Anous stolidus*) at Kouaré islet, southern lagoon



Adult White Tern (*Gygis alba*) nesting on Walpole island

© SCO/Nicolas Barre

© SCO/Robert Aublin

© IRD/Philippe Borsig

© IRD/Philippe Borsig

© SCO/Nicolas Barre

© SCO/Robert Aublin

© SCO/Jérôme Spaggiari

© SCO/Jérôme Spaggiari



Adult Silver Gull (*Larus novaehollandiae forsteri*), Grande Terre



Grey Noddies (*Procelsterna albivitta*) on Matthew island



Roseate Tern (*Sterna dougalli*) on Signal island, southern lagoon



Fairy Tern (*Sterna nereis exsul*), southern lagoon



Dead coral and sand banks devoid of vegetation: Sèche-Croissant cayes, southern lagoon



Low islet of coral sand covered by vegetation: Redika islet, southern lagoon



Rocky islet: Mato islet, southern lagoon



Steep and forested slopes of the central chain of mountains: Koniambo massif, Grande Terre

**DOCUMENTS  
SCIENTIFIQUES  
et TECHNIQUES**

**II7**

**Volume spécial  
Deuxième édition**

# **COMPENDIUM of marine species from New Caledonia**



Edited by  
Claude E. PAYRI  
Bertrand RICHER DE FORGES



**Institut de recherche  
pour le développement**

CENTRE DE NOUméA

# *DOCUMENTS SCIENTIFIQUES et TECHNIQUES*

---



Institut de recherche  
pour le développement

Publication éditée par:  
Centre IRD de Nouméa  
BP A5, 98848 Nouméa CEDEX  
Nouvelle-Calédonie  
Téléphone : (687) 26 10 00  
Fax : (687) 26 43 26

L'IRD propose des programmes regroupés en 5 départements pluridisciplinaires :

- I DME Département milieux et environnement
- II DRV Département ressources vivantes
- III DSS Département sociétés et santé
- IV DEV Département expertise et valorisation
- V DSF Département du soutien et de la formation des communautés scientifiques du Sud

**Modèle de référence bibliographique à cette revue :**

Adjeroud M. *et al.*, 2000. Premiers résultats concernant le benthos et les poissons au cours des missions TYPATOLL.  
*Doc. Sci. Tech. II* 3, 125 p.

Première de couverture : Récif corallien (Côte Ouest, NC) © IRD C.Geffray  
Vignettes : voir les planches photographiques

Quatrième de couverture : *Platygyra sinensis* © IRD C.Geffray

Matériel de plongée sous-marine	L'Aldric, moyen naviguant de l'IRD
© IRD/C.Geffray	© IRD/J.-M. Boré
Récoltes et photographies sous-marines en scaphandre autonome	Traitement des récoltes en laboratoire
© IRD/J.-L. Menou	© IRD/L. Mattio

CONCEPTION/MAQUETTE/MISE EN PAGE  
MAQUETTE DE COUVERTURE  
PLANCHES PHOTOGRAPHIQUES  
TRAITEMENT DES PHOTOGRAPHIES

JEAN PIERRE MERMOUD/ISABELLE RITZENTHALER  
CATHY GEOFFRAY/MINA VILAYLECK  
CATHY GEOFFRAY/JEAN-LOUIS MENOU/GEORGES BARGIBANT  
NOËL GALAUD

La traduction en anglais des textes d'introduction, des Ascidies et des Echinodermes a été assurée par EMMA ROCHELLE-NEWALL,  
la préface par MINA VILAYLECK.

Ce document est publié par le Service ISC du Centre IRD de Nouméa et financé par le Ministère de la Recherche et de la Technologie.

# **COMPENDIUM OF MARINE SPECIES FROM NEW CALEDONIA**

SECOND EDITION

Edited by  
CLAUDE E. PAYRI, BERTRAND RICHER DE FORGES

