

Age-structured gametocyte allocation links immunity to epidemiology in malaria parasites

Richard E Paul^{*1,2}, Sarah Bonnet^{3,4}, Christian Boudin⁵,
Timoleon Tchuinkam³ and Vincent Robert^{5,6}

Address: ¹Laboratoire d'Entomologie Médicale, Institut Pasteur de Dakar, 36 Avenue Pasteur, BP 220 Dakar, Sénégal, ²Laboratoire de Génétique de la réponse aux infections chez l'homme, Institut Pasteur, 28 rue de Dr. Roux, F-75724, Paris cedex 15, France, ³Laboratoire IRD de Recherche sur le Paludisme, Organisation de Coordination pour la lutte Contre les Endemies en Afrique Centrale, P.O. Box 288, Yaounde, Cameroon, ⁴Ecole Nationale Vétérinaire de Nantes, Service de parasitologie, UMR ENVN/INRA 1034 Interactions Hôte-Parasite-Milieu, Atlanpole-La Chantrerie, B.P. 40706, 44307 Nantes cedex 03, France, ⁵UR Paludisme Afro-tropical, Institut de Recherche pour le Développement, B.P.1386 Dakar, Sénégal and ⁶Institut de Recherche pour le Développement/Muséum National d'Histoire Naturelle, 61 rue Buffon, case courrier 52, 75231 Paris cedex 05, France

Email: Richard E Paul* - rpaul@pasteur.fr; Sarah Bonnet - bonnet@vet-nantes.fr; Christian Boudin - boudin@ird.sn; Timoleon Tchuinkam - timotchuinkam@yahoo.fr; Vincent Robert - v.robert@mnhn.fr

* Corresponding author

Published: 12 September 2007

Received: 23 March 2007

Malaria Journal 2007, **6**:123 doi:10.1186/1475-2875-6-123

Accepted: 12 September 2007

This article is available from: <http://www.malariajournal.com/content/6/1/123>

© 2007 Paul et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Despite a long history of attempts to model malaria epidemiology, the over-riding conclusion is that a detailed understanding of host-parasite interactions leading to immunity is required. It is still not known what governs the duration of an infection and how within-human parasite dynamics relate to malaria epidemiology.

Presentation of the hypothesis: Immunity to *Plasmodium falciparum* develops slowly and requires repeated exposure to the parasite, which thus generates age-structure in the host-parasite interaction. An age-structured degree of immunity would present the parasite with humans of highly variable quality. Evolutionary theory suggests that natural selection will mould adaptive phenotypes that are more precise (less variant) in "high quality" habitats, where lifetime reproductive success is best. Variability in malaria parasite gametocyte density is predicted to be less variable in those age groups who best infect mosquitoes. Thus, the extent to which variation in gametocyte density is a simple parasite phenotype reflecting the complex within-host parasite dynamics is addressed.

Testing the hypothesis: Gametocyte densities and corresponding infectiousness to mosquitoes from published data sets and studies in both rural and urban Cameroon are analysed. The mean and variation in gametocyte density according to age group are considered and compared with transmission success (proportion of mosquitoes infected). Across a wide range of settings endemic for malaria, the age group that infected most mosquitoes had the least variation in gametocyte density, i.e. there was a significant relationship between the variance rather than the mean gametocyte density and age-specific parasite transmission success. In these settings, the acquisition of immunity over time was evident as a decrease in asexual parasite densities with age. By contrast, in an urban setting, there were no such age-structured relationships either with variation in gametocyte density or asexual parasite density.

Implications of the hypothesis: Gametocyte production is seemingly predicted by evolutionary theory, insofar as a reproductive phenotype (gametocyte density) is most precisely expressed (i.e. is most invariant) in the most infectious human age group. This human age group would thus be expected to be the habitat most suitable for the parasite. Comprehension of the immunology of malaria, a requisite for any vaccine strategies, remains poor. Immunological characterization of the human population stratified by parasite gametocyte allocation would be a step forward in identifying the salient immunological pathways of what makes a human a good habitat.

Background

Mathematical models of infectious diseases and most especially those based on the calculation of R_0 , the basic reproductive number, have proved to be very powerful and robust in interpreting epidemiological trends, most notably for viral and macro-parasitic diseases [1]. However, despite a long history of attempts to model malaria epidemiology, and most especially that of lethal human malaria *Plasmodium falciparum*, the over-riding conclusion is that a detailed understanding of host-parasite interactions leading to immunity is required. The parasite's high degree of genetic diversity and mechanisms of clonal antigenic variation highlight the complex nature of the human-parasite interaction. However, it is still not known what governs the duration of an infection and how within-human parasite dynamics relate to malaria epidemiology. Indeed, controversy still rages over the rate of acquisition of immunity to malaria [2,3].

The classical Ross-Macdonald model

($R_0 = \frac{ma^2bc}{\mu\gamma} \exp^{-\mu T}$) captures the essence of malaria epi-

demiology and consists of mosquito components that determine, to a large extent, the force of infection [mosquito density (m), biting rate (a) and mortality rate (μ) and the duration of sporogonic development in the vector (T)] and human components that directly relate to the human-parasite interaction. Two of these parameters are concerned with within-host parasite dynamics: the human recovery rate from infection (γ) and the infectiousness of an infected individual to mosquitoes (c). Parameter (b) is the proportion of infective bites that lead to an infection in man.

Immunity to *P. falciparum* develops slowly and requires repeated exposure to the parasite. This generates age-structure in the host-parasite interaction and most notably in the duration of episodes of infection [4]. In areas endemic and stable for malaria, recovery rate is lowest in the young/intermediate age groups [5,6]. Recent studies [2,7], however, suggest that the rate of absolute recovery from infection may not increase with age, even though infect-

ing parasite density may be reduced. Epidemiological models incorporating age-dependency, whether through increased absolute recovery from infection or increased parasite death rate, improve markedly the fit with reality [1]. Measures of human infectiousness to mosquitoes have also shown strong age-specificity such that infections in particular age groups most successfully infect mosquitoes [8-11].

In this work, an attempt is made to identify key measurable parameters linking infection and immunity to epidemiology.

Presentation of the hypothesis

Age-dependency in the development of an immune response against the asexual parasite stages would present the parasite with habitats of highly variable quality. Evolutionary theory predicts that, in a spatially heterogeneous environment, natural selection will mould adaptive phenotypes such that they are more precise (less variant) in the better and the more frequent habitats and less so in infrequent habitats and those where reproductive performance is poor [12], i.e. natural selection will act on both the mean and the variance of the phenotype in question. Heterogeneity in habitat quality can severely alter R_0 and it is trivial to note that optimising transmission (c) in the age groups where recovery rate (γ) is the lowest will significantly increase R_0 . If the parasite is indeed optimising transmission according to the immune-dependent quality of individuals, this should be apparent from the within-host parasite dynamics and notably that pertaining to gametocyte production.

Testing the hypothesis

Gametocyte densities and corresponding infectiousness to mosquitoes from published data sets [9,10,13] and studies in both rural [11,14] and urban [15] settings in Cameroon are examined. The mean and variation in gametocyte density (as proxies of the parasite's transmission phenotype) according to age group are considered and compared with actual measures of transmission success (the proportion of mosquitoes that become infected). Maturation of gametocytes occurs slowly and infections with apparent gametocytes may not be infectious at that

exact time but may be so hours later. Therefore, all observed gametocyte densities are included. The age groups chosen in the published and the rural Cameroon data sets reflect the classical age distribution as defined by malaria prevalence rates that provide a gross measure of the force of infection and gradual development of immunity. The decrease in asexual parasite densities with age is a more sensitive indication of the slow acquisition of immunity that generates age structure and is thus presented. By contrast, malaria transmission in urban settings is low and unstable, resulting in a poor acquisition of immunity. Therefore, age-stratification of the urban Cameroon data set would not be expected to reflect age-specific development of immunity or be apparent from asexual parasite densities.

Cameroon study sites and data collection

The study sites and methods have been previously published [11,14,15]. Briefly, in the rural setting, individuals, symptomatic or not, were recruited in two adjacent villages in the district of Mengang, South Cameroon, in an area of degraded forest with seasonal transmission (intensity of 170 infected bites/person per year) and hyperendemic malaria (*P. falciparum* prevalence rates range from 49% to 82%, depending on the season). Mosquitoes were gorged directly on individuals whether positive or not for gametocytes. In the urban setting, patients were recruited at a Public Health Centre in central Yaounde, the capital town of Cameroon. From those individuals positive for gametocytes by thick blood smear, venous blood was immediately placed into a membrane feeder upon which mosquitoes gorged. Previous comparative studies have shown that although direct feeding of mosquitoes on infectious individuals yields higher infections in mosquitoes than feeding through a membrane, there is good concordance between the two tests for both mosquito infection percentages and oocyst loads [16,17]. In both studies, only experiments where at least 20 mosquitoes survived to day of dissection for oocyst counts (day 7 post-gorging) were used in analyses. The mean number of mosquitoes dissected per experiment was 35 and 36 in the rural and urban settings respectively. Blood parasite counts were established by thick smear per 1,000 leucocytes for asexual parasite density and 1,000 and 3,000 leucocytes for gametocytes respectively in urban and rural settings, assuming an average number of 8,000 leucocytes per microlitre of blood. The studies used the same laboratory mosquito strain reared in the same insectary (*Anopheles gambiae* s.s. Yaounde strain; OCEAC insectary).

Statistical analyses

The effects of age group and gametocyte density per infection on transmission success in the Cameroon data sets were analysed in Genstat version 7 using logistic regression specifying a binomial error structure. The effect of age

group on asexual parasite density was analysed by log-linear regression fitting a GLM, specifying a Poisson error structure. Dispersion parameters were estimated and thus the analyses generate an F statistic in the analysis of deviance.

Using data from the Cameroon studies as well as the two published studies, how the mean and the variability in gametocyte density per age group affected transmission success were then examined. This combined data set was analysed by two alternative methods: a REML meta-analysis and by fitting a GLMM (Mixed model) with study specified in the random model and the mean proportion of mosquitoes infected by age group as the response variate. A REML meta-analysis takes into account differences between sites by incorporating the residual variance of each experiment. GLMMs account for variation due to study site and calculate the variance component due to this random factor independently of that due to the factor of interest – i.e. the response variate. Because the measure of variability in gametocyte density per age group was previously given as confidence intervals (CI) in the published studies, this measure was used for the Cameroon data sets. The lower CI subtracted from the upper CI was calculated to generate a value of dispersion in gametocyte density by age group. Both this value and mean gametocyte density per age group were normalized for each study by dividing by the lowest value, thus giving the lower value of 1 per study. Such normalization enables comparison of the general effect of gametocyte density and variability in density by age group without study specific effects, most evident of which are the very different overall gametocyte densities. Both the normalized gametocyte density mean and variability were then fitted as explanatory variates. Because the data were over-dispersed, a dispersion parameter was estimated. Statistical significance is presented as Wald statistics, which approximate to a χ^2 distribution. For the Cameroon data sets, variance/mean ratios and the Standardized Morisita index (I_p) were additionally calculated. Sample size influences both confidence intervals and variances. The larger the sample size the smaller these measures will be for a data set of the same intrinsic variability. The Standardized Morisita index (I_p), however, is considered the best measure of dispersion being independent of sample size [18]. To calculate this index, the Morisita's index of dispersion (I_d) and two critical values, the Uniform index (M_u) and the clumped index (M_c) are calculated as follows:

$$I_d = n \left[\frac{\sum x^2 - \sum x}{(\sum x)^2 - \sum x} \right]$$

$$M_u = \frac{\chi_{.975}^2 - n + \sum x}{(\sum x) - 1}$$

$$M_c = \frac{\chi_{.025}^2 - n + \sum x}{(\sum x) - 1}$$

where n is the sample size, x is the number of gametocytes in each individual, $\chi_{.975}^2$ and $\chi_{.025}^2$ are the values of the chi-squared with (n-1) degrees of freedom that have 97.5% or 2.5% of the area to the right. The Standardized Morisita index (I_p) is then calculated by one of the four following formulae:

(a) when $I_d \geq M_c > 1$, $I_p = 0.5 + 0.5 \times \left(\frac{I_d - M_c}{n - M_c} \right)$, (b) when

$M_c > I_d \geq 1$, $I_p = 0.5 \times \left(\frac{I_d - 1}{M_u - 1} \right)$,

(c) when $1 > I_d > M_u$, $I_p = -0.5 \times \left(\frac{I_d - 1}{M_u - 1} \right)$ and (d) when

$1 > M_u > I_d$, $I_p = -0.5 + 0.5 \times \left(\frac{I_d - M_u}{M_u} \right)$.

The Standardized Morisita index (I_p) ranges from -1 to +1. Random patterns give a value of zero, clumped patterns above zero and uniform patterns below zero. With respect to gametocyte data, the higher the value above zero the more over-dispersed, and thus more variable, are the gametocyte densities.

Results

Mosquito infection rates have been classically shown to correlate positively with individual gametocyte density. This positive correlation was confirmed here for the Cameroon rural data set, including only those individuals who infected at least one mosquito ($F_{1,45} = 13.9 P < 0.001 N = 50$). However, gametocyte density *per se* explained only 16.6% of the variation in transmission success and there was a significant effect of age group on mosquito infection rates ($F_{3,45} = 3.43, P = 0.025 N = 50$). Within any of the published data sets in Kenya and Sri Lanka and the rural Cameroon data set, the mean gametocyte densities were relatively constant across age groups (Table 1) and typical of those observed in rural endemic situations across the globe. It is notable that the most infectious age groups in any of these settings were not those with the highest mean gametocyte densities (Table 1). Indeed, mean gametocyte

density per age group did not correlate with mosquito infection rates (GLMM $\chi^2_1 = 1.03, P = 0.31$). By contrast, it is striking that the age groups that had the lowest variability in gametocyte density infected the highest proportion of mosquitoes (Figure 1) (REML meta-analysis: $\chi^2_1 = 7.95, P = 0.005$; GLMM $\chi^2_1 = 6.72, P = 0.01$). Variation in gametocyte density (i.e. gametocyte precision) explained 17.1% of variation in age group transmission success. Using the rural Cameroon data set, this relationship was confirmed when measuring gametocyte variability using several different dispersion indices (Table 1), including the Standardized Morisita index (I_p), which is considered independent of sample size [18]; all dispersion indices show the same pattern in the Cameroon data set. Moreover, Table 1 shows that, with the exception of *P. vivax* gametocyte data from Sri Lanka, gametocyte variability is lower in the best transmitting age group rather than the age group with the largest sample size. Larger sample sizes would be expected to yield lower values of dispersion if intrinsic variability were the same among groups and could therefore explain degrees of dispersion. Here this is not the case.

Immune-generated age-structure is apparent in rural Cameroon, where asexual parasite densities (and variation in density) decreased significantly with age from a mean of 3965 parasites/ μ l (SE \pm 690) in the youngest age group to 829 parasites/ μ l (SE \pm 168) in the oldest age group ($F_{3,410} = 11.51 P < 0.001 N = 413$)(Table 2). Likewise, the prevalence of infection and the proportion of infections bearing gametocytes also decreased significantly in the eldest age group [11]. By contrast in the urban setting there was no




Figure 1
Age group variation in gametocyte density with corresponding infectiousness to mosquitoes. Data shown in Table 1. The higher minus the lower 95% confidence intervals are taken as a measure of gametocyte density variation and are normalized within each study area by dividing by the lowest value, thus giving the lower value of 1. Circles – *P. vivax* Sri Lanka [9]; Triangles – *P. falciparum* Sri Lanka [9]; Squares – *P. falciparum* Kenya [10]; Crosses – *P. falciparum* Cameroon [11].

Table 1: Age-specific gametocyte densities with measures of transmission success in rural Cameroon [11] and published data sets [9,10].

Study Site	age group	N	Gametocyte density				Mean proportion mosquitoes infected \pm s.e. (n)	Proportional contribution to human infectious reservoir
			Geometric mean/ μ l (\pm CI _{95%})	Normalized 95% Confidence Intervals	Standardized Morisita Index	Variance/mean		
Cameroon	0-5	40	3.56 (0.8-6.32)	1.04	0.521	12.4	0.22 \pm 0.05 (17)	0.32
	6-9	36	3.54 (0.88-6.2)	1.00	0.518	10.0	0.25 \pm 0.05 (14)	0.43
	10-15	28	2.84 (0.04-5.64)	1.05	0.531	11.6	0.13 \pm 0.06 (7)	0.11
	15+	33	3.71 (0.64-6.88)	1.20	0.528	13.6	0.09 \pm 0.04 (12)	0.10
Kenya	1-4	33	29.3 (27.1-31.7)	1.00			0.16 (20)	0.38
	5-9	36	30.7 (27.8-33.8)	1.33			0.14 (20)	0.34
	10-14	17	29 (26.0-32.9)	1.50			0.07 (18)	0.09
Gametocytes/red blood cells ($\times 10^{-2}$) \pm SD								
Sri Lanka <i>P. vivax</i>	0-5	47	2.2 \pm 3	1.85			0.14	0.11
	6-15	122	1.3 \pm 2	1.00			0.32	0.27
	16-25	75	1.9 \pm 2	1.42			0.28	0.32
	26-50	82	1.4 \pm 2	1.11			0.23	0.28
<i>P. falciparum</i>	0-5	26	0.07 \pm 0.3	1.17			0.08	0.01
	6-20	130	0.09 \pm 0.4	1.00			0.49	0.68
	21-50	187	0.13 \pm 0.5	1.27			0.39	0.30
	50+	27	0.16 \pm 0.5	2.20			0.04	0.01

Differences in the upper and lower 95% confidence intervals are normalized within each study by dividing by the lowest value. The proportional contribution to the infectious reservoir is the proportion of an age group that are infectious to mosquitoes multiplied by the proportion of mosquitoes that become infected. In bold, values for the least dispersed gametocyte densities and the highest mosquito infection rates; in *italics* the highest gametocyte densities. N is the number of individuals with gametocyte parasites and n is the number of individuals infecting mosquitoes.

effect of age group on asexual parasite densities ($F_{4,90} = 0.68$ $P = 0.68$, $N = 95$), with a high overall mean of 7626 parasites/ μ l (SE \pm 1013) (Table 2), as would be expected if acquisition of immunity were low. This lack of age-structure is mirrored in mosquito infection rates where there was no effect of age ($F_{4,80} = 1.3$, $P = 0.27$, $N = 86$), nor was there any relationship between gametocyte variability and transmission in the urban sample (Table 3) ($F_{1,3} = 0.12$ $P = 0.752$). However, there was a strong effect of gametocyte density *per se* ($F_{1,84} = 23.1$ $P < 0.001$ $N = 86$), which accounted for 28.1% of variation in mosquito infection rates.

Implications of the hypothesis

These analyses reveal a significant relationship between precision in gametocyte density and the transmission success of age groups in endemic settings of malaria transmission. Importantly, the actual age group most infectious to mosquitoes varied according to the transmission intensity, becoming younger with increasing intensity. Mean

gametocyte densities by age group were not, however, related to transmission success except in urban Cameroon. As expected in endemic conditions, an age-dependent degree of acquired immunity was evident from the gradual decrease in mean asexual parasite densities. By contrast, the relationship between gametocyte variability and transmission success did not occur in an urban setting where individuals of all ages did not differ in their overall degree of immunity, as apparent from their similar asexual parasite densities.

Studies to date have examined human infectiousness to mosquitoes at three distinct levels: (1) Infectiousness of the individual according to gametocyte density [e.g. [19]], (2) the age group reservoir of infection within a site [e.g. [8,11,20]], and most recently (3) across populations of differing intensity [21]. Here, a gametocyte phenotype (low variance of gametocyte density) measured at the level of age group, but which incorporates individual infection data, is consistently related to transmission suc-

Table 2: Age-specific asexual *Plasmodium* spp. prevalence rates and densities.

Study Site	age group	Trophozoite prevalence (%)	N	Mean trophozoite density (\pm CI _{95%})
Cameroon Rural	0-5	66.4	42	3965 (2613-5317)
	6-9	71	119	1726 (1199-2253)
	10-15	69.5	104	1387 (879-1895)
	15+	39.5	148	829 (500-1158)
Cameroon Urban	5-10	NA	14	7600 (2720-12480)
	11-14		25	8471 (4616-12326)
	15-20		20	9721 (5103-14339)
	21-25		19	5842 (2169-9515)
	25+		17	5934 (2020-9848)
Kenya	1-4	89	63	1603 (968-2654)
	5-9	94	83	654 (444-963)
	10-14	90	88	188 (134-264)
Trophozoite Incidence (%)				
Sri Lanka				
<i>P. vivax</i>	0-5	10	753	NA
	6-15	13.8	829	
	16-25	9.4	712	
	26-50	6.5	1080	
	50+	3.9	251	
<i>P. falciparum</i>	0-5	7.4	753	
	6-15	20.1	829	
	16-25	17.4	712	
	26-50	12.8	1080	
	50+	7.6	251	

N is the number of individuals contributing to the calculation of mean trophozoite densities or, for the case of Sri Lanka, the incidence rates. NA is Not Available.

cess across populations from differing transmission intensities. This relationship is consistent with the thesis that natural selection should mould phenotypes (gametocyte density) such that they are less variant in the most productive habitat. Why certain age groups are the most productive (i.e. infectious) is less clear. Gametocyte epidemiology is complex but does display some age-structure (For an excellent review see [22]). In semi-immune populations, gametocyte prevalence does not always decrease with age [20] and, when it does, tends to reflect age-asexual parasite prevalence profiles [8,10,23]. This suggests that low gametocyte prevalence reflects the development of immunity against the asexual stages [24] rather than any effective anti-gametocyte immune response. However, a recent reappraisal suggests that gametocyte prevalence may decrease more rapidly with age than asexual parasite prevalence rates and that there might be addi-

tional anti-gametocyte immune responses [22]. Moreover, when gametocytes are present in older age groups, their densities, relative to the asexual parasite density whence they arise, are generally increased [22]. This is consistent with the known influence of both specific and non-specific anti-asexual parasite immune mechanisms on the rate of conversion from asexual to gametocyte stage parasites [25-27].

How such variation in gametocyte prevalence rates and densities per human age group relate to transmission success is unclear. Although increasing gametocyte density tends to result in greater infectiousness to mosquitoes [19], it has been repeatedly demonstrated that high gametocyte densities do not guarantee high mosquito infection rates [15,19,20]. Recent work within the context of developing a transmission-blocking vaccine has highlighted concurrent gametocyte density, but not age, as being positively correlated with transmission-blocking immunity [28]. Not unsurprisingly, there exists a minimum gametocyte density necessary for effective transmission, but one which is variable according to study site [29]. Low asexual parasite densities, such as those found in older semi-immune individuals, might therefore generate insufficient gametocytes for transmission. In the rural Cameroon data set here, however, mean and even mode gametocyte densities did not differ among age groups. Conversely, high asexual parasite densities, such as those seen in infants, generate an excess of gametocytes [30] and thus may induce transmission blocking anti-gametocyte specific immunity. Moreover, concurrent fever, induced by high asexual parasitaemia, reduces infectivity [31,32]. An age-dependent pyrogenic threshold of *P. falciparum* parasitaemia has been demonstrated [33]; notably the individuals most tolerant of parasite density were neither the very young nor the older age groups. Such a threshold is expected to vary according to the transmission intensity. This might be an explanation for our observed differences in age of the best-transmitting groups in areas of differing transmission intensity.

Although age-specific transmission success could be explained with reference to age-specific pyrogenic thresholds, it is not clear why variation in gametocyte density should correlate similarly with age. Variation in asexual parasitaemia was found to decrease with age as might be expected with the development of immunity and yet variation in gametocyte density did not. Although variation in gametocyte density in the youngest age groups may thus be explained as a consequence of the variation in asexual parasite densities, the same argument can not simultaneously explain the increased variability in the older age groups. Specific anti-asexual parasite immunity may additionally be involved in older age groups acting indirectly in conflicting ways on gametocyte density: both reducing

Table 3: Age-specific gametocyte densities with measures of transmission success in urban Cameroon [15].

age group	N	Gametocyte density				Mean proportion of mosquitoes infected ± s.e. (n)	Proportional contribution to human infectious reservoir
		Geometric mean/μl (± CI _{95%})	Normalized 95% Confidence Intervals	Standardized Morisita Index	Variance/mean		
5-10	26	78.2 (± 51.2)	1.00	0.518	129.7	0.2 ± 0.05 (17)	0.23
11-14	24	<i>118.7 (± 119.8)</i>	2.34	0.536	369.6	0.16 ± 0.04 (15)	0.17
15-20	26	107.2 (± 93.4)	1.82	0.528	290.0	0.24 ± 0.04 (14)	0.22
21-25	33	76.5 (± 52.4)	1.02	0.519	168.1	0.21 ± 0.05 (24)	0.26
25+	29	48.9 (± 59.4)	1.16	0.539	233.6	0.12 ± 0.03 (16)	0.11

Differences in the upper and lower 95% confidence intervals are normalized by dividing by the lowest value. The proportional contribution to the infectious reservoir is the proportion of an age group that are infectious to mosquitoes multiplied by the proportion of mosquitoes that become infected. In bold, values for the least dispersed gametocyte densities and the highest mosquito infection rates; in *italics* the highest gametocyte densities. N is the number of individuals with gametocyte parasites and n is the number of individuals infecting mosquitoes. No age was recorded for 4 of the 90 urban individuals infecting mosquitoes.

the asexual parasite source and increasing the rate of conversion to gametocytes. This dual effect on gametocyte density could generate increased variability in both gametocyte density and transmission success. Potentially there exists a hitherto unidentified anti-gametocyte specific response that increases with the acquisition of anti-parasite (asexual) immunity.

Hitherto unmentioned, is the documented relationship between gametocyte production and haematological factors [34-36]. Although oxidative stress by cytokine response to infection can lead to alteration of uninfected RBC membranes and thus enhance their removal by the spleen [37], to date no age-specific anti-parasite immune response affecting haemolysis and hence erythropoiesis has been described [38]. Indeed, in regions of high intensity, anaemia occurs in infants and in certain risk groups (pregnant women); older age groups do not apparently suffer from malaria-induced anaemia though haematopoietic activity has not been surveyed as such across a population.

Malaria parasite gametocyte allocation has proven highly informative, revealing adaptive strategies that optimize transmission according to parasite population structure [39,40] and to ensure transmission success [41-43]. Here, examination of variation in gametocyte density reveals a measure that captures the essence of the parasite's behaviour with respect to transmission. The parasite has a very precise gametocyte allocation in individuals of an age from whom transmission to mosquitoes is best. This suggests that the parasite is best adapted to specific age groups, i.e. the parasite is being more precise in the better habitat.

The human immune response characterizes habitat quality of humans for malaria parasites. Infection longevity would be expected to be beneficial for the parasite and thus those individuals with the longest duration of infection be predicted to be the highest quality habitats. The

duration of an infection has proved, however, a very difficult parameter to measure and recently the subject of renewed discussion [2,3,7]. Nevertheless, estimates made in areas of high transmission intensity suggest that the longest duration occurs in intermediate age groups [5,6]; it is not known, however, how this relates to transmission success or what are the principle immunological mechanisms involved. Immuno-tolerance is an accepted feature of malaria parasite sustained infection and a role for the innate immune response has been recently proposed [44]. The age groups most infectious to mosquitoes are those who seemingly have the lowest recovery rate from infection [5,6], when extrapolating from studies in regions of similar transmission intensity to the Kenyan and Cameroon study examined here [10,11]. The longer duration of infection in individuals apparently tolerant to increased parasite densities may provide sufficient time, in a relatively stable environment, with which to produce gametocytes. This would provide the link between immune-dependent habitat quality and transmission success.

Malaria parasite epidemiology may be best understood with reference to reproductive allocation. A deeper understanding of the immunological basis to the host-parasite interaction and what defines a "better host" will be more readily achieved by inspection of host anti-parasite immune responses in age groups defined by the reproductive precision of their infecting parasites. Importantly, the best human habitats were those who had had previous exposure to the parasite. Vaccine strategies designed to reduce morbidity in infants may be at risk of generating of better quality individuals for parasite transmission. In the light of the current widespread vaccine trials this must be carefully monitored to avoid undesirable side-effects.

Authors' contributions

RELPC conceived the idea, analyzed the data and wrote the paper. SB & CB performed all the data collection in rural Cameroon, made comments on and edited the paper. TT

& VR performed all the data collection in Yaounde, TT made comments on and edited the paper and VR co-wrote sections of the paper.

Acknowledgements

This work was supported by the Strategic *Anopheles* Horizontal Research Programme, Institut Pasteur.

References

- Anderson RM, May RM: *Infectious diseases of humans: Dynamics and Control* Oxford: Oxford University Press; 1991.
- Smith DL, Dushoff J, Snow RW, Hay SI: **The entomological inoculation rate and *Plasmodium falciparum* in African children.** *Nature* 2005, **438**:492-495.
- Filion GJ, Paul RE, Robert V: **Transmission and immunity: the importance of heterogeneity in the fight against malaria.** *Trends Parasitol* 2006, **22**:345-348.
- Bruce MC, Donnelly CA, Packer M, Lagog M, Gibson N, Narara A, Walliker D, Alpers MP, Day KP: **Age- and species-specific duration of infection in asymptomatic malaria infections in Papua New Guinea.** *Parasitology* 2000, **121**:247-256.
- Bekessy A, Molineaux L, Storey J: **Estimation of incidence and recovery rates of *Plasmodium falciparum* parasitaemia from longitudinal data.** *Bull World Health Organ* 1976, **54**:685-691.
- Smith T, Beck HP, Kitua A, Mwankusy E, Felger I, Fraser-Hurt N, Irion A, Alonso P, Teuscher T, Tanner M: **Age dependence of the multiplicity of *Plasmodium falciparum* infections and of other malariological indices in an area of high endemicity.** *Trans R Soc Trop Med Hyg* 1999, **93**(Suppl 1):15-20.
- Sama S, Owusu-Agyei S, Felger I, Dietz K, Smith T: **Age and seasonal variation in the transition rates and detectability of *Plasmodium falciparum* malaria.** *Parasitology* 2006, **132**:13-21.
- Muirhead-Thomson RC: **The malarial infectivity of an African village population to mosquitoes (*Anopheles gambiae*).** *Am J Trop Med Hyg* 1957, **6**:971-979.
- Gamage-Mendis AC, Rajakaruna J, Carter R, Mendis KN: **Infectious reservoir of *Plasmodium vivax* and *Plasmodium falciparum* malaria in an endemic region of Sri Lanka.** *Am J Trop Med Hyg* 1991, **45**:479-487.
- Githeko AK, Brandling-Bennett AD, Beier M, Atieli F, Owaga M, Collins FH: **The reservoir of *Plasmodium falciparum* malaria in a holoendemic area of western Kenya.** *Trans R Soc Trop Med Hyg* 1992, **86**:355-358.
- Bonnet S, Paul REL, Gouagna C, Safeukui I, Meunier J-Y, Gounoue R, Boudin C: **Level and dynamics of malaria transmission and morbidity in an equatorial area of South Cameroon: entomological and parasitological data in a population exposed to low and seasonal transmission.** *Trop Med Int Health* 2002, **7**:1-8.
- Kawecki TJ, Stearns SC: **The evolution of life histories in spatially heterogeneous environments: optimal reaction norms revisited.** *Evol Ecol* 1993, **7**:155-174.
- Mendis C, Gamage-Mendis AC, De Zoysa AP, Abhayawardena TA, Carter R, Herath PR, Mendis KN: **Characteristics of malaria transmission in Kataragama, Sri Lanka: a focus for immunoparasitological studies.** *Am J Trop Med Hyg* 1990, **42**:298-308.
- Bonnet S, Gouagna CL, Paul RE, Safeukui I, Meunier J-Y, Boudin C: **Estimation of malaria transmission from humans to mosquitoes in two neighbouring villages in south Cameroon: evaluation and comparison of several indices.** *Trans R Soc Trop Med Hyg* 2003, **97**:53-59.
- Tchuinkam T, Mulder B, Deschering K, Stoffels H, Verhave JP, Cot M, Carnevale P, Meuwissen JH, Robert V: **Experimental infections of *Anopheles gambiae* with *Plasmodium falciparum* of naturally infected gametocyte carriers in Cameroon: factors influencing the infectivity to mosquitoes.** *Trop Med Parasitol* 1993, **44**:271-276.
- Bonnet S, Gouagna C, Safeukui I, Meunier J-Y, Boudin C: **Comparison of artificial membrane feeding with direct skin feeding to estimate infectiousness of *Plasmodium falciparum* gametocyte carriers to mosquitoes.** *Trans R Soc Trop Med Hyg* 2000, **94**:103-106.
- Awono-Ambene HP, Diawara L, Robert V: **Comparison of direct and membrane feeding methods to infect *Anopheles arabiensis* with *Plasmodium falciparum*.** *Am J Trop Med Hyg* 2001, **64**:32-34.
- Krebs CJ: *Ecological Methodology* 2nd edition. Harlow: Addison Wesley Longman; 1999.
- Jeffery GM, Eyles DE: **Infectivity to mosquitoes of *Plasmodium falciparum* as related to gametocyte density and duration of infection.** *Am J Trop Med Hyg* 1955, **4**:781-789.
- Boudin C, Olivier M, Molez JF, Chiron JP, Ambroise-Thomas P: **High human malarial infectivity to laboratory-bred *Anopheles gambiae* in a village in Burkina Faso.** *Am J Trop Med Hyg* 1993, **48**:700-706.
- Killeen GF, Ross A, Smith T: **Infectiousness of malaria-endemic human populations to vectors.** *Am J Trop Med Hyg* 2006, **75**:38-45.
- Drakeley C, Sutherland C, Bousema JT, Sauerwein RW, Targett GA: **The epidemiology of *Plasmodium falciparum* gametocytes: weapons of mass dispersion.** *Trends Parasitol* 2006, **22**:424-30.
- Jones TR, McElroy PD, Oster CN, Beier JC, Oloo AJ, Onyango FK, Chumo DK, Sherwood JA, Hoffman SL: ***Plasmodium falciparum* gametocytaemia in Kenyan children: associations among age, intensity of exposure to transmission, and prevalence and density of subsequent gametocytaemia.** *Am J Trop Med Hyg* 1997, **56**:133-136.
- Carter R, Mendis K: **Transmission immunity in malaria: reflections on the underlying immune mechanisms during natural infections and following artificial immunization.** *Mem Inst Oswaldo Cruz Rio* 1992, **87**(Suppl III):169-173.
- Smalley ME, Brown J: ***Plasmodium falciparum* gametocytogenesis stimulated by lymphocytes and serum from infected Gambian children.** *Trans R Soc Trop Med Hyg* 1981, **75**:316-317.
- Kwiatkowski D, Sambou I, Twumasi P, Greenwood BM, Hill AVS, Manogue KR, Cerami A, Castracane J, Brewster DR: **TNF concentration in fatal cerebral, non-fatal cerebral, and uncomplicated *Plasmodium falciparum* malaria.** *Lancet* 1990, **336**:1201-1204.
- Dyer M, Day KP: **Commitment to gametocytogenesis in *Plasmodium falciparum*.** *Parasitol Today* 2000, **16**:102-107.
- Boudin C, Van Der Kolk M, Tchuinkam T, Gouagna C, Bonnet S, Safeukui I, Mulder B, Meunier JY, Verhave JP: ***Plasmodium falciparum* transmission blocking immunity under conditions of low and high endemicity in Cameroon.** *Parasite Immunol* 2004, **26**:105-110.
- Paul REL, Bonnet S, Boudin C, Tchuinkam T, Robert V: **Aggregation in malaria parasites places limits on mosquito infection rates.** *Infect Genet Evol* in press.
- Davidson G, Draper CC: **Field studies of some of the basic factors concerned in the transmission of malaria.** *Trans R Soc Trop Med Hyg* 1953, **47**:522-535.
- Naotunne TS, Karunaweera ND, Del Giudice G, Kularatne MU, Grau GE, Carter R, Mendis KN: **Cytokines kill malaria parasites during infection crisis: extracellular complementary factors are essential.** *J Exp Med* 1991, **173**:523-529.
- Gouagna LC, Ferguson HM, Okech BA, Killeen GF, Kabiru EW, Beier JC, Githure JJ, Yan G: ***Plasmodium falciparum* malaria disease manifestations in humans and transmission to *Anopheles gambiae*: a field study in Western Kenya.** *Parasitology* 2004, **128**:235-243.
- Rogier C, Commenges D, Trape JF: **Evidence for an age-dependent pyrogenic threshold of *Plasmodium falciparum* parasitemia in highly endemic populations.** *Am J Trop Med Hyg* 1996, **54**:613-619.
- Trager W, Gill GS, Lawrence C, Nagel RL: ***Plasmodium falciparum*: enhanced gametocyte formation *in vitro* in reticulocyte-rich blood.** *Exp Parasitol* 1999, **91**:115-118.
- Price R, Nosten F, Simpson JA, Luxemburger C, Phaipun L, ter Kuile F, van Vugt M, Chongsuphajaisiddhi T, White NJ: **Risk factors for gametocyte carriage in uncomplicated falciparum malaria.** *Am J Trop Med Hyg* 1999, **60**:1019-1023.
- Nacher M, Singhasivanon P, Silachamroon U, Treeprasertsuk S, Tosukh Wong T, Vannaphan S, Gay F, Mazier D, Looareesuwan S: **Decreased hemoglobin concentrations, hyperparasitemia, and severe malaria are associated with increased *Plasmodium falciparum* gametocyte carriage.** *J Parasitol* 2002, **88**:97-101.

37. Ho M, White NJ, Looareesuwan S, Wattanagoon Y, Lee SH, Walport MJ, Bunnag D, Harinasuta T: **Splenic Fc receptor function in host defense and anemia in acute *Plasmodium falciparum* malaria.** *J Infect Dis* 1990, **161**:555-561.
38. Paul REL, Brey PT: **Malaria parasites and red blood cells: from anaemia to transmission.** *Mol Cells* 2003, **15**:139-149.
39. Read AF, Narara A, Nee S, Keymer AE, Day KP: **Gametocyte sex ratios as indirect measures of outcrossing rates in malaria.** *Parasitology* 1992, **104**:387-395.
40. Paul REL, Packer M, Walmsley M, Ranford-Cartwright LC, Paru R, Lagog M, Day K: **Mating patterns in malaria parasite populations of Papua New Guinea.** *Science* 1995, **269**:1709-1711.
41. Paul REL, Coulson TN, Raibaud A, Brey PT: **Sex determination in malaria parasites.** *Science* 2000, **287**:128-131.
42. West SA, Smith TG, Nee S, Read AF: **Fertility insurance and the sex ratios of malaria and related hemosporin blood parasites.** *J Parasitol* 2002, **88**:258-263.
43. Reece SE, Duncan AB, West SA, Read AF: **Host cell preference and variable transmission strategies in malaria parasites.** *Proc R Soc B-Biol Sci* 2005, **272**:511-517.
44. Boutlis CS, Yeo TV, Anstey NM: **Malaria tolerance – for whom the cell tolls?** *Trends Parasitol* 2006, **22**:371-377.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

