

Modelling of bulk density as related to aggregate size distribution in clayey Ferralsols

Nathalie Volland-Tuduri⁽¹⁾, Ary Bruand⁽¹⁾, Michel Brossard⁽²⁾, Luíz Carlos Balbino⁽³⁾, Maria Inês Lopes de Oliveira⁽⁴⁾, Éder de Souza Martins⁽⁴⁾.

¹ ISTO, Institut des Sciences de la Terre d'Orléans (ISTO), Université d'Orléans, Géosciences, BP 6759, 45067 Orléans Cedex 2, France, Nathalie.Volland@univ-orleans.fr; Ary.Bruand@univ-orleans.fr.

² IRD, BP 64501, 34394 Montpellier Cedex 5, France, brossard@ird.mpl.fr.


³ EMBRAPA Arroz e Feijão, CP 179, 75375-000, Santo Antônio de Goiás-GO, Brazil, balbino@cnpaf.embrapa.br.

⁴ EMBRAPA Cerrados, CP 08223, 73301-970 Planaltina-DF, Brazil, eder@cpac.embrapa.br.

Among microaggregated soils, there are Ferralsols that show little or no distinct horizonation. Their macrostructure is weak to moderate and they have typically a strong microstructure. In most Brazilian clayey Ferralsols, physical properties are closely related to the development of microstructure that consists of subrounded microaggregates 50 to 500 μm in diameter. They correspond to the pseudosand, micropeds, granules which were described earlier. Our objective was to show that in these soils bulk density (D_b) can be modelled according to microaggregation development and clay content.. The soils studied are located in the Central Plateau in Brazil. We measured the bulk density, aggregate size distribution, particle size distribution, and main chemical properties from the surface down to 160 cm depth in two soils under native vegetation and three others under pasture. Backscattered electron scanning images (BESI) of microaggregation were recorded using polished thin sections. BESI showed that microaggregates were either in loose assemblage or close assemblage. Aggregates size distribution enabled the measurement of the proportion of microaggregates in loose (Φ_L) and close assemblage (Φ_C) in every soil with respect to depth. Our results also showed that a linear relationship between the reciprocal of D_b and Φ_L or Φ_C can be established. Finally, in order to avoid D_b variation because of texture variation alone, a linear positive relationship between D_b and Φ_L that takes clay content into account is proposed. The latter enables the estimation of the microaggregation development using D_b as single estimator.

Startseite

EUROSOIL 2004
September, 04 - 12
Freiburg/Germany
Abstracts and Full Papers



This CD-Rom provides all abstracts, including oral and poster presentations of the Eurosoil 2004 conference. When a full paper is available, it will be indicated following the title.

Please read the impressum.

For technical requirements refer to the corresponding point in the menu.

Please choose from the menu:

technical requirements impressum abstracts by author abstracts by symposia

Impressum

Unedited Publication

The Full Papers on this CD have not been reviewed. Therefore, the scientific content of these articles is the sole responsibility of the respective authors.

Editors:

Dipl. Forstw. Nicole Wöhrle and Dipl. Forstw. Maik Scheurer
Institute of Soil Science and Forest Nutrition
Albert-Ludwigs-University Freiburg
D-79085 Freiburg i. Br.
Germany

technical requirements start page abstracts by author abstracts by symposia