

## PREVALENCE OF INTESTINAL PARASITES INCLUDING MICROSPORIDIA IN HUMAN IMMUNODEFICIENCY VIRUS-INFECTED ADULTS IN CAMEROON: A CROSS-SECTIONAL STUDY

CLAUDINE SARFATI,\* ANKE BOURGEOIS, JEAN MENOTTI, FLORIAN LIEGEOIS, ROGER MOYOU-SOMO, ERIC DELAPORTE, FRANCIS DEROUIN, EITEL MPOUDI NGOLE, AND JEAN-MICHEL MOLINA

Laboratory of Parasitology and Department of Infectious Diseases, Saint-Louis Hospital, Assistance Publique-Hopitaux de Paris, Paris, France; Institut de Recherche et Développement, Montpellier, France; Institute of Medical Research and Study of Medicinal Plants and Laboratory of Parasitology, Military Hospital, Yaounde, Cameroon

**Abstract.** To assess the prevalence of intestinal parasites in a cohort of human immunodeficiency virus (HIV)-infected adults in Cameroon, a cross-sectional study was conducted. Detection of parasites was performed in 181 stool samples from 154 HIV-infected patients with a mean CD4 cell count of 238 cells/mm<sup>3</sup>. Only 35 patients (22%) were receiving antiretroviral therapy at the time of stool sampling, and 46 (29%) had diarrhea. Opportunistic protozoa were found in 15 patients (9.7%), 8 of whom (53%) had diarrhea. *Enterocytozoon bienewisi* was found in eight patients, *C. parvum* in six patients, and *Isospora belli* in three patients. All *E. bienewisi* isolates tested belonged to the same genotype. The prevalence of opportunistic protozoa among patients with CD4 cell counts less than 50/mm<sup>3</sup> was 32%.

### INTRODUCTION

The prevalence of intestinal pathogens among human immunodeficiency virus (HIV)-infected individuals has dramatically decreased in countries where antiretroviral agents are widely available.<sup>1,2</sup> However, in most African countries, few patients have access to antiretroviral therapy. Intestinal pathogens still represent a frequent cause of diarrhea, weight loss, and wasting. Prevalence rates of intestinal pathogens, including opportunistic protozoa, in Africa vary from study to study depending on the diagnostic technique used and the study population.<sup>3–23</sup> The diagnosis of some of these pathogens is difficult and requires special staining methods<sup>24,25</sup> and trained personnel. However, application of techniques such as the polymerase chain reaction (PCR) for the detection of microsporidia in stools could increase the diagnostic sensitivity.<sup>26</sup> To assess the prevalence of intestinal parasites in Cameroon, we performed a cross-sectional study in HIV-infected adults followed in an urban center with a special focus on opportunistic protozoa.

### PATIENTS AND METHODS

This cross-sectional study was performed at the Yaounde Military Hospital in Cameroon between January 2001 and July 2003. The Cameroon National Ethics Committee reviewed and approved the study. Adult patients  $\geq$  18 years of age previously enrolled in the Projet Prevention du SIDA au Cameroun set up in 1997 in Yaounde were asked to volunteer for this study and to provide at least one stool sample for detection of ova and parasites, regardless of the presence of diarrhea, during their scheduled visit in the cohort. For each patient, data regarding age, sex, use of antiretroviral agents, and CD4 cell counts were recorded. Diarrhea was defined by laboratory staff on the basis of the appearance of stools. No bacterial cultures of stools were prepared.

For the detection of parasites, fresh stool samples were separated into three samples. The first sample was analyzed

by light microscopy in the laboratory of parasitology in Yaounde for the detection of helminths and protozoa by use of both direct and formalin/ether (10%) concentration procedures. The Baermann technique, a larval concentration technique and an alternative to the charcoal culture, was also performed for the detection of *Strongyloides stercoralis* larvae. This sample was also analyzed by modified trichrome (Meridian, Cincinnati, OH) and fluorochrome staining methods for detection of microsporidial spores,<sup>24,25</sup> and by modified Ziehl-Neelsen staining for the detection of *Cryptosporidium* sp. and *Isospora belli*.<sup>27</sup>

The second sample (in 10% formalin) as well as stained slides, were shipped to France for confirmation of the results obtained in Yaounde.

The third sample was diluted in potassium bichromate and used for the detection of microsporidia by a PCR. The DNA was extracted as previously described.<sup>26</sup> Polymerase chain reaction amplification of the internal transcribed spacer of the *E. bienewisi* rRNA gene was performed as described by Velasquez and others.<sup>28</sup> Genotyping of *E. bienewisi* strains isolated from stool specimens was performed by digestion of the amplification products with *Nla* III and *Fnu* 4HI endonucleases (New England Biolabs, Beverly, MA) as described by Liguori and others.<sup>29</sup>

### RESULTS

One hundred eighty-one stool samples from 154 patients were obtained during the study period. Only 57% (154 of 270) of the patients in the cohort agreed to participate in the study. The mean age of the patients was 36 years (range = 23–58), 63% were women, and their mean CD4 cell count was 238 cells/mm<sup>3</sup>. Twenty-two patients (14%) were severely immunosuppressed and had less than 50 CD4 cells/mm<sup>3</sup>. Only 35 patients (22%) were receiving antiretroviral therapy at the time of the study. Forty-six patients (29%) had diarrhea.

Intestinal parasites were found in 51 (33%) of the 154 patients studied. Among the 85 parasites identified, 42 were pathogenic. Helminths were identified in 19 patients, of whom 5 had *S. stercoralis* larvae. Non-opportunistic protozoa, mainly non-pathogenic amoeba, were identified in 28 patients. Seven patients had *Entamoeba histolytica*/*E. dispar* and one

\* Address correspondence to Claudine Sarfati, Laboratory of Parasitology, Saint-Louis Hospital, 1 avenue Claude Vellefaux 75010, Paris, France. E-mail: claudine.sarfati@sls.ap-hop-paris.fr

patient had *Giardia intestinalis*. *Cyclospora cayetanensis* was not detected in any of the 181 stool samples examined.

Opportunistic protozoa including *C. parvum*, microsporidia, and *I. belli* (Table 1) were found in 15 patients (9.7%), 8 of whom (53%) had diarrhea. The median CD4 cell count of these patients with opportunistic protozoa was 72 cells/mm<sup>3</sup> (range = 4–246). All three patients with *I. belli* spores had diarrhea. *Cryptosporidium parvum* oocysts were found in six patients, and direct microscopy detected microsporidia spores in the stools of six patients. The PCR detected *E. bienewisi* DNA in the stools of two other patients in whom no spores could be seen by direct microscopy. Neither of these two patients had diarrhea. Four of the *E. bienewisi* strains isolated in this study showed genotype IV, which is a rare genotype in France.<sup>29</sup>

Overall, among the 22 patients in this study who had less than 50 CD4 cells/mm<sup>3</sup>, 7 (32%) were infected with either *E. bienewisi* or *C. parvum*. Only three patients receiving antiretroviral therapy had opportunistic parasites identified in stools (*Microsporidia*, *Cryptosporidium*, and *I. belli*, respectively). However, the remaining 32 patients receiving antiretroviral therapy were free of intestinal parasites.

DISCUSSION

In this cross-sectional study of HIV-infected adults in Cameroon, the overall prevalence of intestinal parasites in stool samples was 33%. This prevalence is lower than those previously reported in other studies conducted in Africa.<sup>12,19,30</sup> This low prevalence could be related to the location in which our study was conducted, an urban center, and to the population studied, adults not selected for the presence of diarrhea. Only 29% of our patients had diarrhea at the time of stool examination. Furthermore, since no bacterial cultures of stools were prepared in this study, any association between parasites and diarrhea in this study should be made cautiously.

Various intestinal parasites were found in the stools of our patients. Pathogenic helminths were found in 12% of the patients, a prevalence similar to that reported in other studies of HIV-infected patients in Cameroon and Mali.<sup>23,30</sup> Only seven (4.5%) patients had *E. histolytica*/*E. dispar* cysts in their stools, a prevalence that was lower than the prevalence rate of 12% reported in Cameroon in patients with diarrhea.<sup>23</sup>

Our study focused mainly on opportunistic protozoa, a frequent cause of chronic diarrhea and wasting in HIV-infected patients in developing countries in Africa.<sup>3–23,31</sup> *Cryptosporidium parvum*, microsporidia, and *I. belli* accounted for the most of these opportunistic protozoa. Their diagnosis is dif-

ficult because of the special staining methods required. In this study, where only 22% of the patients were receiving antiretroviral therapy, the prevalence of opportunistic protozoa was 9.7%. However, among patients with less than 50 CD4 cells/mm<sup>3</sup>, this prevalence increased to 32%. Previous studies in Africa have found prevalence rates ranging from 3.5% to 28%, but were conducted mainly in patients with diarrhea.<sup>3–15</sup>

*Cryptosporidium parvum* oocysts were found in 3.9% of the patients, half of whom had diarrhea. This was in the range of the prevalence of 8.7% reported in Côte d'Ivoire by Assoumou and others<sup>20</sup> among 217 patients who were not selected for diarrhea. However, higher prevalence rates up to 25% have been reported in other studies usually conducted in patients with diarrhea. The low prevalence reported in our study is not unexpected because the patient population was not selected on the basis of gastrointestinal symptoms.

Microsporidia, which were identified by PCR as *E. bienewisi*, were detected in eight patients. The prevalence reported in our study (5.2%) was lower than those reported in previous studies.<sup>8,10,12,14–16,18,19,23</sup> Less than half of our patients with microsporidia had diarrhea. This suggests that either a significant proportion of our patients were asymptomatic carriers of this opportunistic protozoa, a condition previously described in patients with high CD4 cell counts, or that they had other clinical symptoms frequently associated with microsporidia such as cachexia, weight loss, and malabsorption. Genotype IV of *E. bienewisi* identified in this study was rarely detected in a previous study in France, suggesting differences in *E. bienewisi* strain distribution.<sup>29</sup> Moreover, genotype IV corresponds to genotype K described by Dengjel and others<sup>32</sup> from a cat sample. Genotype K was found in only one human case in Europe,<sup>33</sup> but in 6 of 10 sequences from human cases in Uganda.<sup>34</sup> However, further studies that test a larger number of *E. bienewisi* strains from Africa are needed to confirm these findings.

In conclusion, in this cross-sectional study of 154 HIV-infected adults in Yaounde, the overall prevalence of intestinal parasites was 33%, and that of opportunistic protozoa was 9.7%. However, the prevalence of opportunistic protozoa increased to 32% in patients with less than 50 CD4 cells/mm<sup>3</sup>. The results of our study should therefore prompt physicians caring for HIV-infected patients in Cameroon to request stool examination and specific tests for microsporidia, *Cryptosporidium*, and *Isospora*, especially in patients with low CD4 cell counts.

Received February 22, 2005. Accepted for publication September 1, 2005.

Acknowledgments: We thank Alphonse Poupou for technical assistance in Cameroon, Samia Hamane for expert advice, and Corinne Delalande, Marie-Christine Serugue, and Caroline Verret for technical assistance in Paris.

Financial support: This study was supported by a grant from the French Ministry of Research and Agence Nationale de Recherches sur le SIDA (ANRS)—Program VIH PAL.

Authors' addresses: Claudine Sarfati, Jean Menotti, and Francis Derouin, Laboratory of Parasitology, Saint-Louis Hospital, 1 Avenue Claude Vellefaux 75010, Paris, France, E-mails: claudine.sarfati@sls.ap-hop-paris.fr, jean.menotti@slc.aphp.fr, and francis.derouin@sls.aphp.fr. Anke Bourgeois, Florian Liegeois, and Eric Delaporte, UR 36 IRD/Département Universitaire Maladies Infectieuses et Santé Internationale, 911 Avenue Agropolis, BP 64501, 34394 Montpellier Cedex 5 France, E-mails: anke.bourgeois@mpl.ird.fr,

TABLE 1

Opportunistic protozoa (OP) identified in 154 human immunodeficiency virus-infected adults in Yaounde, Cameroon

	No. of patients (%) <sup>*</sup>	No. of patients (%) without diarrhea	No. of patients (%) with diarrhea
Total	154	108 (70)	46/154 (30)
Patients with OP	15 (9.7)	7/108 (6.4)	8/46 (17.3)
Cryptosporidia	6 (3.8)	3/108 (2.8)	3/46 (6.5)
Microsporidia	8 (5.2)	5/108 (4.6)	3/46 (6.5)
<i>Isospora belli</i>	3 (2)	0/108 (0)	3/46 (6.5)

<sup>\*</sup> One patient can have more than one pathogen.

florian.liegeois@mpl.ird.fr, and eric.delaporte@mpl.ird.fr. Roger Moyou-Somo and Eitel Mpouidi Ngole, Hôpital Militaire, Yaounde, Cameroon. Jean-Michel Molina, Department of Infectious Diseases, Saint-Louis Hospital, 1 Avenue Claude Vellefaux 75010, Paris, France, E-mail: jean-michel.molina@sls.aphp.fr.

## REFERENCES

1. Le Moing V, Bissuel F, Costagliola D, Eid Z, Chapuis F, Molina JM, Salmon-Ceron D, Brasseur P, Lepout C, 1998. Decreased prevalence of intestinal cryptosporidiosis in HIV-infected patients concomitant to the widespread use of protease inhibitors. *AIDS* 12: 1395-1397.
2. Carr A, Marriott D, Field A, Vasak E, Cooper DA, 1998. Treatment of HIV-1-associated microsporidiosis and cryptosporidiosis with combination antiretroviral therapy. *Lancet* 351: 256-261.
3. Fisseha B, Petros B, Wolde Michael T, 1998. Cryptosporidium and others parasites in Ethiopian AIDS patients with chronic diarrhoea. *East Afr Med J* 75: 100-101.
4. Mengesha B, 1994. Cryptosporidiosis among medical patients with the acquired immunodeficiency syndrome in Tikur Anbessa teaching Hospital, Ethiopia. *East Afr Med J* 71: 376-378.
5. Mwachari C, Batchelor BI, Paul J, Waiyaki PG, Gilks CF, 1998. Chronic diarrhoea among HIV-infected adult patients in Nairobi, Kenya. *J Infect* 37: 48-53.
6. Ravera M, Reggioni A, Riccioni G, 1996. Prevalence of *Isospora belli* in AIDS and immunocompetent patients in Uganda. *Presse Med* 25: 1170.
7. Cegielski JP, Ortega YR, McKee S, Madden JF, Gaido L, Schwartz DA, Manji K, Jorgensen AF, Miller SE, Pulipaka UP, Msengi AE, Mwakyusa DH, Sterling CR, Reller LB, 1999. *Cryptosporidium*, *Enterocytozoon*, and *Cyclospora* infections in pediatric and adult patients with diarrhea in Tanzania. *Clin Infect Dis* 28: 314-321.
8. Kadende P, Nkurunziza T, Floch JJ, Mpizzi B, Laroche R, Ndebanezo Aubry P, 1989. Infectious diarrhea in african acquired immunodeficiency syndrome AIDS. A propos of 100 patients studied in Bujumbura (Burundi). *Med Trop (Mars)* 49: 129-213.
9. Cranendonk RJ, Kodde CJ, Chipeta D, Zijlstra EE, Sluiter JF, 2003. *Cryptosporidium parvum* and *Isospora belli* infections among patients with and without diarrhoea. *East Afr Med J* 80: 398-401.
10. Drobniewski F, Kelly P, Carew A, Ngwenya B, Luo N, Pankhurst C, Farthing M, 1995. Human microsporidiosis in African AIDS patients with chronic diarrhea. *J Infect Dis* 171: 515-516.
11. Hunter G, Bagshawe AF, Baboo KS, Luke R, Prociw P, 1992. Intestinal parasites in Zambian patients with AIDS. *Trans R Soc Trop Med Hyg* 86: 543-545.
12. Kelly P, Davies SE, Mandanda B, Veitch A, McPhail G, Zulu I, Drobniewski F, Fuchs D, Summerbell C, Luo NP, Pobee JO, Farthing MJ, 1997. Enteropathy in Zambians with HIV related diarrhoea: regression modelling of potential determinants of mucosal damage. *Gut* 41: 811-816.
13. Coulon CP, Pinching AJ, Perera CU, Moody A, Luo NP, Lucas SB, 1990. HIV-related enteropathy in Zambia: a clinical, microbiological, and histological study. *Am J Trop Med Hyg* 42: 82-88.
14. Gumbo T, Sarbah S, Gangaidzo IT, Ortega Y, Sterling CR, Carville A, Tzipori S, Wiest PM, 1999. Intestinal parasites in patients with diarrhea and human immunodeficiency virus infection in Zimbabwe. *AIDS* 13: 819-821.
15. van Gool T, Luderhoff E, Nathoo KJ, Kiire CF, Dankert J, Mason PR, 1995. High prevalence of *Enterocytozoon bienewsi* infections among HIV-positive individuals with persistent diarrhoea in Harare, Zimbabwe. *Trans R Soc Trop Med Hyg* 89: 478-480.
16. Maiga I, Doumbo O, Dembele M, Traore H, Desportes-Livage I, Hilmarsdottir I, Giboyau E, Maiga L, Kassambara L, el Fakhry Y, Dattray A, Gentilini M, Pichard E, 1997. Human intestinal microsporidiosis in Bamako (Mali): the presence of *Enterocytozoon bienewsi* in HIV seropositive patients. *Sante* 7: 257-262.
17. Dieng T, Ndir O, Diallo S, Coll-Seck AM, Dieng Y, 1994. Prevalence of *Cryptosporidium sp* and *Isospora belli* in patients with acquired immunodeficiency syndrome (AIDS) in Dakar (Senegal). *Dakar Med* 39: 121-124.
18. Dieng Y, Dieng TH, Diouf G, Coll-Seck AM, Diallo S, 1998. Recherche des spores de microsporidies chez des patients sidéens au CHU de Fann à Dakar (Sénégal). Résultats préliminaires. *Med Mal Infect* 28: 265-267.
19. Lebbad M, Norrgren H, Naucler A, Dias F, Andersson S, Linder E, 2001. Intestinal parasites in HIV-2 associated AIDS cases with chronic diarrhoea in Guinea-Bissau. *Acta Trop* 80: 45-49.
20. Assoumou A, Kone M, Penali LK, Coulibaly M, N'Draman AA, 1993. Cryptosporidiosis and HIV in Abidjan (Ivory Coast). *Bull Soc Pathol Exot* 86: 85-86.
21. Therizol-Ferly PM, Tagliente-Saracino J, Kone M, Konan A, Ouhon J, Aassoumou A, Aka K, Aasse G, 1989. Chronic diarrhea and parasitoses in adults suspected of AIDS in the Ivory Coast. *Bull Soc Pathol Exot Filiales* 82: 690-693.
22. Nwokediuko SC, Bojuwoye BJ, Onyenekwe B, 2002. Apparent rarity of cryptosporidiosis in human immunodeficiency virus (HIV)-related diarrhea in Enugu, south eastern, Nigeria. *Niger Postgrad Med J* 9: 70-73.
23. Same-Ekobo A, Lohoue J, Mbassi A, 1997. A clinical and biological study of parasitic and fungal diarrhea in immunosuppressed patients in an urban and suburban area of Yaounde. *Sante* 7: 349-354.
24. Weber R, Bryan RT, Owen RL, Wilcox CM, Gorelkin L, Visvesvara GS, 1992. Improved light-microscopical detection of microsporidia spores in stool and duodenal aspirates. *N Engl J Med* 326: 161-166.
25. van Gool T, Snijders F, Reiss P, Eeftinck Schattenkerk JK, van den Bergh Weerman MA, Bartelsman JF, Bruins JJ, Canning EU, Dankert J, 1993. Diagnosis of intestinal and disseminated microsporidial infections in patients with HIV by a new rapid fluorescence technique. *J Clin Pathol* 46: 694-699.
26. Liguory O, David F, Sarfati C, Schuitema AR, Hartskeerl RA, Derouin F, Modai J, Molina JM, 1997. Diagnosis of infections caused by *Enterocytozoon bienewsi* and *Encephalitozoon intestinalis* using polymerase chain reaction in stool specimens. *AIDS* 11: 723-726.
27. Henriksen S, Pohlenz J, 1981. Staining of *Cryptosporidia* by a modified Ziehl-Neelsen technique. *Acta Vet Scand* 22: 594-596.
28. Velasquez JN, Carnevale S, Guarnera EA, Labbe JH, Chertcoff A, Cabrera MG, Rodriguez MI, 1996. Detection of the microsporidian parasite *Enterocytozoon bienewsi* in specimens from patients with AIDS by PCR. *J Clin Microbiol* 34: 3230-3232.
29. Liguory O, David F, Sarfati C, Derouin F, Molina JM, 1998. Determination of types of *Enterocytozoon bienewsi* strains isolated from patients with intestinal microsporidiosis. *J Clin Microbiol* 36: 1882-1885.
30. Dolo A, Cancrini C, Traore F, Traore S, Kassambara L, Diakite M, Camara F, Doumbo O, 1996. Protozoan infections and intestinal helminthiasis among the population of a village in the northern Sudan savannah area of Mali. *Parasitologia* 38: 585-589.
31. Molina JM, Sarfati C, Beauvais B, Lemann M, Lesourd A, Ferchal F, Casin I, Lagrange P, Modigliani R, Derouin F, Modai J, 1993. Intestinal microsporidiosis in human immunodeficiency virus-infected patients with chronic unexplained diarrhea: prevalence and clinical and biologic features. *J Infect Dis* 167: 217-221.
32. Dengjel B, Zahler M, Hermanns W, Heinritzi K, Spillmann T, Thomschke A, Loscher T, Gothe R, Rinder H, 2001. Zoonotic potential of *Enterocytozoon bienewsi*. *J Clin Microbiol* 39: 4495-4499.
33. Sadler F, Peake N, Borrow R, Rowl PL, Wilkins EG, Curry A, 2002. Genotyping of *Enterocytozoon bienewsi* in AIDS patients from the north west of England. *J Infect* 44: 39-42.
34. Tumwine JK, Kekitiinwa A, Nabukeera N, Akiyoshi DE, Rich SM, Widmer G, Feng X, Tzipori S, 2003. *Cryptosporidium parvum* in children with diarrhea in Mulago Hospital, Kampala, Uganda. *Am J Trop Med Hyg* 68: 710-715.

Sarfati C., Bourgeois Anke, Menotti J., Liégeois Florian,  
Moyou-Somo R., Delaporte Eric, Derouin F., Mpoudi Ngole E.,  
Molina J.M. (2006)

Prevalence of intestinal parasites including microsporidia in  
human immunodeficiency virus-infected adults in Cameroon  
: a cross-sectional study

American Journal of Tropical Medicine and Hygiene, 74 (1),  
162-164

ISSN 0002-9637