

Medium-term and long-term *in vitro* conservation and safe international exchange of yam (*Dioscorea* spp.) germplasm

Bernard Malaurie¹

GeneTrop – Unité de Génétique et d'Amélioration des Plantes - Centre ORSTOM** – 911 avenue Agropolis - BP. 5045 -F. 34032. Montpellier Cedex 1, France.

Tel: Bureau (33)-- 4 67 41 62 44

Tel: Standard (33)-- 4 67 41 61 00

Fax: (33)-- 4 67 54 78 00

E-mail : Bernard.Malaurie@mpl.orstom.fr

Marie-France Trouslot

GeneTrop – Unité de Génétique et d'Amélioration des Plantes - Centre ORSTOM** – 911 avenue Agropolis - BP. 5045 -F. 34032. Montpellier Cedex 1, France. E-mail : Bernard.Malaurie@mpl.orstom.fr

Julien Berthaud

GeneTrop – Unité de Génétique et d'Amélioration des Plantes - Centre ORSTOM** – 911 avenue Agropolis - BP. 5045 -F. 34032. Montpellier Cedex 1, France. E-mail : Bernard.Malaurie@mpl.orstom.fr

Mustapha Bousalem

Laboratoire de Phytovirologie des Régions Chaudes (LPRC) – CIRAD-ORSTOM** – avenue du val de Montferrand - BP. 5035 -F. 34032 - Montpellier Cedex 1, France. E-mail : dubern@melusine.mpl.orstom.fr

Agnès Pinel

Laboratoire de Phytovirologie des Régions Chaudes (LPRC) – CIRAD-ORSTOM** – avenue du val de Montferrand - BP. 5035 -F. 34032 - Montpellier Cedex 1, France. E-mail : dubern@melusine.mpl.orstom.fr

Jean Dubern

Laboratoire de Phytovirologie des Régions Chaudes (LPRC) – CIRAD-ORSTOM** – avenue du val de Montferrand - BP. 5035 -F. 34032 - Montpellier Cedex 1, France. E-mail : dubern@melusine.mpl.orstom.fr

<http://www.mpl.orstom.fr>

Yam edible tubers feed million of peoples in the intertropical area, where they represent 12% of human feeding. However, as a vegetatively propagated crop, yam is seriously affected by an accumulation of pathogens. Establishing *in vitro* germplasm collection is a process that cleans the plants from all diseases but viruses. It gives a good control on the preservation of the yam genetic resources and facilitates international exchanges of healthy plant material.

Two kinds of *in vitro* germplasm preservation were considered: slow growth condition culture for mid-term preservation, and cryopreservation using the encapsulation/dehydration technique for long-term preservation. Virus eradication was approached by meristem culture and chemo and thermotherapy. Production of virus-free plants was controlled by ELISA.

We succeeded in the introduction and maintenance of 20 yam species, under slow growth conditions. Cryopreservation was applied successfully on two edible yam species, *Dioscorea. alata* L and *D. bulbifera* L. Virus-free plants were obtained by meristem culture in *D. cayenensis-D. rotundata* complex and *D. praeheensis*. Indexation allowed the detection of different virus (poty-, potex-, badna- and cucumovirus), where the most important potyvirus was YMV.

Mid-term conservation of yam germplasm is used routinely, and from these conditions a direct acclimatization is possible. On the cryopreservation aspect, experiments are under way to apply the optimized protocol to genotypes which are more representative of the diversity, to insure a routinely use. More work can be conducted now on virus eradication,

¹ Corresponding author

** ORSTOM is now called IRD, Institut de Recherche pour le Développement.

based on knowledge accumulated on potyvirus diversity, on several tests available for yam indexing (ELISA, rt/PCR, monoclonal antibodies) and on new sanitation techniques.

Yam belongs to *Dioscorea* genus which has more than 600 species (Coursey, 1967) most of them distributed in the intertropical humid area. We will distinguish two types of yam: 1) medicinal yams, 2) edible yams and relatives. Medicinal yams concern about fifty species characterized by their sapogenin content, which are steroidal components. For the edible yams and relatives, we will observe two

groups: 1) domesticated species, 2) and wild species.

For domesticated species we consider that forty to fifty species are occasionally used (Martin and Degras, 1978). From these only eleven are cultivated (Table 1). From the 11 cultivated, 6 represent an important part of feeding (*D. alata*, *D. cayenensis*-*D. rotundata* complex, *D. bulbifera*, *D. dumetorum*, *D. esculenta*, *D. trifida*), 3 are scattered in all the intertropical humid area (*D. alata*, *D. bulbifera*, *D. esculenta*), and yam belonging to the *D. cayenensis*-*D. rotundata* complex take place most of them in West Africa and some in the Caribbean area. The other yams are cultivated in their origin area (Degras, 1986).

Table 1 : Main edible species of yam

Species ¹	Zone of origin	Zone of culture
Enantiophyllum Section		
<i>D. alata</i> L.	South East Asia	Inter-tropical humid
<i>D. cayenensis</i> Lamk.	West Africa	West & Central Africa, and Caribbean
<i>D. rotundata</i> Poir. complex ²		
<i>D. nummularia</i> Lamk.	Indonesia, Oceania	Indonesia, Oceania and Micronesia
<i>D. opposita</i> Thunb.	Temperate area from:	Temperate area from:
<i>D. japonica</i> Thunb. complex ³	China, Korea, Taiwan	China, Korea, Taiwan
	Japan	Japan
<i>D. transversa</i> Br.	South Pacific	South Pacific
Lasiophyton Section		
<i>D. dumetorum</i> (Kunth) Pax.	West Africa	West Africa
<i>D. hispida</i> Dennst.	India, South-China, New Guinea	India, South-China, New Guinea
<i>D. pentaphylla</i> L.	Himalaya and Oceania	Himalaya and Oceania
Combilium Section		
<i>D. esculenta</i> (Lour.) Burk.	South East Asia	Inter-tropical humid
Opsophyton Section		
<i>D. bulbifera</i> L.	South East Asia and Africa	Inter-tropical humid
Macrogynodium Section		
<i>D. trifida</i> L.	Guyana, Amazonian basin	Caribbean

Sources: Malaurie et al. (1998a)

¹ Species have been regrouped in Section by Knuth (1924), completed by Burkill (1960)

² Grouping together species of *D. cayenensis* and *D. rotundata* in a Complex has been proposed by Ayensu and Coursey (1972), Martin and Rhodes (1978), Miège (1982)

³ Grouping together species of *D. opposita* and *D. japonica* in a Complex has been proposed by Tanaka (1977).

Main edible species of yams are: 1) native of a continent, 2) and cultivated in the same continent, 3) or/and cultivated in an other. This observation implies very strong links to exchange problems. In Table 2, 36 countries have been

observed by IBPGR in 1986 with *Dioscorea* germplasm. These countries are supposed to be concerned by an international exchange of yam germplasm. Some of them,

have, in our knowledge, already developed *in vitro* germplasm collection.

Table 2 : Countries¹ and geographic zones where yam collections have been observed

Europe	West Indies	America	Pacific	Asia	Africa
France ²	Barbados *	Brazil ²	Cook Islands	Bengladesh	Bénin
United Kingdom ²	Cuba	Colombia	Fiji	India	Burkina Faso
	Guadeloupe ²	Costa Rica	Niue Islands	Indonesia	Cameroun
	Jamaïca	Guatemala	New Calédoniace ²	Japan ²	Ivory Coast ²
	Saint-Dominique	Mexico	Papua NewGuinea	Malaisia	Ghana
	Trinidad-Tobago	Panama	Salomon Islands	Nepal	Nigeria ²
	USA	Tonga	Philippines	South Africa	
		Vanuatu	Sri Lanka	Togo	
		Western Samoa ²	Thailand	Uganda	
			Viet Nam		

(Sources : IBPGR 1986, FAO 1996, Malaurie et al 1998a)

* *in vitro* maintenance for production purpose

¹ This country listing is not exhaustive, and take into account only sources in our possession

² Countries with *in vitro* collection (according to sources in our possession)

Different genebank preservation levels exist. A first group concerns non aseptic germplasm conservation with in field genebank and seed genebank, where important disadvantages and heavy constraint of quarantine measures explain the choice of *in vitro* germplasm conservation (Hanson, 1986; Malaurie et al., 1998a) (Table 3).

Table 3: Non aseptic germplasm conservation

	Non aseptic Genebanks	
	In field Genebanks	Seed Genebanks
Disadvantages		
Genetic erosion	+++	
Expensive	+++	
Hard to manage	+++	
Do not bread true		+++
Tuber shape		+++
Dormancy		+++

× a. conservation

Three levels of *in vitro* genebank preservation levels could be considered: 1) short term conservation: this conservation

under normal growth conditions is suitable for temporary storage of germplasm collections, and for international distribution, 2) medium term conservation, which could be considered as an active *in vitro* genebank, 3) long term conservation, considered also as a base *in vitro* genebank.

These *in vitro* genebanks have been previously introduced *in vitro* from tuber or seed. These introductions have to be linked to an obligatory phytosanitary control from mother plants and from *in vitro* material after introduction. Medium term conservation, which correspond to *in vitro* culture under slow growth conditions, could be obtained by several ways: 1) physiological stage of the explant, 2) addition of osmotic agents and growth moderators, 3) low storage temperature, 4) low mineral or sucrose concentrations, 5) low oxygen pressure, 6) encapsulation in alginate (Charrier et al., 1991; Withers, 1991; Engelmann, 1991; Malaurie et al., 1998a).

Medium-term conservation

At ORSTOM, we choose to maintain the *in vitro* yam

collection in a medium with low mineral nutrient and a low sucrose concentration. We succeeded in the introduction and maintenance of 14 species of yam (Malaurie et al., 1993). Since this time, this collection is continuously enriched by new genotypes and comprises 20 species (Table 4).

For yam, this simple solution of slow growth is used routinely and from these culture conditions a direct acclimatization is possible. This mode of conservation allows an international distribution of the material and corresponds to an active genebank (Malaurie et al. 1993, 1998c, Malaurie and Trouslot 1995c).

This *in vitro* germplasm collection of yam is maintained in test tubes, at ORSTOM (Montpellier, France), with a total of 6 test tubes by accession, with two different places of storage for the replicates; the minimal growth conditions allow to maintain most of the accessions up to 2 years. Technical constraints in the collection management lead to subculture the accessions every 6-8 months (Malaurie et al., 1998c).

Table 4 : Listing of different species of yam maintained in an ² a. collection, under slow growth culture condition *
(GeneTrop, GAP unit, ORSTOM, Montpellier, France)

Species	Number of accessions
<i>D. abyssinica</i> Hochst. Ex Kunth	6
<i>D. alata</i> L.	91
<i>D. bulbifera</i> L.	8
<i>D. cayenensis</i> Lamk.	63 (+ 17)
<i>D. rotundata</i> Poir. complex	
<i>D. burkilliana</i> J. Miège	11
<i>D. dumetorum</i> (Kunth) Pax.	2
<i>D. esculenta</i> (Lour.) Burk.	10
<i>D. hirtiflora</i> Benth.	1
<i>D. mangelotiana</i> J. Miège	15
<i>D. minutiflora</i> Engl.	2
<i>D. opposita</i> Thunb.	1
<i>D. japonica</i> Thunb. complex	
<i>D. praehensilis</i> Benth.	17
<i>D. preussii</i> Pax	1
<i>D. sansibarensis</i> Pax	1
<i>D. schimperana</i> Hochst. Ex Kunth	1
<i>D. smilacifolia</i> De Wild	2
<i>D. togoensis</i> Knuth	8
<i>D. transversa</i> Br.	1
<i>D. trifida</i> L.	2 (+ 1)
Interspecific Hybrids: <i>D. cayenensis</i> - <i>D. rotundata</i> complex cv. 'Krengle' X <i>D. praehensilis</i> so-called 'Ignose de Pilimpikou'	14 (+ 9)

* (+): Accessions recently introduced

Different species maintained in the *in vitro* collection, such as *D. cayenensis*-*D. rotundata* complex are going to be enriched by cultivars from Burkina Faso for a sanitation program, and from Benin for a genetic program linked to virologic aspect. Others species supposed to be links to *D. cayenensis*-*D. rotundata* complex, such as *D. mangenotiana*, *D. praehensilis*, *D. minutiflora*, or *D. abyssinica*, *D. praehensilis*, are going to be enriched by further introduction.

Orstom virologists are interested by *D. trifida* because of its strong sensibility to virus, which provoked in Guadeloupe, French West-Indies, its quite disappearance. Serological and molecular works are developed to explain this virus sensibility.

Long-term conservation

Long term conservation correspond to cryopreservation in liquid nitrogen, at -196°C . Plant cryobiology, which begun in 1971 by Latta works on carrot cell suspension, benefited from results on animals cell by Polge et al in 1949. Since these dates, different techniques have been set up: 1) on one hand, the so-called conventional techniques, using two steps of slow freezing, with the addition of cryo-protector (Sakai, 1984), 2) and on the other hand, new techniques, characterized by a very rapid freezing, about $1000^{\circ}\text{C}/\text{min}$, by direct immersion in liquid nitrogen (Table 5) (Dereuddre et al., 1990, 1991; Tannoury et al., 1991; Uragami, 1993). The aim of these techniques is to try to control water flow and ice formation, and tend to a vitrified state, avoiding crystal formation during thawing, and to protect the cell from thermic shocks.

Table 5 : Long term conservation : cryopreservation in liquid nitrogen, -196°C

Steps	Conventional techniques	New techniques		
		Air-drying	Vitrification	Encapsulation / Dehydration
Encapsulation				+
Sucrose pretreatment	+/-	+ (+ABA)		+
Cryoprotector	+		++++	
Desiccation		+		+
Slow-freezing	+ 0°C to -40°C (0.3 to $1^{\circ}\text{C}/\text{min}$)			
Rapid-freezing	+ -40°C to -196°C ($200^{\circ}\text{C}/\text{min}$)	+ $+25^{\circ}\text{C}$ to -196°C ($720^{\circ}\text{C}/\text{min}$)	+ $+25^{\circ}\text{C}$ to -196°C (400 to $1100^{\circ}\text{C}/\text{min}$)	+ $+25^{\circ}\text{C}$ to -196°C ($720^{\circ}\text{C}/\text{min}$)
Thawing	$500^{\circ}\text{C}/\text{min}$	$120^{\circ}\text{C}/\text{min}$	$120^{\circ}\text{C}/\text{min}$	$120^{\circ}\text{C}/\text{min}$

Sources : Uragami (1993), Malaurie et al. (1998a).

Most of the results about cryopreservation have been obtained from conventional techniques on suspension cells of medicinal yam, *D. deltoidea* being the most used (Butenko et al., 1984; Popov and Fedorovskii, 1992; Popov and Volkova, 1994). More recent works have been done on rapid cryopreservation of callus (Chulafich et al., 1994), by

direct immersion in liquid nitrogen, of two other medicinal yams (*D. balcanica*, *D. caucasica*).

Since 1996, new results have been obtained by two different research teams, using encapsulation/dehydration of shoot apices. On the one hand, Mandal *et al.* (1996) compared the survival capacities of apices after the osmotic

and thermic stress of the technique of four species of yam - three edible (*D. alata*, *D. bulbifera*, *D. wallichii*), and one medicinal (*D. floribunda*). Four of them have survived after immersion in liquid nitrogen, with 26 to 71%, depending on the species. Meanwhile, only two of them (*D. alata*, *D. wallichii*) allowed the recovery into shoots after immersion in liquid nitrogen, with 21 and 37%, respectively.

On the other hand, ORSTOM ability in different aspects of the long term conservation on tropical plants (Engelmann, 1991), and on encapsulation/dehydration technique applied on coffee, cassava, oil palm...etc, permitted to apply the process on apical shoot-tips of *in vitro* plantlets of yam (Malaurie and Trouslot 1996). Malaurie et al. (1998b) obtained survival rates over 50% for the two species (*D. alata*, *D. bulbifera*), and recovery to rooted leafy shoots after immersion in liquid nitrogen of at least 60% for *D. bulbifera* and 20% for *D. alata*, three months culture after

thawing.

Comparatively to previous works on cryopreservation using encapsulation/dehydration technique, Malaurie et al. (1998b) have used higher sucrose concentration (0.9, 1.0 and 1.1M), a wider range of dehydration duration, up to 23h and a new and more accurate method for measuring of dry weight.

The new and more accurate method for measuring of dry weight used in our experiments consisted of desiccating alginate beads for 30d in airtight boxes containing dry silica gel, to avoid mass loss due to caramelization of sugar when drying at a temperature higher than 100°C. We obtained a strong linear correlation between dry mass (DW_{30}) and sucrose molarity for sucrose-pretreated alginate beads. During the whole experiment, we used DW_{30} values estimated by linear regression (Table 6).

Table 6: Dry mass and water content of sucrose-pretreated alginate beads, determined after 30d of drying with silica gel in airtight boxes at room temperature ⁽¹⁾.

Sucrose concentration	DW_{30} (% FW) estimated by linear regression ⁽²⁾	Water content before dehydration ($g \cdot g^{-1}$ DW)
0.75M	28.8	2.47
0.9M	33.3	2.00
1.0M	36.3	1.76
1.1M	39.3	1.54

(1) Source: Malaurie et al. (1998b)

(2) >From mean values over 13 to 15 replicates for each of the four sucrose concentrations ($y = 6.4319 + 29.872x$; $N = 4$; $r = 0.999$). Similar results were obtained from replicate data ($y = 6.4177 + 29.883x$; $N = 55$; $r = 0.960$). Data not shown.

For the best sucrose pretreatments depending species, figure 1 (Malaurie et al., 1998b) shows that *D. bulbifera* still has high survival with high sucrose concentration and after long duration dehydration (up to 23h). For the two species, the water content of encapsulated apices had to be decreased down to $0.15g \text{ H}_2\text{O} \cdot g^{-1} \text{ DW}$ in order to obtain high survival after freezing. The percentage of water loss was of 67, 62, 58 and 55% FW ($\pm 1\%$) for 0.75, 0.9, 1 and 1.1M sucrose pretreatments, respectively. Our results

demonstrated that, in most cases, survival increased when dehydration was extended to a defined threshold, around $0.13-0.15g \cdot g^{-1} \text{ DW}$, which was obtained after desiccation periods from 10 to 18h. It seemed that, with this soft dessication process, we could rub out differences in residual water-free, which certainly exist between apices from a same plot.




Figure 1. Effect of dehydration duration on the water content of beads and survival rate of control (-LN □) and cryopreserved (+LN ■) encapsulated apices of *D. alata* 'Brazo Fuerte' after pretreatment with sucrose (0.9M+1M)- (A), and *D. bulbifera* 'Nouméa Imboro' after pretreatment with sucrose (1M+1.1M)- pretreated ones (B). Each point corresponds to a mean over 2 sucrose concentrations and all the 6 or 8 pretreatment durations used depending on the clone (Malaurie et al., 1998b).

Indexation and Disease-free germplasm production

Indexation

In vitro germplasm conservation presents different advantages such as: 1) to be free from genetic erosion, 2) to have the possibility for the establishment of core collection with long term genebanks, 3) to be free from fungus and bacteria, 4) to be not expensive, when *in vitro* facilities are already present, 5) easy and convenient for international

distribution. But International exchanges need more for safe international exchange. We need to know the plant material on genetic level, and over all on the phytosanitary level. On the phytosanitary level, various viruses have been described on edible and medicinal yams on their production area. Different works, depending virus and virus group, are reported (Table 7). Indexing techniques allow to highlight a certain number of viruses on yam: Poty, potex, badna and cucumo-viruses, where yam mosaic virus (YMV) provokes the most important loss.

Table 7: Viruses of yam: group and type viruses, yam species affected and reference works

Virus Group	Virus	Yam species affected	Geographic spreading	Disease importance	Authors
Cucumovirus	CMV	<i>D. alata</i> , <i>D. cayenensis</i> - <i>D. Rotundata</i> complex, <i>D. Trifida</i>	Caribbean and West-Africa	-	Migliori, 1977; Fauquet and Thouvenel, 1987
'Carlavirus'	ChYNMV	<i>D. batatas</i>	Japan	-	Fukamoto and Tochiara, 1978;

cf.	(Chinese yam necrotic mosaic virus)				Shirako and Ehara, 1986
Badnavirus	DBV (<i>Dioscorea</i> bacilliform badnavirus)	<i>D. alata</i>	Barbade	+/-	Mantell and Haque, 1978
	DaBV (<i>D. alata</i> bacilliform virus)			-	Degras, 1986
	DbBV (<i>D. bulbifera</i> bacilliform virus)			-	Degras, 1986
Potexvirus	DLV (<i>Dioscorea</i> latent virus)	<i>D. floribunda</i> <i>D. composita</i>	Puerto Rico	-	Hearon et al., 1978; Phillips and Brunt, 1988; Waterworth et al., 1974
	PVX (Potato virus X)		<i>In vitro</i> collection	-	Urbino et al., 1998
Potyvirus	YMV (yam mosaic virus)	<i>D. alata</i> , <i>D. Cayenensis</i> - <i>D. Rotundata</i> complex,	All the intertropical area	+++	Thouvenel and Fauquet, 1979; Goudou-Urbino, 1995; Goudou-Urbino et al., 1996a,b
	YMMV ¹ (yam mild mosaic virus)	<i>D. alata</i> , <i>D. Cayenensis</i> - <i>D. Rotundata</i> complex,	West-Africa	+++	Mumford and Seal, 1997
	<i>D. trifida</i> virus ²	<i>D. trifida</i>	Guadeloupe	+++	Migliori, 1977
	DGBMV ² (<i>Dioscorea</i> green banding mosaic virus)		Togo	++	Porth and Nienhaus, 1983
	DaRMV ³ (<i>D. alata</i> ring mottle virus)	<i>D. alata</i>	Togo	++	Porth and Nienhaus, 1983
	DaV ⁴ (<i>D. alata</i> virus)	<i>D. alata</i> <i>D. rotundata</i>	Togo	+	Reckhaus and Nienhaus, 1981
	<i>D. dumetorum</i> potyvirus		South-Pacific	-	Mumford and Seal, 1997
	<i>D. esculenta</i> potyvirus		South-Pacific	-	Mumford and Seal, 1997
	DGBV (<i>Dioscorea</i> greenbanding potyvirus)	<i>D. composita</i> <i>D. floribunda</i>	Puerto Rico	-	Hearon et al., 1978; Phillips et al. 1986
	PVY (Potato virus Y)		<i>In vitro</i> collection		Urbino et al., 1998

¹ YMMV: is it a new potyvirus or a strain of the YMV?

² *D. trifida* virus and DGBMV have been shown as YMV strains (Porth et al., 1987)

³ DaRMV should be a 'yam strain' of the beet mosaic potyvirus transmissible on *N. benthamiana* (Porth et al., 1987)

⁴ DaV is serologically links to YMV but differ by is non-transmissibility (Porth et al., 1987)

During the establishment of the yam *in vitro* germplasm collection, in the biotechnology laboratory of ORSTOM (Maurie et al., 1993), afterwards IIRSDA - Adiopodoumé research station, near Abidjan, Ivory Coast - clones were systematically indexed by ELISA directed to YMV, when introduced *in vitro* (Maurie and Thouvenel, 1988; Maurie et al., 1988a,b; Charrier and Hamon, 1991).

Later on, one indexation was carried out by a virologist team on the duplicate of the yam *in vitro* germplasm collection enriched by introduction of new genotypes. 92

samples, belonging to several yam species, were used for the indexation : *D. alata*, *D. bulbifera*, *D. cayenensis-rotundata* complex, *D. dumetorum*, *D. esculenta*, *D. mangenotiana*, *D. praezensilis*, *D. shimperiana*, *D. togoensis*, *D. trifida*. These samples were originating from various geographic areas: Africa, Caribbean, South America and Asia. Four viruses were fetched by ELISA technique: PVX (potato virus X, potexvirus) PVY (potato virus Y, potyvirus), CMV (cucumber mosaic cucumovirus), and YMV (Urbino et al. 1998). Results are presented in Table 8.

Table 8: Indexation of the ORSTOM ² a. yam collection

ELISA results	PVX	PVY	CMV	YMV
% Positives	5,5	6,3	2,1	7,7
% Negatives	94,5	93,7	97,9	92,3

Source: Urbino et al. (1998)

This study allowed to show that detection of viruses serologically linked to PVX and to PVY, in different yam species, was possible, even with the same frequencies than with YMV. Further works have to be done for a precise characterization of these virus isolates, and check their respective importance in natural environment. Experiments using more sophisticated techniques for virus diagnostic (IC/rt/PCR) are developed (Bousalem, 1995). Yam on molecular characterization and molecular diversity on potyvirus of the yam mosaic virus (YMV) have been developed on the ILTAB/ORSTOM-TSRI laboratories (Aleman, 1996; Aleman et al 1996a,b) and from the LPRC

laboratory (Bousalem, 1995; Urbino et al., 1998).

Virus eradication techniques

The use of *in vitro* techniques allows to be free from fungus, bacteria, and other pests. Only viruses could be present on the plant and have to be eradicated. Different techniques exist and are already applied on yam. There are meristem culture, thermotherapy and/ or chemotherapy (36°C during 1 to 2 weeks on *in vivo* or *in vitro* plants, use of chemicals such as vidarabine, ribavirin and 2-thiouracil). They could be used alone or associated (Table 9).

Table 9 : Yam disease eradication techniques

	Eradication techniques	Species	Authors	Type of use*	Virus eradication
(A)	meristem culture	<i>D. cayenensis-rotundata</i> , <i>D. japonica</i> , <i>D. opposita</i> , <i>D. praezensilis</i> , <i>D. rotundata</i> , <i>D. trifida</i> , <i>Dioscorea</i> spp.	Cortes Monllor et al., 1982; Kobayashi, 1991; Maurie et al., 1988a,b, 1992, 1995a,b; Maurie and Thouvenel, 1988; Matsubaru and Ishira, 1988; Mikami, 1984; Saleil et al., 1990;	E	+ / -
(B)	Thermotherapy <i>in vivo</i> + meristems culture	<i>D. alata</i>	Mantell et al., 1980	E	+
(C)	Nodal microcutting or apices + Thermotherapy	<i>D. alata</i> , <i>D. Trifida</i>	Balagne, 1985; Salazar and Fernandez 1988	E, R (+ / -)	+
(D)	Nodal microcutting +	<i>D. alata</i>	Mantell, 1993	E, R (+ / -)	+

(E)	Chemotherapy Nodal microcutting + Thermotherapy &/or Chemotherapy	<i>D. praehensilis</i>	Malaurie, unpublished results	E	+ / -
(F)	Meristem culture + Thermotherapy &/or Chemotherapy	<i>D. cayenensis-rotundata</i> , <i>D. praehensilis</i>	Malaurie, unpublished results	E	+ / -

* E: experimental use; R: routine use

Success in meristem culture depends on the size and location of the explant excised, and on the growth regulator ratio. 'Meristem culture', on Table 9, concerns works using meristem-tips (0.2-0.5 mm long) as well as shoot-tips (0.6-2.5 mm long). Experiments on viability and *in vitro* morphological development of meristem-tips of two sizes, 'small' (0.3-0.5 mm) and 'large' (0.6-0.8 mm), have shown that it was better to use large meristem size to increase the shoot elongation percentage. The use of axillary or apical meristems did not induce difference and should allow an important yield in micropropagating such material from excised meristem-tips. Eleven months after meristem excision, production of plantlets was observed with a rate of 82% and 39% from the survivors, for a clone of *D.*

cayenensis-D. rotundata complex and *D. praehensilis* genotype, respectively (Malaurie et al., 1995a,b).

Meristem cultures have been done on 8 clones of 5 *Dioscorea* species belonging to the *in vitro* germplasm collection. Morphological development has been observed and data were recorded 60 days after meristem inoculation. In our case, the production of rooted leafy shoots, 60 days after meristem inoculation, occurred in five clones out of eight, with percentage shoot leaf production of 5 to 26 %, depending on the clone. Six months later, the excised meristems of all clones developed into rooted leafy shoots, where *D. bulbifera*, and *D. dumetorum* was not, to our knowledge, mentioned in the literature (Table 10).

Table 10: Genotypic effect on morphogenetic orientation 2 months after meristem excision of *Dioscorea* spp *

	Total meristems observed	Necrosis ¹⁾ %	Organogenesis ²⁾ %	Regeneration ³⁾ %
<i>D. alata</i>	67	27	55	18
<i>D. bulbifera</i>	252	46	37	18
<i>D. bulbifera</i>	146	52	48	0
<i>D. cayenensis-D. rotundata</i> complex	23	65	35	0
<i>D. cayenensis-D. rotundata</i> complex	81	24	51	26
<i>D. cayenensis-D. rotundata</i> complex	81	26	54	20
<i>D. dumetorum</i>	24	83	17	0
<i>D. praehensilis</i>	117	41	54	5

1) Necrosis. 2) Organogenesis: callusing, rooting, swelling were added together, 3) Regeneration: meristem development into rooted leafy shoots and axillary bud development or bud neoformation.

* (Malaurie, unpublished results)

Works about production of virus-free *in vitro* plants of yam through yam meristem culture alone are very rare. Saleil et al. (1990) on *D. trifida* obtained YMV-free plants, after ELISA indexation with a 27% rate through the total indexed plants. Nevertheless, other unpublished results on 2 genotypes YMV-infected of 2 species (*D. cayenensis-D. rotundata* complex, *D. praehensilis*) showed that meristem culture allowed the production of virus-free plants with 76% and 17% plants indexed, respectively (Malaurie, unpublished results).

Production of virus-free *in vitro* plants of yam has been attempted through thermotherapy, chemotherapy associated or not, from *in vivo* mother plants, nodal cuttings or apices (Balagne, 1985; Mantell, 1993; Mantell et al., 1980; Salazar and Fernandez, 1988). None of them described clearly the percentage of virus-free plants obtained through these techniques. Meanwhile, the production of plantlets free from virus is described by Mantell (1993) on *D. alata* cv. Kinabayo, after the action of antiviral agents (vidarabin, ribavirin) on nodal microcuttings infected by a potyvirus. The production of virus-free plants have been obtained 210

days after *in vitro* inoculation, after 3 subcultures of 60, 120 and 30 days on a liquid/solid biphasic culture system with 10^{-5} M of antiviral agent.

Other available techniques could be electrotherapy used on potato, with 60 to 100% success as compared to 25-40% with thermotherapy (Lozoya-Saldaña et al., 1996; Bernal et al., 1998), or apex micrografting, used on Lemon tree or vine, routinely.

If different works have already been done on yam sanitation, only a few of them conducted to an eradication of virus with more or less importance.

Safe international exchange

Exchange and distribution of plant material could be done by two ways: 1) with non aseptic plant material (tubers, aerial tubers, seeds, nodal cuttings from the vine), 2) with plant material in aseptic conditions (micro-nodal cuttings,

microtubers, aerial microtubers, apices, zygotic or somatic embryos, callus and cells suspension).

Exchange in non-aseptic conditions was used in the past, but required severe quarantine measures. Since 1989, with the FAO/IBPGR technical guidelines for the safe movement of yam germplasm, recommendation has been given to use *in vitro* conditions for exchange and distribution. For that, safe movement of yam germplasm could be done easily by three ways: 1) micro-nodal cuttings, 2) micro-tubers, 3) or encapsulated apices.

Safe movement of yam germplasm by micro-nodal cuttings is the most common way and has been frequently used (Malaurie et al., 1998a). In Table 11, the use of laboratories with *in vitro* and quarantine facilities allowed the indexation, *in vitro* introduction and micropropagation for a safe diffusion of various genotypes from different geographical origin.

Table 11: Enrichment of the genetic diversity of a country by transfer and introduction of yam genotypes from different geographic origins*

Species	Number of accession	Sending countries	Receiving countries
<i>D. alata</i>	6	Ivory Coast	New Caledonia
<i>D. alata</i>	5	West Indies	New Caledonia
<i>D. alata</i>	1	Brazil	New Caledonia
<i>D. alata</i>	5	West Indies	Ivory Coast
<i>D. alata</i>	3	New Caledonia	Ivory Coast
<i>D. alata</i>	3	Brazil	Ivory Coast
<i>D. bulbifera</i>	1	New Caledonia	Ivory Coast
<i>D. cayenensis-D. rotundata</i> complex	4	Ivory Coast	New Caledonia
<i>D. cayenensis-D. rotundata</i> complex	1	Brazil	Ivory Coast

* All plant material from the sending countries were, at first, tubers sent to laboratories with quarantine and *in vitro* culture facilities (1988-89: Orstom & Iirsda, Adiopodoumé, Côte d'Ivoire; 1992-95: Orstom, LRGAPT, Montpellier) for their *in vitro* introduction and micropropagation, preliminary to all safe international exchange.

Tuber potentiality shown by a great number of *in vitro* yams (aerial and basal micro-tubers) could be also used for a safe transfer of yam germplasm. They could increase the percentage of success during their acclimatation in field (John et al., 1993; Malaurie et al., 1993; Mantell, 1993; Ng, 1988; Ng and Mantell, 1997). These tubers developed *in vitro* are dormant at maturity and they still keep their dormancy from 2 to 5 months, as tubers developed *in vivo*. Recently a new method, experimented over three yam species (*D. alata*, *D. opposita*, *D. rotundata*), has been proposed by Hasan and Takagi (1995). They use encapsulation technique, with the embeddment of nodal cuttings in alginate beads, for a concept of a material transfer. This process allow to maintain in the dark for at least 2 weeks. These 2 weeks in the dark allow to envisage

a safe and easy international exchange of genetic resources.

Concluding Remarks

This paper tries to describe different studies done and to be done on yam *in vitro* germplasm conservation and its safe international exchange. Yam *in vitro* culture contributes to the safeguard of the biodiversity of the genus *Dioscorea*. An application of the results obtained on cryopreservation to more species should allow a transfer of technology. The use of new techniques, in a one hand, for pathogen eradication (electrotherapy, micrografting), in addition to the existent ones, and in the other hand, for the obtention of plants resistant to some viruses (transformation), should guarantee to yam a state of virus-free plant and allow

international exchanges, and in long term, distribution to the farmer of cultivar free from virus.

To conclude we can say that we are already able to manage routinely yam *in vitro* genebanks in slow growth culture, to index for more viruses, and to produce some virus-free *in vitro* plantlets.

For an efficient distribution - transfer - utilisation of yam germplasm, we should develop: 1) virus-free germplasm, 2) restricted size collection, with large diversity, so-called core-collections. For that, *in vitro* conservation under slow growth condition and cryopreservation, have to be applied routinely to more genotypes; virus-indexing has to be done with more precise techniques (rt/PCR); therapy has to be done with several combined techniques to become genotype independant.

But, we should not forget, as previously said by Hanson (1986), that, for a better security of germplasm conservation, different methods of conservation have to be combined (*in situ* - Field Genebanks - , *ex situ* - Seed Genebanks, *in vitro* Genebanks).

References

Aleman, M.E. (1996). Caractérisation moléculaire, diversité génétique et contrôle du virus de la mosaïque de l'igname (YMV). These, Doctorat en sciences agronomiques. ENSAM, Montpellier, France, 138p.

Aleman, M.E., Goudou-Urbino, C., Dubern, J. and Fauquet, C. (1996). Analysis of sequence variations in the P1, HC, P3, N1b and CP regions of Yam Mosaic Potyvirus isolates: implications for potyvirus intra-species molecular diversity. *Journal of General Virology* 78: 1253-1264.

Aleman, M.E., Marcos, J.F., Brugidou, C., Fux, C.I., Goudou-Urbino, C. and Dubern, J. (1996b). Genomic Organization, Molecular Diversity and Control of Yam (*Dioscorea*) Mosaic Virus. *ILTAB Review* 1991-1996, p. 81-82.

Ayensu, E.S. and Coursey, D.G. (1972). Guinea yams. The botany, ethnobotany, use and possible future of yams in West Africa. *Economic Botany* 26:301-318.

Balagne, M. (1985). Le microbouturage *in vitro* de l'igname cousse couche *Dioscorea trifida* en vue de son application pour la guérison des variétés atteintes de viroses. Thèse de 3ème cycle, USTL, Montpellier, 144 p.

Bernal, F.J.M., Hernández, R.P., Noa, J.C., Pichardo, T.P., Igarza, Y.C., Sarria, Z. and Agromonte, D.P. (1998). Técnicas de saneamiento para la obtención de papa (*Solanum tuberosum* L.) var. Desirée libre del virus del enrollamiento de la hoja. REDBIO'98-3rd Latin-American Meeting on Plant Biotechnology, June 1-5, La Habana, Cuba, p. 137.

Bousalem, M. (1995). Mise au point d'outils immunologiques et moléculaires pour la détection et la caractérisation des potyvirus infectant l'igname- Caractérisation moléculaire préliminaire de l'isolat BFC 56 du Yam Mosaic Virus. Laboratoire de Phytovirologie des Régions Chaudes (LPRC), Montpellier, Cirad-Orstom, 23p. (document interne).

Burkill, I.H. (1960). The organography and the evolution of Dioscoreaceae, the family of the yams. *J. Linn. Soc. (Bot.)* 56:319-412.

Butenko, R.G., Popov, A.S., Volkova, L.A., Chernyak, N.D. and Nosov, A.M. (1984). Recovery of cell cultures and their biosynthetic capacity after storage of *Dioscorea deltoidea* and *Panax ginseng* cells in liquid nitrogen. *Plant Science Letters* 33: 285-292.

Charrier, A., Dereuddre, J., Engelmann, F. (1991). The implications of biotechnology in germplasm conservation and utilization. *Crop Genetic Resources of Africa*. Vol II. N.Q. Ng, P. Perrino, F. Attere, H. Zedan ed., Ibadan, Nigeria, IITA/IBPGR/UNEP/CNR, In: Proceedings of an International Conference on Crop Genetic Resources of Africa. pp. 279-286.

Charrier, A. and Hamon, S. (1991). Germplasm collection, conservation and utilization activities of the Office de la Recherche Scientifique et Technique d'Outre-Mer (ORSTOM). *Crop Genetic Resources of Africa*. Vol II. N.Q. Ng, P. Perrino, F. Attere, H. Zedan ed., Ibadan, Nigeria, IITA/IBPGR/UNEP/CNR. In: Proceedings of an International Conference on Crop Genetic Resources of Africa. pp. 41-52.

Chulafich, L., Grubishich, D., Vuiichich, R., Volkova, L.A. and Popov, A.S. (1994). Somatic embryo production *in vitro* in *Dioscorea caucasica* Lipsky and *Dioscorea balcanica* Kosanin and cryopreservation of their organogenic callus tissue. *Russian Journal of Plant Physiology* 41:821-826.

Cortes Monllor, A., Liu, L.J. and Arroyo, E. (1982). An improved medium for tissue culture of yam *Dioscorea* sp. *in vitro* in Puerto Rico. *Phytopathology* 72:171.

Coursey, D.G. (1967). *Yams*, London, Longmans, 230 p.

Degras, L. (1986). L'igname, Techniques Agricoles et Productions Tropicales, Ed. G.P. Maisonneuve et Larose & Agence de Coopération Culturelle et Technique, 409 p.

Dereuddre, J., Scottez, C., Arnaud, Y. and Duron, M. (1990). Résistance d'apex caulinaires de vitroplants de poirier (*Pyrus communis* L. cv Beurré Hardy), enrobés dans l'alginat, à une deshydratation puis à une congélation dans l'azote liquide: effet d'un durcissement préalable au froid. *C. R. Acad. Sc. Paris, Série III*, 310:317-323.

- Dereuddre, J., Blandin, S. and Hassen, N. (1991). Resistance of alginate-coated somatic embryos of carrot (*Daucus carota* L.) to desiccation and freezing in liquid nitrogen: 1- Effects of preculture. *Cryo-Letters* 12:1125-134.
- Engelmann, F. (1991). *In vitro* conservation of tropical plant germplasm - a review. *Euphytica* 57:227-243.
- Fauquet, C. and Thouvenel, J.C. (1987). Cucumber mosaic on sweet potato and yam. *In: Plant Viral Diseases in the Ivory Coast, Documentations Techniques n° 46, Editions de l'Orstom, Paris, p. 29.*
- FAO/IBPGR (1989). Technical guidelines for the safe movement of yam germplasm. A.A. Brunt, G.V.H. Jackson, E.A. Frison, ed., Rome, pp. 1-20.
- FAO (1996). The State of the World's Plant Genetic Resources for Food and Agriculture. Background Documentation prepared for the International Technical Conference on Plant Genetic Resources, Leipzig, Germany, 17-23 June, 1996. FAO : Rome, 336 p.
- Fukamoto, F. and Tochihara, H. (1978). Chinese yam necrotic virus. *Annals of the . Phytopathological Society of Japan* 44:1-5.
- Goudou-Urbino, C. (1995). La mosaïque de l'igname: aspects épidémiologiques au Burkina-Faso et variabilité du virus. Thèse de l'Université Montpellier II, 147 p.
- Goudou-Urbino, C., Givord, L., Quiot, J.B., Boeglin, M., Konate, G. and Dubern, J. (1996a). Differentiation of yam virus using symptomatology, western blot assay, and monoclonal antibodies. *Journal of Phytopathology* 144:235-240.
- Goudou-Urbino, C., Konate, G., Quiot, J.B. and Dubern, J. (1996b). Etiology and ecology of yam mosaic disease in Burkina Faso. *Tropical Sciences*, 36:34-40.
- Hanson, J. (1986). Methods of storing tropical root crop germplasm with special reference to yam. *FAO/IBPGR Plant Genetic Ressources Newsletter* 64:24-32.
- Hasan, S.M.Z. and Takagi, H. (1995). Alginate-coated nodal segments of yam (*Dioscorea* spp.) for germplasm exchange and distribution. *Plant Genetic Resources Newsletter* 103:32-35.
- Hearon, S.S., Corbett, M.K., Lawson, R.H., Gillespie, A.G. and Waterworth H.E. (1978). Two flexuous-rod viruses in *Dioscorea floribunda*: symptoms, identification and ultrastructure. *Phytopathology* 68:1137-1146.
- IBPGR (1986). 2. Root and Tuber Crops (2nd Ed.), *In Directory of Germplasm Collections*. J. Toll, T. Lawrence, D.H. Van Sloten Ed. International Board for Plant Genetic Resources, Rome, pp. 130-155.
- John, J.L., Courtney, W.H. and Decoteau, D.R. (1993). The influence of plant growth regulators and light on microtuber induction and formation in *Dioscorea alata* L. cultures *Plant Cell, Tissue and Organ Culture* 34:245-252.
- Knuth, R. (1924). *Dioscoreaceae*. A. Engler. *Das Pflanzenreich*, IV-43, 87. Heft, 1-387.
- Kobayashi, N. (1991). Production of virus-free plants and its mass propagation by tissue culture in Chinese yam, *Dioscorea opposita* Thunb. *Bulletin of the Saitama Horticultural Experiment Station*. N° 18, 81-99.
- Latta, R. (1971). Preservation of suspension culture of plant cells by freezing. *Canadian Journal of Botany* 40: 253-1254.
- Lozoya-Saldana, H, Abello, F.J. and Garcia G. de la R. (1996). Electrotherapy and shoot tip culture eliminate potato virus X in potatoes. *American Potato Journal* 73:149-154.
- Malaurie, B., Tardieu, F. and Thouvenel, J.C. (1988a). Rapid production of disease-free germplasm of *Dioscorea* spp. (Monocotyledonous). *In: Proceedings of 8th International Biotechnology Symposium, Paris, 17-22 July 1988, p. 248.*
- Malaurie, B., Tardieu, F. and Thouvenel, J.C. (1988b). Improving Yam (*Dioscorea* spp.), using Biotechnology. 1) Rapid production of disease free germplasm. *In: Proceedings of the International Society for Root Crops Symposium. Bangkok, 30/10-5/11/1988, p. 51.*
- Malaurie, B. and Thouvenel, J.C. (1988). Production de plants d'igname, dépourvus de viroses, par culture de méristèmes. 2° Salon International de la Coopération et de l'Aide au Développement (SICAD). 7-11/12/1988, Montpellier, Poster Session, 3 p.
- Malaurie, B., Pungu, O., Dubern, J. and Thouvenel, J.C. (1992). Determination of the best conditions for the regeneration of microplants and the elimination of YMV from excised meristems of yam nodal cuttings (*Dioscorea* spp.). *Association of Applied Biologists. Plant Virology in the Tropics. University of York. 9-10 April 1992.*
- Malaurie, B., Pungu, O., Dumont, R. and Trouslot, M.F. (1993). The creation of an *in vitro* germplasm collection of yam (*Dioscorea* spp.) for genetic resources preservation. *Euphytica* 65: 113-122. *Corrigendum* 66: 243.
- Malaurie, B., Pungu, O. and Trouslot, M.F. (1995a). Effect of growth regulators concentrations on morphological development of meristem-tips in *Dioscorea cayenensis-D.*

rotundata complex and *D. praehensilis*. Plant Cell, Tissue and Organ Culture 41:229-235.

Malaurie, B., Thouvenel, J.-C. and Pungu, O. (1995)b. Influence of meristem-tip size and location on morphological development in *Dioscorea cayenensis*-*D. rotundata* complex 'Grosse Caille' and one genotype of *D. praehensilis*. Plant Cell, Tissue and Organ Culture 42: 215-218.

Malaurie, B. and Trouslot, M.F. (1995c). Les Igname. In Biotechnologies végétales BV9D, CNED-AUPELF-UREF ed., chapitre 10:49-77.

Malaurie, B. and Trouslot, M.F. (1996). Cryopreservation of *in vitro* yam (*Dioscorea* sp.) apices by the encapsulation-dehydration technique. Eucarpia, Meeting on Tropical Plants, Marsh 11-15 1996, France, Abstract, p 252.

Malaurie, B., Trouslot, M.F. and Berthaud, J. (1998a). Conservation et échange de germoplasme chez les ignames (*Dioscorea* spp.). 'L'igname : Plante Séculaire et Culture d'Avenir'. eds. J. Berthaud, N. Bricas, J.-L. Marchand, Montpellier 3-6 Juin 1997, In: Acte du séminaire international Cirad-Inra-Orstom-Coraf, France. 135-161.

Malaurie, B., Trouslot, M.F., Engelmann, F. and Chabrilange, N. (1998b). Effect of pretreatment conditions on the cryopreservation of *in vitro*-cultured yam (*Dioscorea alata* 'Brazo Fuerte' and *D. bulbifera* 'Nouméa Imboro') shoot apices by encapsulation-dehydration. Cryo-Letters 19:15-26.

Malaurie, B., Trouslot, M.F., Berthaud, J., Chabrilange, N., Récalt, C. and Dussert, S. (1998c). The use of slow growth condition culture and cryopreservation in liquid nitrogen for medium and long term conservation and utilisation of *in vitro* yam (*Dioscorea* spp.) germplasm. In: Proceedings of the Workshop on Conservation & Utilisation of Cassava, Sweetpotato and Yam Germplasm in Sub-Saharan Africa. Nairobi, November 11 to 13, 1997, (in press).

Mandal, B.B., Chandel, K.P.S. and Dwivedi, S. (1996). Cryopreservation of yam (*Dioscorea* spp.) shoot apices by encapsulation-dehydration. Cryo-Letters 17:165-174.

Mantell, S.H. and Haque, S.Q. (1978). Incidence of internal brown spot disease in White Lisbon (*D. alata*) during storage. Experimental Agriculture 14:167-172.

Mantell, S. H., Haque, S. Q. and Whitehall, A. P. (1980). Apical meristem tip culture for eradication of flexuous rod viruses in yams (*Dioscorea alata*). Tropical Pest Management 26 : 170-179.

Mantell, S.H. (1993). Integrated use of micropropagation and conventional propagation techniques for production of certified seed tubers of tropical yams (*Dioscorea* spp.).

Adapted propagation techniques for commercial crops of the tropics, IFS, Vietnam, 02/1993. In: Proceedings of the Southeast Asian Regional Workshop on Propagation Techniques for Commercial Crops of the Tropics 66-93.

Martin, F.W. and Degras, L. (1978). Tropical yams and their potential. Part 6. Minor cultivated *Dioscorea* species. USDA agriculture handbook n° 538. 23 p.

Martin, F.W. and Rhodes, A.M. (1978). The relationship of *Dioscorea cayenensis* and *D. rotundata*. Tropical Agriculture, Trin., 55:193-206.

Matsubara, S. and Ishihara, M. (1988). Production and vegetative propagation of virus-free plants of *Dioscorea* species. Scientific Reports of the Faculty of Agriculture, Okayama University, N° 72, 19-26.

Miège, J. (1982). Etude chimiotaxonomique de dix cultivars de Côte d'Ivoire relevant du complexe *Dioscorea cayenensis*-*D. rotundata*. Yams-Igname, J. Miège, S.N. Lyonga, Ed., Claredon Press Oxford, pp. 197-231.

Migliori, A. (1977). Maladies à virus de l'igname *Dioscorea* spp. Nouvelles agronomiques Antilles Guyane 3:428-435.

Mikami, T. (1984). Virus-free plant propagation through meristem-tip culture of Japanese yam and garlic. Research Journal of Food and Agriculture 7:17-20.

Mumford, R.A. and Seal, S.E. (1997). Detection and characterization of yam potyviruses. In: Yam viruses, Annual Collaborators Meeting, IITA - JIC - NRI, Gatsby-funded Biotechnology Projects, 7-9 February 1997, IITA, Ibadan, Nigeria, Extended Abstracts, pp. 14-15.

Ng, S.Y.C. (1988). *In vitro* tuberization in white yam (*Dioscorea rotundata* Poir.). Plant Cell, Tissue and Organ Culture 14:121-128.

Ng, S.Y.C. and Mantell, S.H. (1997). Technologies for Germplasm Conservation and Distribution of pathogen-free *Dioscorea* yams to National Root Crop Research Programmes. ODA Project R4886 (H) Final Report. Wye College University of London, UK, 12p.

Phillips, S. and Brunt, A.A. (1988). *Dioscorea* latent virus. AAB Descriptions of Plant Viruses, N°. 335. Association of Applied Biologists, Wellesbourne.

Phillips, S., Pigott, J. d'A. and Brunt, A.A. (1986). Further evidence that *Dioscorea* latent virus is a potyvirus. Annals of Applied Biology 109:137-145.

Polge, C.A., Smith, A.V. and Parkes, A.S. (1949). Revival of spermatozoa after vitrification and dehydration at low temperature. Nature 164: 666.

- Popov, A.S. and Volkova, L.A. (1994). Cryopreservation and some characteristics of *Dioscorea deltoidea* cell cultures in the vitamin-free medium. Russian Journal of Plant Physiology 41:815-820.
- Popov, A.S. and Fedorovskii, D.N. (1992). Injuries to the plasmalemma of *Dioscorea* cells cultured *in vitro* incurred in the process of their cryopreservation. Soviet Plant Physiology 39:211-216.
- Porth, A., Lesemann, D.-E. and Vetten, H.J. (1987). Characterization of potyvirus isolates from West Africa yams (*Dioscorea* spp.). Journal of Phytopathology 120:166-183.
- Porth, A. and Nienhaus, F. (1983). *Dioscorea alata* ring mottle virus, a new potyvirus in yam in Togo. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 90:352-362.
- Reckhaus, P. and Nienhaus, F. (1981). Etiology of a virus disease of white yam *D. rotundata* in Togo. Journal of plant disease and protection 88:492-509.
- Sakai, A. (1984). Cryopreservation of apical meristems. In Horticultural Reviews, J. JANICK Ed., Purdue University, pp. 357-372.
- Salazar, S.S and Fernandez, R.Z.Z. (1988). Thermotherapy, shoot tip culture, axillary bud proliferation and plant regeneration in yam (*Dioscorea trifida* L.). In VIth Symposium of the International Society for Tropical Root Crops, Gosier (Guadeloupe), 1-6 July 1985, INRA Ed., Paris, p. 439-445.
- Saleil, V., Degras, L. and Jonard, R. (1990). Obtention de plantes indemnes du virus de la mosaïque de l'igname (YMV) par culture *in vitro* des apex chez l'igname américaine *Dioscorea trifida* L. Agronomie 10:605-615.
- Shirako, Y. and Ehara, Y. (1986). Rapid diagnosis of chinese yam necrotic mosaic virus infection by electro-blot immunoassay. Annals of the Phytopathological Society of Japan 52: 453-459.
- Tanaka, S. (1977). Chinese yam *Dioscorea opposita* Thunb. in Japan. Tropical Root and Tuber Crops Newsletter n°10, pp. 4-5.
- Tannoury, M., Ralambossa, J., Kaminske, M. and Dereuddre, J. (1991). Cryoconservation par vitrification d'apex enrobés d'oeillets (*Dianthus caryophyllus* L.) cultivés *in vitro*. C. R. Acad. Sc. Paris, Série III, 313:633-638.
- Thouvenel, J.-C. and Fauquet, C. (1979). Yam mosaic, a new potyvirus infecting *Dioscorea cayenensis* in the Ivory Coast. Annals of Applied Biology 93:279-283.
- Uragami, A. (1993). Cryopreservation of cultured cells and organs of vegetables. In Reprint from JICA GRP REF N°6, 111-135.
- Urbino, C., Bousalem, M., Pinel, A., Fargette, D. and Dubern, J. (1998). Les Virus de l'igname: aspects épidémiologiques et variabilité. 'L'igname: Plante Séculaire et Culture d'Avenir'. eds. J. Berthaud, N. Bricas, J.-L. Marchand, Montpellier 3-6 Juin 1997, In: Acte du séminaire international Cirad-Inra-Orstom-Coraf, France. 205-211.
- Watterworth, H.E., Lawson, R.H. and Kahn, R.P. (1974). Purification, electron microscopy, and serology of *Dioscorea* latent virus. Journal of Agriculture of the University of Puerto Rico 58:351-357.
- Withers, L.A. (1991). Crop strategies for roots and tubers: potato- A model for refinement, Yam- a problem for development. In Workshop on Conservation of Plant Genetic Resources. Becker B. Ed., Bonn. ATSAF/IBPGR, pp. 11-14.