

Biochar and *Trichoderma aureoviride* applied to the sandy soil: effect on soil quality and watermelon growth

Erika V. MEDEIROS^{1*}, Marcelle C.H.S. MORAES¹, Diogo P. COSTA¹,
Gustavo P. DUDA¹, Jenifer S.A. SILVA¹, Julyana B. OLIVEIRA¹,
José R.S. LIMA¹, Romulo S.C. MENEZES², Claude HAMMECKER³

¹Universidade Federal do Agreste de Pernambuco, Garanhuns, Pernambuco, Brazil; evmbio@gmail.com (*corresponding author);
marcele-moraes@hotmail.com; diogocosta1@yahoo.com.br; gpduda@gmail.com; jenifersthephanie12@hotmail.com;
jully_bragaa@live.com; romualdo_solo@yahoo.com.br

²Universidade Federal de Pernambuco, 1235 - Cidade Universitária, 50i70-901, Recife, Pernambuco, Brazil; rmenezes@ufpe.br

³Institut de Recherche pour le Développement (IRD), 2, Place Pierre Viala, Montpellier, France; claud.hammecker@gmail.com

Abstract

Trichoderma spp. are an alternative to increase plant growth and as biological control agents of diseases. Biochar added to soil and *Trichoderma* can result in the enhancement of crop development and can aid in preventing fertilizer run-off, improving plant health, retaining soil moisture, and helping plants through drought periods. However, a knowledge gap remains regarding the combined effect of biochar and *Trichoderma* on soil quality and crop growth. The objective of the present study was to evaluate this combined effect and show a new approach of biochar as a route of *T. aureoviride* (T) inoculation. We evaluated three sources of biochar, bean husks, coffee grounds, and coffee husks, with or without T and an additional control that was devoid of biochar and T. The association of all biochar sources with T showed a significant increase in watermelon growth and soil quality. Biochar coffee grounds with T increased urease, β -glucosidase, and total organic carbon, showing the potential of this combination in these processes. Watermelon plants cultivated in soil with coffee ground biochar and T showed plant length and shoot dry biomass increases of 129% and 192%, respectively. Thus, this study found that biochar use plays an important role in the effectiveness of *T. aureoviride* applications and is a new instrument for sustainable agriculture.

Keywords: *Citrullus lanatus*; cycling of nutrients; microbial biomass; soil fertility; soil metabolism; vegetative growth

Introduction

Biochar is a stable product obtained from the pyrolysis of organic materials, such as rice straw, wood, manure, and leaves under a low oxygen (O₂) concentration. Originally, it was found in Amazonian regions in soil bands known as "Black Indian lands" (Saxena *et al.*, 2013). During the last several years, this material has received prominence because of its potential for carbon sequestration, heavy metals adsorption, and increased

availability of phosphorus (P) in the soil because of the increase in pH and the adsorption of this nutrient on its surface (Elzobair *et al.*, 2016; Yang *et al.*, 2017). Biochar also increases cation exchange capacity, enzymatic activities, and water retention in soil, mitigating saline stress and improving the development of plants and microorganisms (Patel *et al.*, 2017).

Among the microorganisms, we highlight the beneficial fungi of the genus *Trichoderma*, which act as agents for the biological control of phytopathogens and promote plant growth (Silva *et al.*, 2016). Among the mechanisms conferred by these fungi is the increase in the area of root absorption because of the *Trichoderma* root association, resulting in a higher nutritional absorption, followed by an increase in the pathogen resistance factor (Oskiera *et al.*, 2017). This fungus assists in conserving soils, increasing seed germination rates, mitigating abiotic interference effects, and characterizing a systemic defence accompanied by increases in soil productivity (Mendoza-Mendoza *et al.*, 2018). The fungus *T. aureoviride* showed physiological and functional versatility mentioned in previous studies, such as the capacity to produce the enzyme chitinase that acts on the cell wall of several fungi and in the direct biological control of phytopathogens. (Silva *et al.*, 2016).

Considering the ease of cultural adaptation and economic relevance, the watermelon (*Citrullus lanatus* Thumb. Mansf.) culture constitutes a relevant model to evaluate the effects of biochar inoculated with *Trichoderma* because it has a short cycle, facilitating experimental trials, and its pulp, bark, and seeds are rich in minerals, vitamins, and proteins (Poduria *et al.*, 2013).

There are several studies relating the beneficial effects of biochar applied to soil (Patel *et al.*, 2017; Lima *et al.*, 2018) and the use of *Trichoderma* as instrument to promote plant growth or biological control (Silva *et al.*, 2017). Nonetheless, there is still a gap in knowledge regarding the effect of the combined application of both on soil quality and plant growth. There is no report regarding the use of this combined application of biochar and *Trichoderma* in watermelon culture; thus, this is a pioneering study in this context. This study presents the hypothesis that biochar is a potential vehicle for inoculation of *Trichoderma* spp., improving several soil attributes that in turn increase crop development. Therefore, the objective of the present study was to evaluate the combined effect of biochar and *T. aureoviride* on the improvement of soil quality and the initial development of watermelon plants.

Materials and Methods

Trichoderma aureoviride strain and biochar application

Biochar was obtained from industrial bean and coffee residues, namely, bean husks (BH), coffee grounds (CG), and coffee husks (CH). They were charred for 10 to 12 h under oxygen-limited conditions during a slow pyrolysis process where the temperature reached 530 °C in a metallic kiln and were then sieved using a 2 mm sieve. The quality of the biochar has been assessed by Lima *et al.* (2018).

T. aureoviride URM 5158 isolate was obtained from the Micoteca URM collection of the Mycology Department of the Biological Sciences Center of the Federal University of Pernambuco (<https://www.ufpe.br/micoteca/>). The fungus conidia reactivated through three successive subcultures in an Erlenmeyer flask containing 50 ml of potato dextrose agar liquid medium, grown at 26 °C ± 2 °C for eight days. Each treatment with *T. aureoviride* was sprayed with 100 ml (1 × 10⁶ conidia per ml) applied to each biochar source. The biochar and *T. aureoviride* were mixed with the soil before planting.

Experimental setup

A pot experiment was conducted in a greenhouse in Pernambuco, Brazil. The soil used for the experiment was a topsoil (the 0-20 cm layer) collected in a patch of native forest in São João (08° 48' 34.2" S,

36° 24' 29.3" W) at an elevation of 705 m above the mean sea level. The soil is a typical entisol with nearly 90 % sand and less than 5 % clay with extremely low values of cation exchange capacity.

The chemical attributes of the soil and biochar before the experiment are shown in Lima *et al.* (2018). This soil was previously treated with a formulated fertilizer NPK (6:24:12) to prevent variations related to plant nutritional deficits. We added biochar corresponding to the application of 32 t ha⁻¹ for each biochar, considering its incorporation in the first 20 cm of the soil.

The experimental design used was completely randomized in a double factorial scheme with an additional treatment (3 × 2 + 1), with four repetitions. The first factor represented three sources of biochar, BH, CG, and CH, and the second considered inoculation (T+) and absence (T-) of *T. aureoviride* on these substrates. The additional treatment represented the absolute control without biochar or *T. aureoviride*.

We used a commercial variety of watermelon (variety 'Vereda style') used by the farmers in the region where the experiment was conducted. The seeds were previously treated with fungicide, and three seeds were sown per pot. Five days after emergence, thinning was performed to leave one plant per pot. The plants were irrigated daily with distilled water to maintain the soil at field capacity.

At the end of the experiment, 45 days after planting, the plants were removed to measure the diameter of the base (DIA), main branch length (LEN), shoot (SFB) and root fresh biomass (RFB), followed by their respective dry biomass after drying in a forced ventilation oven at 65 °C until reaching the consistency of the dehydrated biomasses (shoot dry biomass, SDB, and root dry biomass, RDB). The experiments were repeated two times and the means of both were showed.

After removal of the plants from the pots, soils were collected for the determination of pH values in water (1:2.5) and the content of P, Na⁺, K⁺, Ca²⁺, Mg²⁺, Al³⁺, and H+Al³⁺, according to the methodology of Silva (2009). The inorganic labile P, Na⁺, and K⁺ were extracted using Mehlich's solution 1 (H₂SO₄ 0.125 mol L⁻¹ + HCl 0.5 mol L⁻¹). The P was quantified using spectrophotometric colorimetry and the Na⁺ and K⁺ were determined using flame photometry. The Ca²⁺, Mg²⁺, and Al³⁺ were extracted with a KCl 1.0 mol L⁻¹ solution; the first two were determined using atomic absorption spectrometry (Raij *et al.*, 2001). Potential acidity (H+Al³⁺) was determined using calcium acetate (Ca (CH₃COO)₂ H₂O) at pH 7.0. In addition, Al³⁺ was determined via titration with NaOH 0.025 mol L⁻¹ using bromothymol blue as an indicator. In all extractions, 5.0 g of soil was used for 50 mL of extractive solution (1:10).

Total organic carbon content (TOC) was obtained using the C oxidation method in a wet form with potassium dichromate followed by determination of remaining Cr₂O₇²⁻ via titration with ammonium iron (II) sulfate (Yeomans and Bremner, 1988). The microbial biomass carbon (MBC) was determined using the methodology of irradiation (Mendonça, 2005) with the extraction carried out by the addition of 80 mL of K₂SO₄ 0.5 M to each 20 g of soil (Tate *et al.*, 1988). The C content was determined using colorimetry (Bartlett and Ross, 1988).

The evolution of CO₂ released by microbial respiration was also measured for the calculation of soil basal respiration and the soil metabolic quotient (*q*CO₂) using the alkaline adsorption method. For this purpose, at the end of the experiment, the soil was maintained at 60% of field capacity based on gravimetry (Anderson and Domsch, 1985). Then, 30 g of each sample were hermetically sealed in containers to capture CO₂ with 10 mL of NaOH 0.5 M solution. After 72 h, 2 mL of BaCl 10% (m/v) solution were added for total precipitation of CO₂, followed by two drops of phenolphthalein diluted in 100 ml of ethyl alcohol 95% (v/v the determinations being carried out by titration with HCl 0.25 M).

Enzymatic soil activities were determined by estimating the gene expression of the main processes related to acidic (P. Aci) and alkaline phosphatases (P. Alk) (E.C. 3.1.3), β-glucosidase (Beta) (E.C. 3.2.1.21), and urease (Ure) (E.C. 3.5.1.5) according to the colorimetric analysis of the products released by the samples subjected to incubation in an adequate substrate (Sigma-Aldrich). The activities of P. Aci and P. Alk were estimated according to the methodology of Eivazi and Tabatabai (1977). Beta was estimated using the methodology of Eivazi and Tabatabai (1988), and Ure was estimated using the method by Kandeler and Gerber (1988). Product absorbance was measured using a spectrophotometer (Libra S22, Biochrom, Cambridge, UK).

Statistical analyses

Statistical and multivariate analyses were conducted using the R language platform (v.3.4.3, 2017) based on the data of the biometric attributes of the plants, chemicals, and enzymatic activities of the soils. Multivariate exploratory analyses including nonmetric multidimensional scaling (nMDS), analysis of similarity (ANOSIM), heatmaps, and Mantel tests were completed based on the tools of the vegan and heatmaply libraries. After adjusting the models and removing the outliers, the means, standard deviations, and variation coefficients were calculated using the *doBy* library. Homogeneity tests, analysis of variances (ANOVA), and comparisons between averages were completed with the balanced data, according to the tools contained in the stats, AxpDes, multcomp, and agricolae libraries. Residues from all datasets were considered normal, with the exception of SDB and K, which were adequately transformed for ANOVA.

Results

The results showed the influence of the biochar source types and *T. aureoviride* on watermelon plant attributes (Table 1), soil chemical attributes (Table 2) and metabolic pattern of the soil microbial community (Table 3).

Table 1. Watermelon plants attributes (cv. Vereda style) treated with biochar from coffee grounds (CG), coffee husks (CH), and bean husks (BH), with (+T) and without (-T) inoculation of *Trichoderma aureoviride*

	CG		CH		BH	
	1. Length (cm)					
+T	<u>95.6</u>	aA	79.3	b	<u>91.7</u>	a
-T	69.3	bB	75	b	<u>91.3</u>	a
	C = 73.56					
	2. Diameter (cm)					
+T	7.2	aA	6.4	bB	7.4	aA
-T	6.7	abB	7.1	aA	6.6	bB
	C = 6.78					
	3. SFM (g)					
+T	<u>76.7</u>	aA	<u>60</u>	bA	<u>73.3</u>	a
-T	<u>60.6</u>	bB	47.8	cB	<u>75</u>	a
	C = 44.78					
	4. RFM (g)					
+T	<u>4.2</u>	bA	2.5	cA	<u>6.1</u>	aA
-T	2.3	bB	<u>0.7</u>	cB	<u>4</u>	aB
	C = 2.36					
	5. SDM (g)					
+T	<u>28.9</u>	aA	15.6	bA	<u>28.3</u>	a
-T	14.4	bB	<u>9.4</u>	bB	<u>25.8</u>	a
	C = 15.00					
	6. RDM (g)					
+T	<u>0.21</u>	b	<u>0.14</u>	cA	<u>0.44</u>	aA
-T	<u>0.17</u>	b	0.06	cB	<u>0.32</u>	aB
	C = 0.047					

LEN= plant length; DIA= diameter; SFM= shoot fresh mass; RFM= root fresh mass; SDM= shoot dry mass; RDM= root dry mass. Means followed by the same lowercase in the line (between different sources of biochar) and upper case in the column (between the effects of *Trichoderma aureoviride*), do not differ statistically by Tukey's test at 5%. The absence of letters indicates that the means are statistically equal to the 5% level using the F test. The underlined averages differ significantly from the respective control treatments (C) by Dunnett test at 5%

The inoculation of *T. aureoviride* with biochar also demonstrated a significant increase that was, higher than that in the control treatment, with the exception of the diameter. However, the results demonstrated a significant positive effect of *T. aureoviride* inoculation on the vegetative development of watermelon plants grown in a regolithic neosol treated using different biochar sources.

Table 2. Chemical attributes of soil cultivated with watermelon (cv. Vereda style) treated with biochar from coffee grounds (CG), coffee husks (CH), and bean husks (BH), with (+T) and without (-T) inoculation of *Trichoderma aureoviride*

	CG		CH		BH	
	<i>1. pH</i>					
+T	<u>5.57</u>	b	<u>7</u>	aB	<u>6.74</u>	a
-T	<u>5.48</u>	c	<u>7.89</u>	aA	<u>6.65</u>	b
	C = 4.54					
	<i>2. Na²⁺ (cmolc kg⁻¹)</i>					
+T	0.82	ab	0.93	a	0.82	b
-T	0.89	ab	0.89	a	0.85	b
	C = 0.821					
	<i>3. K⁺ (cmolc kg⁻¹)</i>					
+T	<u>2.13</u>	c	<u>6.7</u>	a	<u>3.97</u>	bB
-T	2.13	b	<u>5.93</u>	a	<u>5.19</u>	aA
	C = 0.596					
	<i>4. P (mg kg⁻¹)</i>					
+T	<u>2.03</u>	c	<u>3.02</u>	b	<u>4.14</u>	aA
-T	<u>2.3</u>	b	<u>3.03</u>	a	<u>3.4</u>	aB
	C = 0.875					
	<i>5. H⁺+Al³⁺ (cmolc dm⁻³)</i>					
+T	<u>1.13</u>	aB	<u>0.52</u>	b	<u>0.63</u>	b
-T	<u>1.4</u>	aA	<u>0.47</u>	b	<u>0.52</u>	b
	C = 1.84					
	<i>6. Ca²⁺ (cmolc dm⁻³)</i>					
+T	0.39	aB	<u>0.45</u>	a	<u>0.44</u>	aA
-T	<u>0.46</u>	aA	<u>0.42</u>	ab	0.39	bB
	C = 0.346					
	<i>7. Mg²⁺ (cmolc dm⁻³)</i>					
+T	0.36	b	<u>0.38</u>	b	<u>0.6</u>	a
-T	<u>0.4</u>	b	<u>0.4</u>	b	<u>0.57</u>	a
	C = 0.250					

Means followed by the same lowercase in the line (between different sources of biochar) and upper case in the column (between the effects of *Trichoderma aureoviride*), do not differ statistically by Tukey's test at 5%. The absence of letters indicates that the means are statistically equal to the 5% level using the F test. The underlined averages differ significantly from the respective control treatments (C) by Dunnett test at 5%.

Table 3. Enzyme activities, total organic carbon, microbial biomass carbon and methabolic quotient of soil cultivated with watermelon plants treated with biochar from coffee grounds (CG), coffee husks (CH), and bean husks (BH), with (+T) and without (-T) inoculation of *Trichoderma aureoviride*

	CG		CH		BH	
	1. Ure ($\mu\text{g NH}_4\text{-N g}^{-1}\text{dwt } 2\text{h}^{-1}$)					
+T	30.6	b	<u>44.4</u>	a	<u>40.4</u>	a
-T	<u>31.3</u>	b	<u>43.9</u>	a	<u>44.3</u>	a
	C = 25.32					
	2. Beta ($\mu\text{g p-Nitrof. g}^{-1}\text{soil h}^{-1}$)					
+T	<u>58.6</u>	aA	43.4	bA	48.1	bA
-T	<u>53</u>	aB	35.9	bB	40.3	bB
	C = 39.75					
	3. P.Alk ($\mu\text{g p-Nitrof. g}^{-1}\text{soil h}^{-1}$)					
+T	3.1	aA	1.7	b	2.4	ab
-T	1.1	bB	2.5	a	2.6	a
	C = 2.26					
	4. P.Aci ($\mu\text{g p-Nitrof. g}^{-1}\text{soil h}^{-1}$)					
+T	1.9	a	1.5	b	2.3	ab
-T	3.7	a	1.1	b	2.4	ab
	C = 1.40					
	5. TOC ($\text{g kg}^{-1}\text{ soil}$)					
+T	<u>12.8</u>	a	8.7	b	<u>10.5</u>	b
-T	<u>12.9</u>	a	<u>9.4</u>	b	8.7	b
	C = 6.88					
	6. MBC ($\text{g kg}^{-1}\text{ soil}$)					
+T	0.1	b	0.3	ab	0.3	aB
-T	0.2	b	0.2	b	<u>0.5</u>	aA
	C = 0.16					
	7. $q\text{CO}_2$ ($\text{mgC-CO}_2\text{g}^{-1}\text{BMS-C h}^{-1}$)					
+T	2.4	abA	2.5	aB	2	bA
-T	1.5	abB	2.9	aA	0.7	bB
	C = 2.04					

Ure= soil Urease; Beta= β -glycosidase; P.alk= alkaline phosphatase; P.aci=acid phosphatase activities; TOC= total organic carbon; MBC= microbial biomass carbon; $q\text{CO}_2$ = soil metabolic quotient. Means followed by the same lowercase in the line (between different sources of biochar) and upper case in the column (between the effects of *Trichoderma aureoviride*), do not differ statistically by Tukey's test at 5%. The absence of letters indicates that the means are statistically equal to the 5% level using the F test. The underlined averages differ significantly from the respective control treatments (C) by Dunnett test at 5%

The plant biometric variables significantly responded to both biochar type and T in contrast with the control treatment. Most of the plant parameters were strongly influenced (increased) by CG or BH and T application into the soil (Table 1). For example, the application of CG biochar with *Trichoderma aureoviride* to the soil increased the plant length, diameter, shoot fresh and dry matter and root dry matter (1.3; 1.06; 1.7;

1.9; 4.4 times, respectively). Watermelon plants that received BH biochar with T also showed increases, mainly on DIA, RFM, and RDM (9.3 times compared to the control).

The addition of biochar, in general, increased the pH and the P, K⁺, Ca²⁺, and Mg²⁺ and decreased the H+Al³⁺ compared to the control. BH was also generally responsible for the higher levels of P, particularly when applied with *T. aureoviride* (increase of 4.7 times compared to the control), differing from all other treatments, with the exception of CG (T-) (Table 2). The treatments with CG and BH also contributed to a reduction in potential acidity (H+Al³⁺), which was higher than that of CG, in general and was closely correlated with the changes in pH. The Ca²⁺ responses to the treatments were quite varied and were the same as the treatments with CG (T-), CG (T+ and T-), and BH (T+), which were all superior to CG (T+) and BH (T-).

The chemical variables pH, Na⁺, K⁺, P, and H+Al³⁺ significantly responded to the isolated effect of different biochar sources, while only the pH showed a significant response to the isolated effect of T inoculation (Table 2). However, T showed a significant interaction with biochar for the variables pH, P, K⁺, Ca²⁺, and H+Al³⁺. The chemical attributes of the control differed from those of all the other treatments, the only exception being Na⁺ content. The application of BH with T increased P (4.7 times), Ca²⁺ (1.2 times) and Mg²⁺ (2.4 times).

Biochar applied to the soil increased Ure, except in the treatment CG (T+). In addition, Ure activity in soils with CH and BH was higher than that of soils with CG and the opposite was observed in Beta, with its activity being significantly more expressed in CG soils (Table 3). This was the only substrate that provided a significant increase in this activity in relation to the control, regardless of the presence of T. Beta activity was also the only variable among the metabolic attributes whose significantly increased because of the inoculation with T for all biochars, mainly the application of CG with T that increase 1.5 times the Beta activity, showing its relevant participation in the hydrolysis of carbon sources.

The activity of P. Alk showed significant responses in the interaction of biochar x T. In this manner, the treatments diversified the responses of this enzyme, emphasizing that, in soils with CG (T+), there was higher expression of P. Alk, significantly contrasting with CG (T-), with less activity (Table 3).

Soils that received CG showed superior TOC (almost 2 times) content in relation to the control treatments, CH and BH, similar to Beta (Table 3). The soil MBC significantly varied with the interaction between biochar and T because of the highest accumulation of microbial carbon in the soils with BH (T-), significantly contrasting BH (T+) and the control. The metabolic quotient ($q\text{CO}_2$) also showed varied responses, highlighting its significantly higher expression in CG. The T inoculation on CG and BH significantly increased $q\text{CO}_2$, though the inverse was observed in CG. Ordering nMDS had a low stress ($S_m = 0.11$), showing an explanation of the treatment effects (biochar and T) on the metabolic and sensory activities of chemical vectors and biometric attributes (Figure 1). A control treatment was positioned in a quadrant opposite to the vectors related to the P and the main soil bases (Mg²⁺, Ca²⁺, and K⁺), and to the growth attributes of plants, in which it showed that this treatment had the lowest values. These variables showed a higher association of BH with MBC and with Ure activity in the soil that pointed to a higher contribution of BH in nutrient accumulation and N cycling in the development of plants and soil microbial communities.




Figure 1. Dissimilarities between the metabolic activities of microorganisms in soils enriched with biochar from bean husks (BH), coffee grounds (CG), and coffee husks (CH), with (T+) and without (T-) *T. aureoviride*. Nonmetric multidimensional scaling (nMDS) showed a reasonable fit of the explanatory model, according to the stress measure ($S_m = .113$, $K_{(dimension)} = 3$) by Kruskal (1964). The vectors corresponding to the chemical attributes (black) and biometrics (gray) followed by asterisks (*) have significant correlations ($\rho \leq 0.05$)

Urease (Ure), β -glycosidase (Beta), alkaline phosphatase (P.Alk), acid phosphatase (P.Aci), total organic carbon (TOC), microbial biomass carbon (MBC), soil metabolic quotient (qCO_2); root dry matter (RDM) and root fresh matter (RFM), shoot dry mass (SDM), and shoot fresh matter (SFM), length (LEN) and diameter (DIA) of watermelon branches

Soils treated with CG biochar showed a higher RFB and levels of $H+Al^{3+}$, as these vectors are negatively correlated with soil pH. Because the independent vectors RFB and Al^{3+} were opposed to each other, the negative effect of the increased Al^{3+} exchangeable on the root development was reiterated, showing that the approximation between RFB and $H+Al^{3+}$ is more related to the H^+ concentrations in the soil and not to Al^{3+} , as expected. In addition, soils with CG showed the highest responses for P. Aci, Beta, and TOC. In general, the metabolic activities of the soils with CG distanced themselves from the soils with CH and BH; these dissimilarities are primarily explained by the differences that these sources exert in soil chemical attributes.

The ANOSIM test confirmed the occurrence of strong segregations among the biometric and chemical attributes (Table 4). In general, all the attributes observed in the control distanced themselves without overlapping their correspondents in the other treatments ($R_{Anosim} > .75$), with the exception of the biometric attributes of plants grown on soils with CH (T-), although they have still distanced themselves ($R_{Anosim} = .56$, Table 4).

Table 4. Analysis of similarity of attributes from coffee grounds (CG), coffee husks (CH), and bean husks (BH) with (+T) and without (-T) inoculation of *Trichoderma aureoviride*, under cultivation of watermelon plants of the variety Vereda style

	CH (T+)	CG (T+)	BH (T+)	CH (T-)	CG (T-)	BH (T-)
<i>a. Metabolic attribute (R-Global = 0.86; p < 0.0001)</i>						
Control	<u>1.00</u>	<u>0.93</u>	<u>1.00</u>	<u>1.00</u>	<u>1.00</u>	<u>1.00</u>
CH (T+)		<u>1.00</u>	0.70	0.56	<u>1.00</u>	0.41
CG (T+)			<u>1.00</u>	<u>1.00</u>	0.56	<u>1.00</u>
BH (T+)				<u>0.96</u>	<u>1.00</u>	0.67
CH (T-)					<u>1.00</u>	<u>0.96</u>
CG (T-)						<u>1.00</u>
<i>b. Biometric attribute (R-Global = 0.58; p < 0.0001)</i>						
Control	<u>1.00</u>	<u>1.00</u>	<u>1.00</u>	0.56	<u>1.00</u>	<u>1.00</u>
CH (T+)		<u>0.96</u>	<u>0.93</u>	0.56	0.22	<u>0.93</u>
CG (T+)			0.30	0.56	<u>1.00</u>	-0.30
BH (T+)				0.56	<u>0.89</u>	0.19
CH (T-)					0.56	0.56
CG (T-)						<u>1.00</u>
<i>c. Chemical attribute (R-Global = 0.59; p < 0.0001)</i>						
Control	<u>1.00</u>	<u>0.89</u>	<u>1.00</u>	<u>1.00</u>	<u>1.00</u>	<u>1.00</u>
CH (T+)		<u>1.00</u>	0.44	0.56	<u>1.00</u>	0.41
CG (T+)			0.37	<u>1.00</u>	-0.22	<u>1.00</u>
BH (T+)				0.37	0.26	-0.04
CH (T-)					<u>1.00</u>	0.30
CG (T-)						<u>1.00</u>

The analyses were made through the Euclidean distance applied to the normalized matrices of the metabolic (a), biometric (b), and chemical (c) attributes. These attributes were modeled in a double factorial scheme ($Y \sim \text{biochar} * T. \text{aureoviride}$) and submitted to permutational multivariate analysis of variance through the Adonis pseudo-F significance test, where segregations (r^2) with $p \leq .05$ are significant: Metabolic (a) biochar ($r^2 = .64, p \leq .001$), *T. aureoviride* ($r^2 = .008, p \leq .785$), and biochar x *T. aureoviride* ($r^2 = .10; p \leq .021$); Biometrics (b) biochar ($r^2 = .57, p \leq .001$), *T. aureoviride* ($r^2 = .09, p \leq .003$), and biochar x *T. aureoviride* ($r^2 = .11; p \leq .006$); ($r^2 = .022, p \leq .665$), and biochar x *T. aureoviride* ($r^2 = .022, p \leq .665$), and *T. aureoviride* ($r^2 = .025, p \leq .231$)

Analyses of the correlations between the scores of the two major components of nMDS showed that the first relevant ordering (nMDS1) had significant positive correlations with the chemical attributes pH, P, K⁺, Ca²⁺, and Mg²⁺ and had negative correlations with Al³⁺ (Table 5). The second ordering (nMDS2) showed strong positive correlations with all biometric attributes, and were significant for LEN, SFB, SDB, and RDB (Table 5). Thus, the dispersions of the metabolic pattern of the microbial communities are primarily explained by the chemical attributes of the soil and then by the responses of the plants in terms of development because of the chemical changes in the soil.

Table 5. Correlation coefficients (r^2) between the explanatory variables and the first two axes of non-parametric multidimensional scaling (nMDS: non-metric multidimensional scaling)

	nMDS1	nMDS2	$r^2(*)$	$P_r(>r^+)$	
<i>a. Metabolic attribute</i>					
Ure	0.999	-0.051	0.80	0.001	***
Beta	-0.967	0.255	0.81	0.001	***
P.Alk	-0.472	-0.882	0.63	0.001	***
P.Aci	-0.586	0.810	0.50	0.002	**
TOC	-0.374	0.928	0.53	0.001	***
MBC	0.678	0.735	0.56	0.001	***
qCO ₂	-0.209	-0.978	0.85	0.001	***
<i>b. Biometric attribute</i>					
LEN	0.153	0.988	0.33	0.026	*
DIA	0.218	0.976	0.17	0.174	ns
SFM	0.115	0.993	0.45	0.002	**
RFM	-0.591	0.807	0.07	0.538	ns
SDM	0.130	0.992	0.29	0.033	*
RDM	0.133	0.991	0.39	0.008	**
<i>c. Chemical attribute</i>					
pH	0.995	-0.103	0.72	0.001	***
Na ⁺	-0.267	-0.964	0.02	0.865	ns
K ⁺	1.000	-0.025	0.59	0.002	**
P	0.848	0.530	0.49	0.002	**
H+Al	-0.999	-0.055	0.65	0.001	***
Ca ²⁺	0.911	0.412	0.36	0.024	*
Mg ²⁺	0.819	0.574	0.56	0.003	**
Al ³⁺	0.304	-0.953	0.13	0.300	ns

*Adjustment statistics by squared correlation coefficient (r^2). + Significance intervals (Pr): 0 *** 0.001 ** 0.01 * 0.05 ns

1.Ure= soil Urease; Beta= β -glycosidase; P.alk= alkaline phosphatase; P.aci=acid phosphatase activities; TOC= total organic carbon; MBC= microbial biomass carbon; qCO₂= soil metabolic quotient; LEN= plant length; DIA= diameter; SFM= shoot fresh mass; RFM= root fresh mass; SDM= shoot dry mass; RDM= root dry mass

The Mantel complementary test confirmed significant positive correlations between the metabolic and biometric attributes ($r = 0.32, \rho = 0.003$), metabolic and chemical attributes ($r = 0.53, \rho = 0.001$), and chemical and biometric attributes ($r = 0.34, \rho = 0.009$). By internally analysing these interactions through the matrices with the Pearson coefficient (R_p), it was possible to evaluate the degree of dependence between the variables associated with the pairs (Figure 2).




Figure 2. Correlations between enzymatic activities, soil chemical attributes, and biometric data of watermelon plants grown on soils under different biochar treatments with and without *Trichoderma aureoviride*. The heatmap and Pearson correlation coefficients (R_p) station were used to associate the paired samples. The underlined correlations are significant ($p \leq .05$). In general, the Mantel test revealed significant positive correlations between soil metabolic attributes and plant biometric (A, $r = 0.32$, $p = 0.003$), soil metabolic, and chemical attributes (B, $r = .53$, $p = 0.001$), and chemical and biometric attributes (C, $r = 0.34$, $p = 0.009$)

Urease (Ure), β -glycosidase (Beta), alkaline phosphatase (P.alk), acid phosphatase (P.aci), total organic carbon (TOC), microbial biomass carbon (MBC), soil metabolic quotient (qCO₂); root dry matter (RDM) and root fresh matter (RFM), shoot dry mass (SDM), and shoot fresh matter (SFM), length (LEN) and diameter (DIA) of watermelon branches

Regarding the interaction between metabolic and biometric attributes, soil MBC showed positive correlations with RDB, SDB, and LEN of the watermelon plants (Figure 2A). The increase in the TOC in the soil was accompanied by an increase in SFB and P. alk correlated with SDB. Metabolic and chemical attribute interactions highlighted the significant and positive correlations of soil Ure activity with pH, Mg²⁺, P, K⁺, and Ca²⁺ content and the negative correlation with H+Al³⁺. MBC also showed positive correlations with Mg²⁺ and P (Figure 2B). Regarding the interaction between chemical and biometric attributes (Figure 2C), the significant and positive correlations of Mg²⁺ with RDB and SDB and P with SDB were highlighted. The only remaining significant correlation was between Al³⁺ and SDB, demonstrating the negative effect of exchangeable Al³⁺ on the labile P and on the growth of the aerial part of the plants.

All the metabolic variables significantly responded to the isolated effect of the biochar source, the only exception begin P. Alk. The isolated effect of *T. aureoviride* inoculation was significant only for Beta and qCO₂ (metabolic attributes). However, the effect of the interaction of biochar $\times$ T was significant for Ure, Beta, TOC, and MBC, showing the potential of this combination in these processes.

Discussion

The different sources of organic matter used to produce the biochar generated products with different properties, mainly in the chemical attributes of the soils that received the biochar (Figure 1). Several studies showed that biochar sources as manure, coffee bark, and bean hulls that are rich in nutrients, sawdust, pine bark, and sugar cane bagasse can provide diverse agronomic and environmental benefits such as a high content of KHCO₃ and CaCO₃ associated with the ash of these materials, increasing the pH and Ca²⁺ content in the soils. The soil attributes can vary by source as shown in previous studies (Domingues *et al.*, 2017; Lee *et al.*,

2019; Song *et al.*, 2019). Here, we used three different sources, two from coffee and one from bean residues, that also showed differences in these and other soil chemical attributes, confirming that biochar production source influences soil fertility. The sandy soil cultivating watermelon that received biochar CG showed fewer macronutrients because it is a thermally pre-processed source; part of its nutrients is removed after infusion into water. However, its participation in the activation of the enzymes β -glucosidase and acid phosphatase was significantly higher than that found in CH. B-glucosidase facilitates the decomposition of low molecular weight carbohydrates, producing glucose, which is fundamental in the terrestrial cycle of C to supply the energy necessary for the proliferation of soil microorganisms (Eivazi and Tabatabai, 1990). Therefore, CG was a significant stimulant of β -glucosidase producing microbial communities. Notably, the increase in the activity of this enzyme accompanied the TOC (substrate) and the activity of acid phosphatase, according to the positive and significant correlations of these variables with the second ordering of nMDS (Figure 1, Table 5).

The similarity between the expressions of β -glucosidase and acid phosphatase in soils that received CG can be explained by the fact that these enzymes mitigate the discharge of bioavailable nutrients of organic carbon and phosphorus in the soil (Sinsabaugh *et al.*, 2014), which were stimulated by the addition of organic materials as shown in two European long-term field experiments (Böhme *et al.*, 2005). Positive and significant correlations between phosphatases and organic carbon are already well known from a previous study (Kumari *et al.*, 2017). However, explanations for these mechanisms are not well justified.

In view of these positions and different enzymatic expressions, the present study proposed the hypothesis that the biochar of CG selects groups of distinct microorganisms that are favoured by biochars with more nutrients available - CG and BH. The higher aggregate expression of β -glucosidase and acid phosphatase in soils with CG corroborates this assertion, as these biochars led to a higher metabolism of organic materials by microorganisms in search of energy. In addition to the nature of the carbon source from the coffee biochars, the optimal pH range provided by these sources may explain the higher enzymatic activities. Acosta-Martínez and Tabatabai (2000) studied enzymatic activity in soils under different doses of limestone in Iowa (USA) and demonstrated that the activity of acid phosphatase was significantly and negatively correlated to soil pH, with optimal performance at pH 6.5. This result corroborates that of the present study (Figure 2B), where soils with CG and BH showed a relative increase in acid phosphatase activity, contrasting with the CH biochar, which greatly increased the soil pH (Tables 2, 3).

Organic waste accounts for most of the monoester phosphates in the soil, which is among the main organic sources of P (Turner, 2008). Thus, in acidic soils with low levels of nutrients and inorganic P, microorganisms are more dependent on the P forms associated with organic materials (Gatiboni *et al.*, 2008). However, these metabolized compounds have a high affinity for soil colloids, where most of the microbiota cannot access them (Rheinheimer *et al.*, 2001). In this aspect, biochar is once again advantageous because it can adsorb the soil nutrients on its surface, keeping them bioavailable for incorporation by microbial biomass or plants during phenolic stages of higher energy demand (Elzobair *et al.*, 2016; Yang *et al.*, 2017). In addition, the fungi *Trichoderma* spp. can also assist in plant nutrition by promoting the solubilization of some minerals, producing phytohormones, and increasing the efficiency of water use, defence, health, and plant development (Shoresh *et al.*, 2010; Khan and Mohiddin, 2018).

In this context, MBC showed a relationship with total organic C and with acid phosphatase through approximations between both in the heatmaps (Figure 2A, 2B) and its strong positive and significant correlations with the second nMDS axis (Table 5). This relationship was discussed in Zhu *et al.* (2017), in which the addition of corn straw biochar under low pyrolysis (450 °C) in a dryland corn field in Northwest China increased the microbial biomass and the activity of phosphatases. Anders *et al.* (2013) showed that these increases were primarily attributed to the increase in C and the nutrients provided by biochar in the soil or by root activity. The present study also showed a greater approximation of RFB gain with TOC, in addition to acid phosphatase (Figure 1). This corroborates the work by Amendola *et al.* (2017), who verified the increase

in organic carbon and biomass of fine roots of grapevine plants immediately after the application of only 10 t ha⁻¹ of biochar.

Urease was the enzyme that most directly correlated with the soil chemical attributes (Figure 2B), which is similar to the result of Kandeler *et al.* (2006) in the 0-10 cm layer of a pasture soil in a semiarid region of northern Colorado (EUA, 25 a 60 NH⁴⁺-Ng⁻¹ 2 h⁻¹). Urease is a metalloenzyme (EC 3.5.1.5) that performs the hydrolysis of urea, releasing CO₂ and NH₃ with increased activity to pH 10 (Kandeler and Gerber, 1988). This explains their higher activity in soils with CH and BH, sources that increased the soil pH, tending to the optimal level of performance (pH 10). This phenomenon is primarily attributed to of Ca²⁺ and K⁺ carbonates, abundant in CH and BH biochar, which are also rich in P and Mg²⁺, such that the pH tends to increase along with the content of these nutrients in the soil, as its potential acidity decreases (H⁺+Al³⁺, Figure 2B).

Notably, there are high positive and significant correlations of labile P and Mg²⁺ with CBM (Figure 2B). Regarding P, this correlation is justified by its importance as a mediator in the energy transfer processes in the form of ATP and in the synthesis of nucleic acids, phospholipids, and molecular signals in the path of enzymatic phosphorylation. Mg²⁺ is also a cofactor and fundamental regulator of countless proteins with the presence of fundamental P in its transport to the interior of the prokaryotic cells. This is explained by extracellular Mg²⁺ with ATP as cofactors of Mg²⁺ hydrolase, importing ATPase (EC 3.6.3.2), the enzyme responsible for the release of phosphate and M²⁺ inside cells (Tao *et al.*, 1995). In addition, in the present study, these nutrients were the only ones that significantly affected the dry mass gain of the shoot and roots (in the case of Mg) and the fresh mass of the aerial part in the case of P (Figure 2C). In summary, the mechanisms of these interactions represent some of the several that reflect the importance of PO₄³⁻ and Mg²⁺ availability for the development of watermelon plants and associated microbial communities.

In view of the variability between the responses to the different biochar sources on the chemical and enzymatic attributes of the soils, the choice of the best source is dependent on the purpose of its use. Although the CH and BH biochars provide more labile P and bases to the soil, increasing the pH, the CG biochar is more relevant for the activation of acid phosphatase and β -glucosidase and an increase in soil organic carbon. In addition, all sources presented different relevance when considering inoculation with *T. aureoviride* (Figure 1), primarily on the development of plants (Table 1).

In general, the most significant contributions of *T. aureoviride* inoculation were the following (Table 1, 2, 3): I - increased fresh root mass, regardless of biochar; II - increased fresh and dry mass of the aerial part, in treatments with CG and BH; III - increased length of the main branches in treatments with CG; IV - increased branch diameter in treatments with CG and BH; and V - increased root dry biomass in treatments with CG and primarily with BH.

Similarly, Jalali *et al.* (2017) demonstrated the potential to promote *Arabidopsis thaliana* growth because of increased tolerance to salinity stress after inoculation with several species of *Trichoderma* spp., protecting plants against oxidative damage. Kiriga *et al.* (2018) also demonstrated that *Trichoderma* spp. can endophytically colonize the pineapple plant roots, protecting the culture against infestations of *Meloidogyne javanica* and fungi in the soil and increasing root mass and plant development. In general, there are numerous studies mentioning other benefits provided by several species of *Trichoderma* spp., such as increased root uptake (Oskiera *et al.*, 2017), seed germination (Mendoza-Mendoza *et al.*, 2018), and soil metabolic activity (Elzobair *et al.*, 2016), among others. These references corroborate the significant increase in β -glucosidase activity as a function of *T. aureoviride* inoculation, independent of the biochar used, with greater relevance for CG (Table 3).

Considering only the chemical contributions, the greatest change was observed in soils with *T. aureoviride* inoculated with BH, where there was a significant increase in the levels of labile P (Mehlich⁻¹) and Ca²⁺ (Table 2). Corroborating this result, Saravanakumar *et al.* (2013) demonstrated that all ten species of *Trichoderma* spp., isolated from the rhizosphere *Avicennia marina*, a native species of mangroves, solubilized tricalcium phosphate (Ca₃(PO₄)₂). This insoluble source of P induces the synthesis of the extracellular enzyme

phytase by the isolates. The same study also showed that the re-inoculation of these isolates increased the biomass of *A. marina* cultivated in poor soil (*in vitro*) without phosphate supplementation by 19 %. Thus, this reference explains the simultaneous increases in exchangeable Ca and P content in soils with *T. aureoviride* and BH, in addition to the enormous enzymatic versatility and potential of vegetal promotion of this species.

These changes in the means of the variables analyzed in response to inoculation with *T. aureoviride* are because of the physicochemical specificity of each material used as they affect soil aggregation (Amendola *et al.*, 2017) and microbial community structure (Zhang *et al.*, 2018).

Conclusions

This study showed a new approach of biochar as a route of *T. aureoviride* inoculation. Watermelon plants cultivated in soil with coffee ground biochar and inoculated with *T. aureoviride* have increased the plant length and shoot dry biomass by 129% and 192%, respectively, compared to the control. The use of *T. aureoviride* via inoculation using different biochar sources showed sufficient characteristics for the development of a biological substrate with a high potential for the development of watermelon plants and improvement in the soil quality. This is even more relevant when considering the low production costs of these substrates compared to that of chemical fertilizers.

Acknowledgements

We thank fellowships and grants from CNPq (306401/2015-0 465764/2014-2 CNPq-441305/2017-2), CAPES (88887.736369/2017-00), FACEPE (APQ-0223-5.01/15; APQ-0419-5.01/15; APQ-0431-5.01/17; APQ-0498-3.07/17) and Micoteca URM collection of the Mycology Department of the Biological Sciences Center of the Federal University of Pernambuco for provided us the *Trichoderma aureoviride* URM 5158 isolate.

Conflict of Interests

The authors declare that there are no conflicts of interest related to this article.

References

- Acosta-Martínez V, Tabatabai MA (2000). Enzyme activities in a limed agricultural soil. *Biology and Fertility of Soils* 31:85-91. <https://doi.org/10.1007/s003740050628>
- Amendola C, Montagnoli A, Terzaghi M, Trupiano D, Oliva F, Baronti S, Scippa GS (2017). Short-term effects of biochar on grapevine fine root dynamics and arbuscular mycorrhizae production. *Agriculture, Ecosystems and Environment* 239:236-245. <https://doi.org/10.1016/j.agee.2017.01.025>
- Anders E, Watzinger A, Remt F, Kitzler B, Wimmer B, Zehetner F (2013). Biochar affects the structure rather than the total biomass of microbial communities in temperate soils. *Agricultural and Food Science* 22:404-423. <https://doi.org/10.23986/afsci.8095>
- Anderson TH, Domsch KH (1985). Determination of eco-physiological maintenance carbon requirements of soil microorganisms in a dormant state. *Biology and Fertility of Soils* 1(2):81-89. <https://doi.org/10.1007/BF00255134>

- Bartlett RJ, Ross DS (1988). Colorimetric determination of oxidizable carbon in acid soil solutions. Soil Science Society of America Journal 53:191-192. <https://doi.org/10.2136/sssaj1988.03615995005200040055x>
- Böhme L, Langer U, Böhme F (2005). Microbial biomass, enzyme activities and microbial community structure in two European long-term field experiments. Agriculture, Ecosystems and Environment 109:141-152. <https://doi.org/10.1016/j.agee.2005.01.017>
- Domingues RR, Trugilho PF, Silva CA (2017). Properties of biochar derived from wood and high-nutrient biomasses with the aim of agronomic and environmental benefits. Plos One 12 (5):e0176884. <https://doi.org/10.1371/journal.pone.0176884>
- Eivazi F, Tabatabai MA (1990). Factors affecting glucosidase and galactosidase activities in soils. Soil Biology and Biochemistry 22:891-897. [https://doi.org/10.1016/0038-0717\(90\)90126](https://doi.org/10.1016/0038-0717(90)90126)
- Eivazi F, Tabatabai M A (1977). Phosphatases in soils. Soil Biology and Biochemistry 9:167-172. [http://dx.doi.org/10.1016/0038-0717\(77\)90070-0](http://dx.doi.org/10.1016/0038-0717(77)90070-0)
- Eivazi F, Tabatabai MA (1988). Glucosidases and galactosidases in soils. Soil Biology and Biochemistry 20(5):601-606. [https://doi.org/10.1016/0038-0717\(88\)90141-1](https://doi.org/10.1016/0038-0717(88)90141-1)
- Elzobair KA, Stromberger ME, Ippolito JA, Lentz RD (2016). Contrasting effects of biochar versus manure on soil microbial communities and enzyme activities in an Aridisol. Chemosphere 142:145-152. <https://doi.org/10.1016/j.chemosphere.2015.06.044>
- Gatiboni LC, Rheinheimer DS, Brunetto GB (2008). Soil microbial biomass phosphorus and activity of acid phosphatases during decline of soil available phosphorus. Pesquisa Agropecuária Brasileira 43(8):1085-1091.
- Jalali F, Zafari D, Salari H (2017). Volatile organic compounds of some *Trichoderma* spp. increase growth and induce salt tolerance in *Arabidopsis thaliana*. Fungal Ecology 29:67-75.
- Khan MR, Mohiddin FA (2018). *Trichoderma*: its multifarious utility in crop improvement. In: Crop Improvement Through Microbial Biotechnology. Elsevier pp 263-291.
- Kandeler E, Gerber H (1988). Short-term assay of soil urease activity using colorimetric determination of ammonium. Biology and Fertility of Soils 6:68-72. <http://dx.doi.org/10.1007/BF00257924>
- Kandeler, E, Mosier A, Morgan J (2006). Response of soil microbial biomass and enzyme activities to transient elevation of carbon dioxide in a semi-arid grassland. Soil Biology and Biochemistry 38:2448-2460. <https://doi.org/10.1016/j.soilbio.2006.02.021>
- Kiriga AW, Haukeland S, Kariuki GM, Coyne DL, Beek NV (2018). Effect of *Trichoderma* spp. and *Purpureocillium lilacinum* on *Meloidogyne javanica* in commercial pineapple production in Kenya. Biological Control 119:27-32.
- Kumari A, Rao PC, Padmaja G, Madhavi M (2017). Effect of physico-chemical properties on soil enzyme acid phosphatase activity of some soils in vegetable growing soils of Ranga Reddy District of Telangana State, India. International Journal of Current Microbiology and Applied 6(10):3496-3503. <https://doi.org/10.20546/ijcmas.2017.610.412>
- Lee ME, Park JH, Chung JW (2019). Comparison of the lead and copper adsorption capacities of plant source materials and their biochars. Journal of Environmental Management 236:118-124. <https://doi.org/10.1016/j.jenvman.2019.01.100>
- Lima JRS, Silva WM, Medeiros EV, Duda GP, Corrêa MM, Martins Filho AP, Hammecker C (2018). Effect of biochar on physicochemical properties of a sandy soil and maize growth in a greenhouse experiment. Geoderma 319:14-23. <https://doi.org/10.1016/j.geoderma.2017.12.033>
- Mendonca ES, Matos ES (2005). Matéria orgânica do solo: Métodos de análises. Universidade Federal de Viçosa (UFV), Viçosa, Brasil pp 86-92.
- Mendoza-Mendoza A, Zaid R, Lawry R, Hermosa R, Monte E, Horwitz BA, Mukherjee PK (2018). Molecular dialogues between *Trichoderma* and roots: Role of the fungal secretome. Fungal Biology Reviews 32:62-85.
- Oskiera M, Szczec M, Stępowaska A, Smolińska U, Bartoszewski G (2017) Monitoring of *Trichoderma* species in agricultural soil in response to application of biopreparations. Biological Control 113:65-72. <https://doi.org/10.1016/j.biocontrol.2017.07.005>
- Patel A, Khare P, Patra DD (2017). Biochar mitigates salinity stress in plants. In: Shukla V, Kumar S, Kumar N (Eds). Plant adaptation strategies in changing environment. Springer, Singapore, pp 153-182.

- Poduria A, Rateria DL, Sahab SK, Sahac S, Daugherty A (2013). *Citrullus lanatus* 'sentinel' (watermelon) extract reduces atherosclerosis in LDL receptor-deficient mice. The Journal of nutritional biochemistry 24:882-886. <https://doi.org/10.1016/j.jnutbio.2012.05.011>
- Raij B, Van Andrade JC, Cantarella H, Quaggio JA (2001). Análise química para avaliação da fertilidade de solos tropicais. Campinas: Instituto Agronômico pp 284.
- Rheinheimer DS, Anghinoni I (2001). Distribution of inorganic phosphorus fraction in soil management systems. Pesquisa Agropecuária Brasileira 36:151-160.
- Saravanakumar K, Arasu VS, Kathiresan K (2013). Effect of *Trichoderma* on soil phosphate solubilization and growth improvement of *Avicennia marina*. Aquatic Botany 104:101-105. <https://doi.org/10.1016/j.aquabot.2012.09.001>
- Saxena, J, Rana G, Pandey M (2013). Impact of addition of biochar along with *Bacillus* sp. on growth and yield of French beans. Scientia Horticulturae 162:351-356. <https://doi.org/10.1016/j.scienta.2013.08.002>
- Shores M, Harman GE, Mastouri F (2010). Induced systemic resistance and plant responses to fungal biocontrol agents. Annual Review of Phytopathology 48:21-43. <https://doi.org/10.1146/annurev-phyto-073009-114450>
- Silva FC (2009). Manual de análises químicas de solos, plantas e fertilizantes. 2.ed. Brasília, DF: Embrapa Informação Tecnológica; Rio de Janeiro: Embrapa Solos, pp 627.
- Silva JAT, Medeiros EV, Silva JM, Tenório DDA, Moreira KA, Nascimento TCEDS, Souza-Motta C (2016). *Trichoderma aureoviride* URM 5158 and *Trichoderma hamatum* URM 6656 are biocontrol agents that act against cassava root rot through different mechanisms. Journal of Phytopathology 164:1003-1011. <https://doi.org/10.1111/jph.12521>
- Sinsabaugh RL, Belnap J, Findlay SG, Shah JFF, Hill BH, Kuehn KA (2014). Extracellular enzyme kinetics scale with resource availability. Biogeochemistry 121:287-304. <https://doi.org/10.1007/s10533-014-0030-y>
- Song B, Chen M, Zhao L, Qiu H, Cao X (2019). Physicochemical property and colloidal stability of micron-and nano-particle biochar derived from a variety of feedstock sources. Science of the Total Environment <https://doi.org/10.1016/j.scitotenv.2019.01.193>
- Tao T, Snively MD, Farr SG, Maguire ME (1995). Magnesium transport in *Salmonella typhimurium*: mgtA encodes a P-type ATPase and is regulated by Mg²⁺ in a manner similar to that of the mgtB P-type ATPase. Journal of Bacteriology 177(10):2654-62. <https://doi.org/10.1128/jb.177.10.2654-2662.1995>
- Tate KR, Ross DJ, Feltham CW (1988). A direct extraction method to estimate soil microbial C: effects of experimental variables and some different calibration procedures. Soil Biology and Biochemistry 20:329-335. [https://doi.org/10.1016/0038-0717\(88\)90013-2](https://doi.org/10.1016/0038-0717(88)90013-2)
- Turner BL (2008) Resource partitioning for soil phosphorus: a hypothesis. Journal of Ecology 96:698-702. <https://doi.org/10.1111/j.1365-2745.2008.01384.x>
- Yang X, Meng J, Lan Y, Chen W, Yang T, Yuan J, Han J (2017). Effects of maize stover and its biochar on soil CO₂ emissions and labile organic carbon fractions in Northeast China. Agriculture, Ecosystems and Environment 240:24-31. <https://doi.org/10.1007/s11356-017-8500-0>
- Yeomans JC, Bremner JM (1988). A rapid and precise method for routine determination of organic carbon in soil. Communications in Soil Science and Plant Analysis 19:1467-1476. <https://doi.org/10.1080/00103628809368027>
- Zhang P, Sun H, Min L, Ren C (2018). Biochars change the sorption and degradation of thiacloprid in soil: Insights into chemical and biological mechanisms. Environmental Pollution 236:158-167. <https://doi.org/10.1016/j.envpol.2018.01.030>
- Zhu LX, Xiao Q, Cheng HY, Shi BJ, Shen YF, Li SQ (2017). Seasonal dynamics of soil microbial activity after biochar addition in a dryland maize field in North-Western China. Ecological Engineering 104:141-149. <https://doi.org/10.1016/j.ecoleng.2017.04.026>



The journal offers free, immediate, and unrestricted access to peer-reviewed research and scholarly work. Users are allowed to read, download, copy, distribute, print, search, or link to the full texts of the articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

License - Articles published in *Notulae Botanicae Horti Agrobotanici Cluj-Napoca* are Open-Access, distributed under the terms and conditions of the Creative Commons Attribution (CC BY 4.0) License.

© Articles by the authors; UASVM, Cluj-Napoca, Romania. The journal allows the author(s) to hold the copyright/to retain publishing rights without restriction.