# Nod Factor Thin-Layer Chromatography Profiling as a Tool to Characterize Symbiotic Specificity of Rhizobial Strains: Application to *Sinorhizobium saheli, S. teranga,* and *Rhizobium* sp. Strains Isolated from *Acacia* and *Sesbania*

Gilles Lortet,<sup>1,2</sup> Nathalie Méar,<sup>1</sup> Jean Lorquin,<sup>1</sup> Bernard Dreyfus,<sup>1</sup> Philippe de Lajudie,<sup>1</sup> Charles Rosenberg,<sup>2</sup> and Catherine Boivin<sup>1</sup>

<sup>1</sup>Laboratoire de Microbiologie, ORSTOM, BP1386, Dakar, Sénégal; <sup>2</sup>Laboratoire de Biologie Moléculaire des Relations Plantes-Microorganismes, CNRS-INRA, B. P. 27, 31 326 Castanet-Tolosan Cedex, France Received 2 May 1996. Accepted 25 June 1996.

Rhizobia isolated from Acacia or Sesbania belong to several taxonomic groups, including the newly described species Sinorhizobium saheli, Sinorhizobium teranga, and the so-called cluster U. A collection of strains belonging to these different groups was analyzed in order to determine whether the host range of a strain could be correlated with various molecular nodulation determinants. Nodulation tests showed that, independently of their taxonomic position, all the strains isolated from the same plant genus exhibited a similar host range, which was different for Sesbania and Acacia isolates. The fact that S. teranga strains nodulate either Acacia or Sesbania led us to subdivide this species into biovars acaciae and sesbaniae. Thin-layer chromatography (TLC) analysis of the Nod factors synthesized by overproducing strains showed that (i) strains isolated from the same plant genus exhibited similar TLC profiles and (ii) profiles of Acacia and Sesbania symbionts were easily distinguishable, Acacia strains producing, in particular, sulfated molecules. In contrast, no correlation could be established between the host range of a strain and its plasmid content, the nature of the nod gene inducers or the presence of DNA sequences homologous to specific nod genes. We thus propose that Nod factor TLC profiling may be used as an easy and powerful tool for the classification of rhizobial strains on the basis of their symbiotic properties.

#### Additional keywords: nitrogen fixation.

Soil bacteria which symbiotically interact with leguminous plants to form nitrogen-fixing nodules are very diverse in regard to their taxonomic position. These bacteria, termed rhizobia, have been shown to belong to about 15 different named species and several unamed genomic species, distributed in the genera *Rhizobium*, *Bradyrhizobium*, *Azorhizobium*, and *Sinorhizobium* (see Young 1994; de Lajudie et al. 1994). In

phylogenetic trees, these genera are intermixed with nonsymbiotic bacteria like Agrobacterium or Rhodopseudomonas (see Young 1994; Martinez-Romero 1994). The current classification is mainly based on universally approved techniques in microbial taxonomy, like auxanographic tests, DNA:DNA and DNA:rRNA hybridizations, whole-cell protein SDS-PAGE, and 16S rRNA sequencing (Graham et al. 1991). Such a classification no longer considers symbiotic properties as a taxonomic criterion, since the ability to nodulate a given legume species can be shared by taxonomically dissimilar symbionts, especially in the case of tropical legumes. For example, soybean is nodulated by B. japonicum (Jordan 1984), S. fredii (Jarvis et al. 1992), and B. elkanii (Kuykendall et al. 1992). Conversely, strains belonging to the same species can exhibit different host spectra. For example, in R. leguminosarum, three different biovars have been defined on the basis of their host range (Jordan 1984). Therefore, the host range of a given strain cannot be predicted from its taxonomic position. When rhizobial strains need to be classified on the basis of their host specificity, as in agro-ecological studies, or for analysis of biodiversity and plant-bacteria coevolution, their host range is analyzed by nodulation tests on various potential host plants. However, such experiments, though representing heavy work, provide only a partial characterization of the host range, and can be poorly discriminating, depending on the number and on the choice of the hosts tested.

To develop more accurate tools for the characterization of the symbiotic potentialities of a strain, we examined possible correlations between the host range and diverse elements involved in the molecular dialogue which controls the recognition process between the bacterial symbiont and its host plant. The bacterial nodulation genes (*nod, nol,* and *noe*), which are generally located on large plasmids in *Rhizobium* spp., play a central role in this molecular dialogue (Fellay et al. 1995; Dénarié et al. 1996). These genes are expressed in response to plant inducers present in root exudates, generally flavonoids. The induction process is mediated by the products of the *nodD* regulatory genes, which bind to conserved DNA consensus sequences located upstream of the *nod* operons, called *nod* 

Corresponding author: C. Boivin; E-mail: boivinc@belair.orstom.sn

736 / Molecular Plant-Microbe Interactions



Fonds Documentaire ORSTOM Cote: B×15493 Ex: ユ boxes (Spaink et al. 1987; Schultze et al. 1994; Fellay et al. 1995). In the presence of appropriate flavonoids, the NodD proteins activate the transcription of the structural nod genes. The recognition between a given NodD protein and particular flavonoids determines a first level of specificity in the symbiotic interaction. The expression of the structural nod genes leads to the production of extracellular bacterial signals, called Nod factors, which are essential for the infection process and nodule organogenesis (for reviews see Dénarié et al. 1992; Fellay et al. 1995). These Nod factors are lipooligosaccharides basically consisting of a chito-oligosaccharide backbone of three to five B-1,4-linked N-acetyl-D-glucosamine residues, substituted by an N-acyl chain at the nonreducing end. Other diverse substitutions are present at both ends of the oligochitin chain. The nature of all these substitutions, which is controlled by various combinations of specific nod genes, make the Nod factors plant specific (Dénarié and Cullimore 1993; Spaink 1994; Dénarié et al. 1996). For instance, in S. meliloti, the *nodH* gene is involved in the addition of the sulfate group, which is required for alfalfa nodulation (Roche et al. 1991; Schultze et al. 1995). In the same way, the nodS and nodU genes of Sinorhizobium sp. NGR234, respectively, determine Nod factor N-methylation and O-carbamoylation, which enable this strain to nodulate Leucaena leucocephala (Lewin et al. 1990; Jabbouri et al. 1995).

Taking advantage of this current understanding of the molecular basis of host specificity, we looked for possible relationships between host range and several bacterial nodulationrelated features. For this study, a collection of rhizobial strains isolated in Senegal from *Acacia* and *Sesbania* appeared of special interest, since strains isolated from the same host belong to different species, whereas strains isolated from different hosts belong to a same species (de Lajudie et al. 1994). We characterized the host range of these strains by testing their ability to nodulate various legumes, and we attempted to establish correlations between this host range and various nodulation-related determinants such as plasmid content, nature of *nod* gene inducers, Nod factor production and presence of DNA sequences homologous to host-specific *nod* genes.

### RESULTS

## Grouping *Acacia* and *Sesbania* isolates on the basis of their host range.

Tropical legumes of the genus Sesbania are known to be nodulated by the highly specific species Azorhizobium caulinodans (Dreyfus et al. 1988). In a recent report, de Lajudie et al. (1994) performed a taxonomic analysis of a collection of rhizobial strains isolated in Senegal either from Sesbania or from Acacia hosts: All the strains classified into the new species Sinorhizobium saheli were isolated from Sesbania, whereas strains belonging to the genospecies Ua and Ub of the so-called cluster U were all isolated from Acacia. Strains of the new species Sinorhizobium teranga were isolated either from Sesbania or from Acacia. S. teranga and S. saheli are taxonomically closely related to R. meliloti and R. fredii, which have been reclassified in the emended genus Sinorhizobium (de Lajudie et al. 1994). Genospecies Ua and Ub belong to the [R.] loti branch, which is currently proposed to form the new genus Mesorhizobium (Lindström et al. 1995).

To classify these rhizobial strains on the basis of their sym-

biotic properties, 42 strains of *S. teranga*, *S. saheli*, genospecies Ua and genospecies Ub originating from either Acacia or Sesbania (Table 1) were inoculated onto several Acacia (A. nilotica, A. senegal, A. raddiana, and A. seyal) and Sesbania (S. pubescens, S. rostrata, and S. grandiflora) species. As cross inoculation between Acacia, Leucaena, and Prosopis has been observed (Zhang et al. 1991), Prosopis juliflora and Leucaena leucocephala were also tested.

Results (Table 1) showed that Acacia isolates (belonging to S. teranga, genospecies Ua or genospecies Ub) and Sesbania isolates (belonging to S. teranga or S. saheli) exhibit distinct host spectra: (i) all Acacia isolates induced nodules on Acacia species (data not shown for A. seval) and most of them formed efficient nodules fixing nitrogen. They also nodulated P. juliflora and L. leucocephala, but failed to induce nodule formation on any of the Sesbania species tested, in the exception of S. rostrata, on which several strains were able to induce pseudonodules. (ii) Sesbania isolates induced nitrogen-fixing nodules on all Sesbania species tested, but formed non-fixing nodules on Acacia seyal (data not shown), and were unable to nodulate A. nilotica, A. senegal, P. juliflora, and L. leucocephala. A. raddiana was poorly nodulated by S. teranga strains. The strain ORS1013 is atypical, since, though isolated from A. senegal, it exhibited on other hosts the same specificity as S. teranga strains isolated from Sesbania (Table 1).

Therefore, in contrast to S. saheli, which only comprises Sesbania strains, the species S. teranga clusters both Acacia and Sesbania isolates with very distinct host ranges. In the following, we will designate S. teranga strains nodulating Acacia as S. teranga by. acaciae and strains nodulating Sesbania as S. teranga by. sesbaniae (Table 1).

## S. teranga strains isolated from Acacia and Sesbania exhibit similar plasmid profiles.

In most fast-growing Rhizobium, the nod genes are located on a large plasmid called the pSym (Rosenberg et al. 1981; Appelbaum et al. 1985) and it has been suggested that transfer of symbiotic plasmids could be responsible for modifications in host specificity (Martinez-Romero 1994; Segovia et al. 1993). Therefore, strains belonging to different species but isolated from the same legume (Acacia or Sesbania) in the same area could be expected to exhibit similarities in their plasmid content. Conversely, S. teranga strains with differerent host ranges could be expected to have different plasmid contents. We thus visualized the plasmids present in the strains by agarose gel electrophoresis (see Materials and Methods). However, our results showed that S. saheli, S. teranga by. sesbaniae, and S. teranga by. acaciae strains could not be distinguished on the basis of their plasmid content (Fig. 1). One band migrating more slowly than pSyma and pSymb from S. meliloti was detectable in all the Sinorhizobium strains tested. As this band was routinely observed using a technique which does not usually allow chromosome visualization, and as no equivalent band could be detected with any of the control strains, we assigned it to a megaplasmid. All the Sinorhizobium strains also contain another large plasmid of more than 350 kb, and a few of them (ORS611, ORS1071, and ORS22) an additional smaller plasmid (data not shown). In contrast, no plasmids were detected in the cluster U strains studied, except for ORS1018 and ORS1020, which both contained one plasmid (data not shown).

## Luteolin is a good *nod* gene inducer for both *S. teranga* and *S. saheli*.

The specific activation of a NodD protein by plant inducers determines a first level of specificity in the symbiotic interaction. We thus screened various flavonoid compounds for their ability to induce *nod* gene expression in strains chosen as representatives of the different taxonomic groups. Identification of *nod* gene inducers was performed using available transcriptional fusions of *E. coli lacZ* gene with the *nodA* promoter of *R. legu* 

minosarum bv. viciae (plasmid pMP240) or bv. trifolii (plasmid pRT311), and the nodY promoter of *B. japonicum* (plasmid pDT900). These fusions were introduced in streptomycinresistant derivatives of *S. saheli* ORS611, *S. teranga* bv. sesbaniae ORS604, *S. teranga* bv. acaciae ORS1073, genospecies Ua ORS1001 and genospecies Ub ORS1040. Various commercial flavonoids, most of them known as nod gene inducers in other *Rhizobium* species (for a review see Fellay et al. 1995), were tested for their effect on  $\beta$ -galactosidase activity in these constructs.

Table 1. Original host plant and host specificity of members of Sinorhizobium teranga, Sinorhizobium saheli, and Rhizobium sp. from cluster U among Sesbania, Acacia, Prosopis, and Leucaena species<sup>a</sup>

				Plant tested					
Rhizobial	Isolated		Sesbania			Acacia		Prosopis	Leucaena
strain	from	grandiflora	rostrata	pubescens	senegal	raddiana	nilotica	juliflora	leucocephala
Sinorhizobium branch	1								
S. saheli									
ORS 600	S. pachycarpa	+	+	+	_	-	-	-	
ORS 609	S. cannabina	+	+	+	-	-	_	NT	
ORS 611	S. grandiflora	+	+	+	-	-	-		-
S. teranga bv. sesba	niae								
ORS 15	S. sp.	+	+	÷	-	± d	-	NT	_
ORS 19	S. cannabina	+	+	+	_	· +	-	NT	
ORS 604	S. aculeata	+	+	+		± d	-	-	-
ORS 613	S. sesban	+	+	+	-	+	-	_	-
ORS 8	S. rostrata	+	+	+	-	± d		NT	-
ORS 22	S. rostrata	+	+	+	-	$\pm d$	-	-	-
ORS 51	S. rostrata	+	+	÷	_	±d	-	NT	-
ORS 52	S. rostrata	+	+	+		± d	-	-	
ORS 53	S. rostrata	+	+	+	-	±d	-	NT	
ORS 1013	A. senegal	+	+	+	+	± d		-	
S. teranga bv. acaci	ae								
ORS 1071	A. senegal	_	-	-	+	+	+	NT	+
ORS 1072	A. senegal	_	–(pn)	-	+	+	+	ᆂ	+
ORS 1073	A. senegal	-	-	_	+	+	+	÷	+
ORS 1007	A. laeta	-	-(pn)	-	+	+	+	NT	+
ORS 1009	A. laeta	-	–(pn)	-	+	+	+	+	+
ORS 1016	A. laeta	-	-	-	+	+	+	+	+
ORS 1045	A. raddiana	-	-		+	+	+	NT	+
ORS 1047	A. horrida	-	-	-	+	+	+	NT	+
ORS 1057	A. mollissima	_	-	-	+	+	+	NT	+
ORS 1058	A. mollissima	_		-	+	+	+	NT	+
ORS 929	A. sp.	-	(pn)		-	+	+	NT	+
[Rhizobium] loti bran	ch *								
Cluster U, genospec	ies Ua#								
ORS 1001	A. senegal	-	-	-	+	+	+ d	+	+
ORS 1004	A. senegal	-		-	+	+	+	NT	+
ORS 1010	A. senegal	-	-	-	+	+	+	NT	+
ORS 1014	A. senegal	-	_	-	+	+	+	NT	+
ORS 1015	A. senegal	_	_	-	+	÷	+	NT	+
ORS 1018	A. senegal	_	-		+	+	+	NT	÷
ORS 1020	A. senegal	_	_	-	+	+	+	NT	+
ORS 1024	A. senegal	-	-(pn)	_	+	+	+	±	±
ORS 1029	A. senegal			-	+	+	+	NT	+
ORS 1035	A. senegal	_	_	-	+	+	+ d	+	+
ORS 1036	A. senegal	-	_		+	+	+	NT	+
ORS 1037	A. senegal	_	_	-	+	+	+	NT	+
Cluster U. genospecies Ubs									
ORS 1030	A. senegal	_	-	-	+	+	+	+	+
ORS 1031	A. senegal	_	_	-	+	+	+	NT	+
ORS 1032	A. senegal	-	_	_	+	+	+	NT	+
ORS 1038	A. senegal	-	-	_	+	+	+	NT	+
ORS 1040	A. senegal	-	_	_	+	+	+ d	±d	+
ORS 1088	A. seyal	-	-		+	+	+	±	+

<sup>a</sup> +, more than 50% of the plant tested were nodulated. -, none of the plants tested were nodulated. ±, between 10% and 50% of the plants tested were nodulated. d, delayed nodulation. NT, not tested. (pn), occasional formation of pseudonodules. \* A new genus, *Mesorhizobium*, has been proposed for the *R. loti* branch (Lindström et al. 1995). # Ua is constituted of subclusters U1 and U2 described in de Lajudie et al. (1994). § Ub comprises strains from subcluster U3 described in de Lajudie et al. (1994). Ua and Ub represent two different genospecies (P. de Lajudie, unpublished).

No induction of the *nodY-lacZ* fusion could be detected in any of the five strains (data not shown). In contrast, significant induction levels were observed in Sinorhizobium strains containing the plasmid pMP240 carrying the R. leguminosarum by. viciae nodA::lacZ fusion (Fig. 2), and, to a lesser extent, the plasmid pRT311 (data not shown). For these strains, several flavonoids, i.e. apigenin, naringenin, butein, and kaempferol had an inducing effect, but the best inducer was the flavone luteolin (Fig. 2). No induction or a very weak induction was observed with daidzein, genistein (Fig. 2), eriodictyol, umbelliferone, myricetin, and formononetin (data not shown). No induction could be detected for genospecies Ua ORS1001 and genospecies Ub ORS1040, bearing either pMP240, pRT311, or pMP194, a S. meliloti nodABC::lacZ fusion (data not shown, except for ORS1001(pMP240), Fig. 2).

## Construction of derivatives overproducing Nod factors.

For the Sinorhizobium representatives, for which nod gene inducers have been identified, we attempted to characterize Nod factor production. S. saheli ORS611, S. teranga by. sesbaniae ORS604, and S. teranga by. acaciae ORS1073 were grown in a medium containing 10 µM luteolin, and <sup>14</sup>C acetate as a Nod factor precursor. Culture supernatants were analyzed by the TLC system previously described (Spaink et al. 1992). No luteolin-inducible spots were detected on TLC plates after a 2-week exposure, indicating a very low level of Nod factor production. This led us to attempt to increase Nod factor production by introducing plasmids bearing nodD genes from other rhizobial species into wild-type strains representative of the five taxonomic groups. We tested different nodD genes: nodD1 of the broad host range strain Sinorhizobium sp. NGR234 (= plasmid pA28), nodD2 of the tree-nodulating strain S. sp. BR816 (= plasmid pVK31), nodD1 of the taxonomically closely related species S. meliloti (= plasmid pMH901), all of them known to activate nod gene expression through flavonoid induction (Bassam et al. 1988; van Rhijn et al. 1994; Maillet et al. 1990) and the nodD3-syrM genes of S. meliloti (plasmid pMH682), which direct flavonoidindependent nod gene transcription (Maillet et al. 1990).

Results (Table 2) showed that *nodD1* of *Sinorhizobium* sp. NGR234 allowed Nod factor overproduction in the five taxonomic groups. In contrast, the inducing effect of the other *nodD* genes was dependent on the species: *nodD3-syrM* genes

of *S. meliloti* were efficient in the *Sesbania*-nodulating strains *S. saheli* ORS611 and *S. teranga* ORS604, whereas *S. meliloti* nodD1 gene increased Nod factor production in both cluster U representatives ORS1001 and ORS1040, which nodulate Acacia. The nodD2 gene of Sinorhizobium sp. BR816 was only active in the *S. teranga* by. acaciae strain ORS1073.

Moreover, introducing plasmids pMH682, pVK31, or pMH901 into other *Sesbania* symbionts, *S. teranga* bv. *acaciae* and cluster U strains, respectively, resulted in the production of detectable Nod factors (see Fig. 3 for examples). Thus, for taxonomically closely related rhizobia, the ability of a strain to respond to a given exogenous *nodD* gene seems to be linked to its host range. However, such a link cannot be observed for taxonomically distant strains.

It has been reported that the nature of the regulatory *nodD* gene can influence Nod factor structure (Demont et al. 1994). Therefore, when possible, we compared the effect of different exogenous *nodD* genes on Nod factor production. However, for a given strain, similar TLC patterns were obtained, whatever the exogenous *nodD* gene responsible for Nod factor overproduction (data not shown).

## Acacia symbionts produce sulfated Nod factors, whereas Sesbania symbionts do not.

Several overproducing derivatives were constructed for each taxonomic group, and their Nod factor production was analyzed by TLC, using <sup>14</sup>C acetate or <sup>35</sup>S sulfate as precursors (see Materials and Methods). In <sup>14</sup>C-labeled S. teranga by. acaciae strains harboring pVK31, several major flavonoidinducible spots were detected in the culture supernatants (Fig. 3A, lanes 1ai, 2ai and 3ai). With <sup>35</sup>S as precursor, one of these spots was strongly labeled, and one or two others appeared as weaker signals (Fig. 3A, lanes 1si, 2si, and 3si). Flavonoidinduced cultures of the Acacia-nodulating genospecies Ua and Ub containing pMH901, exhibited an analogous TLC profile (Fig. 3B, lanes 4ai, 5ai, and 6ai), with also one major sulfatelabeled spot (Fig. 3B, lanes 4si, 5si, and 6si). However, the faster migrating spots were lacking or less intense than in S. teranga by. acaciae. (Fig. 3D, lanes 6i and 14i). The TLC profile of these cluster U strains was very similar to that obtained with a <sup>14</sup>C-labeled culture of R. tropici CFN299 (Fig. 3D, lanes 6i and 13i), a tropical strain nodulating L. leucocephala (Martinez-Romero et al. 1991), but also A. senegal and A. nilotica (data not shown).



Fig. 1. Plasmid profiles of representative Sinorhizobium teranga by. acaciae (<sup>4</sup>) and by. sesbaniae (<sup>5</sup>) strains. ORS51 (lane 2); ORS52 (lane 3); ORS53 (lane 4); ORS1072 (lane 6); ORS1073 (lane 7); ORS19 (lane 8); ORS1007 (lane 9); ORS1009 (lane 10); ORS1016 (lane 11); ORS1045 (lane 12). Rhizobium leguminosarum by. viciae VF39SM (lane 1, standard plasmid sizes: 135, 160, 205, 220, 350 and 600 kb) and Sinorhizobium meliloti RCR2011 (lane 5) exhibiting two megaplasmids (arrow) were used as standards.

In contrast, *S. saheli* and *S. teranga* bv. *sesbaniae* bearing pMH682 both produced only nonsulfated Nod metabolites, which migrated closely together (Fig. 3C, lanes 7ao, 8ao, and 9ao). Their TLC profiles appeared very similar to that of *A. caulinodans* ORS571 isolated from *S. rostrata* (Fig. 3D, lane 10i, 11o, and 12o). It is worth noting that the strain ORS1013, isolated from *A. senegal*, but able to nodulate *Sesbania* species, exhibited a typical *Sesbania* Nod factor pattern (Fig. 3C, lane 7ao), and thus we classified it as *S. teranga* bv. *sesbaniae* (Table 1).

Thus Acacia and Sesbania symbionts synthesize different Nod metabolites, and can be easily distinguished on the basis of their supernatant TLC profiles (Fig. 3D, lanes 10i, 11o, and 12o for Sesbania strains and 13i, 6i, and 14i for Acacia nodulating strains).

#### Genomic hybridizations using various nod gene probes.

From the recent determination of the structure of the Nod factors produced by representatives of the different taxonomic groups (J. Lorquin, G. Lortet, M. Ferro, N. Méar, B. Dreyfus, J.-C. Promé, and C. Boivin, unpublished), it appeared that for



Fig. 2. Expression of a *Rhizobium leguminosarum nodA::lacZ* fusion in *Sinorhizobium saheli*, *S. teranga* and genospecies Ua in response to several flavonoids. Values represent average of at least three independent experiments.

both Acacia and Sesbania isolates, the nonreducing end was substituted by a N-methyl and a O-carbamoyl group, whereas the reducing end was glycosylated for Sesbania-nodulating strains and partially sulfated for Acacia-nodulating strains. We thus looked for the presence in these strains of DNA sequences hybridizing to probes consisting of the nodS and nodU, nodH, or nodZ genes, which are involved, respectively, in N-methylation, O-carbamoylation (Jabbouri et al. 1995; Geelen et al. 1995; Mergaert et al. 1995b), sulfation (Lerouge et al. 1990; Roche et al. 1991) and glycosylation (Stacey et al. 1994; Mergaert et al., in press) of Nod factors. Results are summarized in Table 3. Surprisingly, hybridization of genomic DNA with an internal nodH probe from S. meliloti revealed that an homologous DNA sequence was present in all tested strains, even those for which no <sup>35</sup>S-labeled spots could be detected. In fact, except for ORS1047 and ORS1058, which form a subgroup in the S. teranga species (M. Neyra, unpublished results), all Senegalese sinorhizobia shared a common unique 5.5-kb nodH-hybridizing band. All of the strains tested, nodulating either Acacia or Sesbania, lacked detectable homology to an internal nodZ probe from A. caulinodans ORS571. Using the nodSU genes from S. fredii as a probe, Acacia and Sesbania strains hybridized differently: the former gave a clear (cluster U strains) or an intense (S. teranga strains) signal, while no hybridization or only a weak trace could be observed with the latter strains.

## DISCUSSION

Current *Rhizobium* taxonomy is based on a polyphasic approach, consisting of a set of biochemical and molecular techniques (Graham et al. 1991). In contrast, the classification of rhizobia as a function of their symbiotic properties still relies on nodulation tests, which are time- and space-consuming, and only provide a partial characterization of the host spectrum. To develop alternative methods for classifying rhizobia strains on the basis of their host range, we examined possible correlations between host spectrum and various molecular nodulation determinants.

Nodulation experiments were performed with a collection of strains isolated either from *Sesbania* or from *Acacia*, and belonging to the species *S. saheli*, *S. teranga* or to the geno-

		TLC re of plas	esponse after intro mid-borne <i>nodD</i> g		Genospecies Ub ORS1040	
Plasmid (genes present)	Inducer	S. saheli ORS611	S. teranga bv. sesbaniae S. teranga bv. ORS604 acaciae ORS107			Genospecies Ua ORS1001
pMH901						
(nodD1 of S. me)	Luteolin	-	-	-	+	+
pMH682						
(nodD3 and syrM of S. me)	None	+	+	-	-	
pVK31						
(nodD2 of S. sp. BR816)	Naringenin	-	-	+		-
pA28	U					
(nodD1 of S. sp. NGR234)	Apigenin	+	+	+	+	+

Table 2. Nod factor overproduction by Sinorhizobium saheli, S. teranga and Rhizobium sp. from cluster U harbouring nodD genes from various sources<sup>a</sup>

<sup>a</sup> Rhizobium and Sinorhizobium derivatives, harboring an exogenous *nodD* gene, were grown in the presence of <sup>14</sup>C acetate and with or without the appropriate inducing flavonoid. Culture supernatants were analyzed by TLC, using octadecyl silica plates. Presence and absence of flavonoid-inducible compounds are, respectively, designated + and -. However, rhizobia derivatives harbouring pMH982 containing the constitutively expressed *nodD3-syrM* genes are compared to the corresponding wild-type strains for the detection of new radiolabeled compounds. In this case, + and - designate, respectively, presence or absence of *nodD*-inducible compounds. Data are the results of at least three independent experiments.

species Ua and Ub from the so-called cluster U (Table 1). All strains isolated from the same plant genus exhibited similar host ranges, independently of their taxonomic position. *Sesbania* isolates nodulate all *Sesbania* species tested as well as the widely compatible *Acacia seyal* (Dreyfus and Dommergues 1981), but only poorly *A. raddiana*, and not other *Acacia* species. In contrast, the host spectrum of the *Acacia* isolates includes all the species of *Acacia* tested, *Prosopis juliflora* and *Leucaena leucocephala*, but not *Sesbania* species. Since strains with different host ranges belong to the species *S. ter*-

anga, we propose to subdivide this species into biovars acaciae and sesbaniae.

Examining different possible host specificity markers, we showed that Nod factor TLC profile is the only character tightly correlated with the host range of the strains. Whatever their taxonomic position, strains isolated from the same plant genus exhibited similar TLC profiles, which are clearly different for *Acacia* and *Sesbania* isolates (Fig. 3D). In particular, TLC analysis of <sup>35</sup>S-labeled overproducing strain supernatants revealed the partial sulfation of Nod factors from *Acacia* iso-



Fig. 3. Thin-layer chromatography (TLC) separation of Nod factors produced by various Acacia and Sesbania nodulating strains. Radioactivity was visualized after 3 to 8 days exposition to a Kodak X-OMAT K film. a, <sup>14</sup>C-labeled compounds; s, <sup>35</sup>S-labeled compounds; i, cells grown in presence of the appropriate flavonoid inducer; n, non induced cultures; w, wild-type strain; o, overproducing strain harboring pMH682. A, Sinorhizobium teranga bv. acaciae overproducing strains: ORS1071(pVK31) (lanes 1), ORS1007(pVK31) (lanes 2) and ORS929(pVK31) (lanes 3). B, Rhizobium sp. overproducing strains from cluster U: ORS1040(pMH901) (lanes 4), ORS1020(pMH901) (lanes 5) and ORS1001(pMH901) (lanes 6). C, S. saheli and S. teranga bv. sesbaniae wild type and overproducing strains: ORS1013 (lanes 7), ORS19 (lanes 8), and ORS609 (lanes 9). D, Comparison of TLC migration of <sup>14</sup>C-labeled compounds produced by Azorhizobium caulinodans wild-type strain ORS571 (lanes 10), S. saheli ORS611 (lanes 11), S. teranga bv. sesbaniae ORS604 (lanes 12), R. tropici wild-type strain CFN299 (lanes 13), Rhizobium sp. ORS1001 from cluster U (lanes 6) and S. teranga bv. acaciae ORS1073 (lanes 14).

lates, whereas no sulfation could be observed for those of *Sesbania* isolates (Fig. 3C). Furthermore, *A. caulinodans* strain ORS571 and *Sinorhizobium* strains nodulating *Sesbania* showed very similar Nod factors TLC profiles. The recent determination of the chemical structure of the Nod factors produced by *Sesbania*-nodulating *Sinorhizobium* (J. Lorquin, G. Lortet, M. Ferro, N. Méar, B. Dreyfus, J.-C. Promé, and C. Boivin, unpublished) confirmed that this similarity of the TLC profiles reflects the resemblance of the molecules. In the same way, *R. tropici* strain CFN299, which nodulates *A. senegal* and *A. nilotica* (unpublished results), exhibited a Nod factor profile similar to that of *Acacia* strains from cluster U. TLC analysis of Nod factors thus appears to be a powerful screening method to group or distinguish strains on the basis of their host spectra.

A technical difficulty of this method could be the detection of the very small amounts of Nod factors produced by many wild-type strains. We solved this problem by the introduction of exogenous nodD genes cloned in multicopy plasmids into the studied strains. Whereas most of the nodD genes tested were only able to activate Nod factor production in a defined group of strains, the introduction of the nodD1 gene from Sinorhizobium sp. strain NGR234 enabled all the strains to overproduce Nod factor in the presence of apigenin (Table 2). This nodD1 gene had a similar effect on Nod factor production when introduced into various other temperate and tropical Rhizobium species (C. Boivin, unpublished data). Thus, in addition to its ability to respond to a wide range of inducing compounds (Bassam et al. 1988; van Rhijn et al. 1994), the NGR234 NodD1 protein seems to be able to recognize the nod boxes of various rhizobia. The R. leguminosarum nodD derivative FITA (flavonoid independent transcription activation) was also shown to allow Nod factor overproduction in several taxonomically different strains (Lopez-Lara et al. 1995a). The construction of overproducing strains carrying such exogenous nodD genes should greatly facilitate Nod factor characterization, whatever the production level of the wild-type strains. Furthermore, the use of 1-14C glucosamine as Nod factor precursor (Price and Carlson 1995) could allow more specific labeling of these molecules, and should improve the sensitivity of the technique. When Nod factor profiles are closely related, highly resolutive techniques are needed. Resolution of TLC plates could be improved by standardizing the culture conditions, the extraction and elution processes, and by performing two-dimensional TLC (Price and Carlson 1995).

We therefore propose to use the TLC profiling technique, which is currently the fastest way to globally characterize Nod factors, for classifying rhizobial strains on the basis of their host range. This method should constitute a more comprehensive approach than nodulation tests, the confidence of which depends on the choice of appropriated plant species, and on their ability to grow and develop symbiotic interactions in laboratory conditions. An immediate application could be its use for the characterization of biovars in rhizobium taxonomy. To our knowledge, this is the first application of the recent advances in the understanding of the molecular basis of symbiotic specificity for ecological or classification purposes.

A potential limitation to this method could be due to the fact that Nod factor structure is not the only determinant of the specificity in the *Rhizobium*-legume interaction. For instance, *R. etli* and *R. loti* produce identical Nod factors (Lopez-Lara et al. 1995a; Cardenas et al. 1995; Poupot et al. 1995), but differ in host specificity presumably because of different perception of plant flavonoids (Cardenas et al. 1995). Thus, to classify isolates as a function of their host range, the identification of nod gene inducers could be a useful complement to Nod factor profiling. In the present work, all Sinorhizobium strains tested exhibited very similar induction patterns, the highest level of induction being obtained with 10 µM of luteolin (Fig. 2). Luteolin was already known as one of the best nod gene inducers for the closely related species S. meliloti (Peters et al. 1986), but was reported as a poor nod gene inducer for A. caulinodans (Goethals et al. 1989). No inducer could be identified for cluster U strains among a collection of commercial flavonoids. This could be due either to the fact that we did not find the right compound, or to the fact that the NodD proteins of cluster U strains were not able to interact with the foreign nod promoters fused to the lacZ gene. However, it is worth noting that in a R. loti strain, which is taxonomically related to the cluster U, no induction of a R. loti nodA-lacZ fusion could be obtained with commercial flavonoids (Zhang et al. 1995). Our results suggest that the nature of the nod gene inducers seems to be more related to the taxonomic position of a given strain than to its symbiotic properties, and therefore plant signal recognition may not interfere with Nod factor production for the determination of the host range of taxonomically close rhizobia.

From an evolutionary point of view, the fact that all *S. saheli* and *S. teranga* strains present similar responses to plant flavonoids suggests that, at least, the C-terminal part of the NodD proteins, which is mostly responsible for flavonoid recognition (Kondorosi 1992), is conserved among these strains. However, the *nodD* gene sequence was expected to vary as a function of the host plant, since it has been reported that the phylogenetic trees established from *nodD* 3'-sequences of various rhizobia strains follow plant taxonomy, and not rhizo-

Table 3. Presence of sequences homologous to S. meliloti nodH, A. caulinodans nodZ and S. fredii nodSU in S. saheli, S. teranga and Rhizobium sp. from cluster U

	Number of EcoRI fragments homologous to					
Strain	S. me nodH	A. ca nodZ	S. fr. nodSU			
S. saheli						
ORS600	1 <sup>a</sup>	NT	0			
ORS609	1ª	0	0			
ORS611	1ª	0	0			
S. teranga bv. ses	baniae					
ORS604	1 <sup>a</sup>	0	0			
ORS613	1 <sup>a</sup>	0	1			
ORS51	1 <sup>a</sup>	NT	0			
ORS19	1 <sup>a</sup>	0	0			
ORS1013	1ª	0	0			
S. teranga by. acc	ıciae					
ORS1009	1ª	0	1			
ORS1016	1 <sup>a</sup>	NT	2			
ORS1073	1ª	0	1			
ORS1047	1	0	1			
ORS1058	1	0	2			
cluster U						
ORS1001	1	0	1			
ORS1010	NT	NT	1			
ORS1030	NT	NT	1			
ORS1035	NT	0	1			

<sup>a</sup> The *Eco*RI hybridizing fragment was 5.5 kb. NT = not tested.

bia taxonomy (Dobert et al. 1994). Thus, the fact that the NodD proteins of *S. teranga* by. *acaciae* and *sesbaniae* and *S. saheli* exhibit a similar pattern of flavonoid recognition could just reflect the fact that *Acacia* and *Sesbania* root exudates contain common inducers.

The synthesis and transfer of the various substitutions present on the Nod factor oligochitin backbone is under the control of the specific nod genes (see Fellay et al. 1995). We therefore expected that it could be possible to discriminate between Acacia and Sesbania strains by detecting in these strains particular host specificity nod genes. TLC analysis of <sup>35</sup>S-labeled cultures indicated that Nod factors from Acacia strains were partially sulfated, whereas those of Sesbania strains were not (Fig. 3). However, both Acacia and Sesbania strains hybridized with a nodH probe. The nodH-like sequence present in Sesbania strains could therefore be a pseudo gene without biological activity. Such an example of nonfunctional nodulation genes has been described in S. fredii USDA257, where a deletion removing part of the upstream nod box is responsible for the loss of expression of the nodSU genes (Krishnan et al. 1992). However, the fact that the internal nodH probe hybridized to a single common band in most of the S. teranga and S. saheli strains tested indicates a high conservation of that sequence. A possible explanation is that the nodH-like gene encodes a protein also involved in the sulfation of other compounds, such as lipopolysaccharides, as it has been suggested for S. meliloti nodQ genes. (Cedergren et al. 1995). In the case of Sesbania strains, this protein would be unable to transfer the sulfate group to the Nod factors, possibly as a result of structural modifications of these factors.

S. saheli and S. teranga by. sesbaniae Nod factors are fucosylated, whereas those of S. teranga by. acaciae and cluster U strains are not (J. Lorquin, G. Lortet, M. Ferro, N. Mear, B. Dreyfus, J.-C. Prome, and C. Boivin, unpublished). As in B. japonicum (Stacey et al. 1994) and A. caulinodans (Mergaert et al., in press), the fucosylation of the Nod factors is under the control of the nodZ gene, we investigated the possibility that DNA homology with this gene could be a criterion to distinguish between Sesbania and Acacia strains. However, in S. saheli and S. teranga bv. sesbaniae, as well as in Acacia strains, no hybridization signal could be detected with a nodZ probe (Table 3). Similarly, nodS and nodU genes are, respectively, responsible for the N-methylation and the 6-O carbamoylation of the nonreducing terminal glucosamine residue in A. caulinodans (Mergaert et al. 1995a, 1995b) and Sinorhizobium NGR234 (Jabbouri et al. 1995). We used the S. fredii nodSU genes, which show a high degree of homology with their NGR234 counterparts (Krishnan et al. 1992), as a probe to detect the presence of homologous sequences in Sinorhizobium and cluster U strains. Strains isolated from Acacia exhibited homology with this probe, while most of the Sesbania strains from either S. saheli or S. teranga vielded no hybridization signal (Table 3). Such a result was unexpected, since Nmethylation and/or O-carbamoylation of Nod factors are very common in tropical strains, and are encountered in both Acacia (Rhizobium sp. GRH2, R. tropici) and Sesbania (A. caulinodans) nodulating strains (Lopez-Lara et al. 1995b; Poupot et al. 1993; Mergaert et al. 1993). Furthermore, 0carbamoylation and N-methylation of Nod factors in both Acacia and Sesbania isolates was recently established (J. Lorquin, G. Lortet, M. Ferro, N. Méar, B. Dreyfus, J.-C. Promé,

and C. Boivin, unpublished). Taken together, these results suggest that absence of hybridization with *Sesbania* strains was caused by the fact that *nodSU* and *nodZ* alleles diverged significantly in terms of DNA sequences from those of *S. fredii* and *A. caulinodans*, respectively, but yet retained full biochemical activity.

More generally, hybridization results show that in a given strain, the presence or the absence of particular *nod* genes, as assessed by Southern hybridization, is neither sufficient to predict the nature of the substituents present on the Nod factors, nor correlated with the host range. Such discrepancies are presumably due to sequence divergences in genes encoding the same function and presence of pseudo genes having lost their functional activity, as a consequence of genetic rearrangements.

The fact that strains belonging to different taxonomic groups show convergent host ranges strongly suggests that lateral gene transfer has occurred, possibly via plasmid transfer, since in most Rhizobia and Sinorhizobia, nod genes are located on transmissible plasmids. All S. teranga by, acaciae and sesbaniae and S. saheli harbor a megaplasmid and one or two additional large plasmids, a profile similar to that of the closely related species S. meliloti (Bromfield et al. 1987). In contrast, Acacia strains from cluster U usually do not contain any plasmids. Thus, plasmid content of a strain appeared not to be related to its host specificity. This result indicates that a simple transfer of plasmids cannot account for the convergence of host ranges in different species nor for the divergence of host specificities within the same rhizobial species. Emergence of Sesbania and Acacia symbionts within the Sinorhizobium branch thus probably results from complex genetic events, possibly including plasmid transfer followed by genetic recombination, for the acquisition of new genes such as nodZ-like genes, and specific inactivation of other genes such as nodH. A given host plant would select bacteria for which these genetic events lead to the production of Nod factors adapted to specific plant receptors.

## MATERIALS AND METHODS

#### Bacterial strains, plasmids, and growth conditions.

All the wild-type rhizobia listed in Table 1, were described by de Lajudie et al. (1994). Other bacterial strains and plasmids used in this study are described in Table 4. The R. meliloti-R fredii branch including Rhizobium sp. NGR234 and Rhizobium sp. BR816 (Martinez-Romero 1994) has been recently reclassified in the emended genus Sinorhizobium (de Lajudie et al. 1994). In the text R. meliloti, R. fredii, R. sp. NGR234, and R. sp. BR816 were thus referred to as Sinorhizobium meliloti, S. fredii, Sinorhizobium sp. NGR234 and BR816. The complete media for the growth of E. coli and Rhizobium strains were, respectively, LB (Sambrook et al. 1989) and yeast extract-mannitol medium YM (de Lajudie et al. 1994). The minimal medium for nod gene induction (V medium) was as described by Roche et al. (1991). TpBroth medium (in g/liter: 4, peptone; 0.5, yeast extract; 0.5, tryptone; 0.2, MgSO<sub>4</sub>, 7H2O; 0.2, CaCl<sub>2</sub>, 2H<sub>2</sub>O) was used for triparental matings and plasmid content. E. coli, S. teranga, S. saheli, and cluster U strains were grown at 37°C on solid medium and at 30°C in liquid medium.

## Microbiological techniques.

All conjugation experiments were realized using as receptor spontaneous streptomycin-resistant derivatives of wild-type strains described in Table 1. Broad host range plasmids were mobilized from E. coli to Sm<sup>r</sup> rhizobia derivatives with the helper plasmid pRK2013. (Sino)rhizobium and E. coli strains were grown overnight at 30°C in Tpbroth. Samples of donor, helper, and receptor cells (1:1:5, ratio) were pooled, centrifugated, and resuspended in TpBroth 1/50 initial volume. Mating mixtures were spread on TpBroth agar and incubated overnight at 37°C. Selection of transconjugants was performed on YM agar medium supplemented with 10 µg of tetracycline per ml (incP vectors) or 100 µg of chloramphenicol per ml (incQ vectors) for plasmid selection, and 100 µg of streptomycin per ml to select against E. coli donors. Presence of plasmids in transconjugants was confirmed by rapid plasmid DNA extraction and digestion with the appropriate enzymes according to standard procedures (Sambrook et al. 1989).

#### Nodulation tests.

Seeds were superficially sterilized with concentrated sulfuric acid for a duration of 30 (*Acacia senegal*), 150 (*A. raddiana*), 120 (*A. nilotica*), 30 (*Leucaena leucocephala, A. seyal*), 15 (*Prosopis juliflora*), or 60 min (*Sesbania grandiflora* pink flowers; *S. pubescens*; *S. rostrata*). Seeds were then abundantly rinsed with sterile water and allowed to soak for 6 to 24 h. Surface-sterilized seeds were germinated at 30°C for 24 to 48 h, and then transferred to Gibson tubes containing Jensen slant agar and liquid medium (Vincent 1970). Plants were grown under continuous light (20 W/m<sup>2</sup>) at 28°C. Five to eight plants were tested for each strain. Plants were observed for nodule formation during 6 to 8 weeks, and effectiveness was estimated by visual observation of plant vigor and foliage color. *Acacia* and *Prosopis* seeds were obtained from DRPF, ISRA, Senegal.

#### β-Galactosidase assays.

Bacteria were grown overnight to mid-log phase in V medium supplemented with tetracycline (5 µg/ml) or chloramphenicol (50 µg/ml). Bacterial suspension was diluted in fresh medium to  $A_{600} = 0.02$  and flavonoids added at a final concentration of 1 µM (apigenin, naringenin), 2 µM (daidzein, genistein), 10 µM (luteolin), or 5 µM (other flavonoid compounds). Induction was performed at 30°C for 24 h and enzyme activity was assayed as previously described (Boivin et al. 1991). The basal level of β-galactosidase activity of *Rhizobium* and *Sinorhizobium* strains containing pMP220 or pMP190 derivatives was high (150 to 300 units). Luteolin, apigenin, naringenin, butein, daidzein, kaempferol, and genistein were purchased from Extrasynthese (Genay, France). Eriodictyol, umbelliferone, myricetin, and formononetin were a gift from J. Dénarié.

#### Detection of Nod factors by thin-layer chromatography.

An overnight culture was diluted in fresh V medium, supplemented with tetracycline (5 µg/ml) for overproducing strains, to  $A_{600} = 0.02$ , in 2.5 ml of final volume. Ten microcuries of (14C) acetate (56 mCi/mmole, Amersham) or 10 µCi of (<sup>35</sup>S) sulphate (1,000 Ci/mmol, Amersham) was added. Cells were simultaneously induced with 10 µM luteolin (strains bearing pMH901 and wild-type S. saheli and S. teranga strains), 2.5 µM naringenin (strains bearing pVK31), 20 µM naringenin (A. caulinodans ORS571, R. tropici CFN299) or 1 µM apigenin (strains bearing pA28), and incubated at 30°C for 24 h. No inducer was added for strains bearing pMH682. Supernatants of labeled cultures were extracted with n-butanol, applied to octadecyl silica TLC plates (Sigma), and visualized as described by Spaink et al. (1992), except that each extract loaded on a plate corresponded to 1.25 ml of the initial culture supernatant.

Table 4. Other bacterial strains and plasmids used in this study

Designation	Relevant characteristics	Source or reference
Rhizobia		
RCR2011	= SU47, Sinohizobium meliloti <sup>a</sup> wild type, Nod <sup>+</sup> , Fix <sup>+</sup> on Medicago sativa	Rosenberg et al. 1981
ORS571	Azorhizobium caulinodans wild type, Nod <sup>+</sup> , Fix <sup>+</sup> on Sesbania rostrata	Drevfus et al. 1988
CFN299	Rhizobium tropici type IIA wild type, Nod <sup>+</sup> , Fix <sup>+</sup> on Phaseolus vulgaris	Martinez-Romero et al. 1991
VF39SM	Rhizobium leguminosarum by, viciae VF39 derivative, Mel <sup>+</sup> , Sm <sup>r</sup>	Hynes et al. 1988
E. coli	3	
KMBL1164	del(lac-proAB) thi F <sup>-</sup>	P. van der Putte
GMI3355	Nal <sup>r</sup> , Rif <sup>r</sup> derivative of K12	Sambrook et al. 1989
Plasmids		
pRK2013	ColE1, helper plasmid for mobilization of IncP and IncQ plasmids, Km <sup>r</sup>	Figurski and Helsinski 1979
pMP240	pMP220 containing the <i>nodABC</i> promoter of <i>R. leguminosarum</i> by. <i>viciae</i> towards <i>lacZ</i> , Tc <sup>r</sup> , IncP	R. J. H. Okker
pRT311	pMP220 containing the nodABC promoter of R. leguminosarum bv. trifolii strain ANU843 towards lacZ, Tc <sup>r</sup> , IncP	Mc Iver et al. 1989
pMP194	pMP190 containing the <i>nodABC</i> promoter of S. <i>meliloti</i> <sup>a</sup> towards <i>lacZ</i> , Cm <sup>r</sup> , IncP	Spaink et al. 1987
pTD900	pPP375 containing the <i>nodY</i> promoter of <i>B. japonicum</i> towards <i>lacZ</i> , Tc <sup>r</sup> , IncP	G. Stacey
pA28	2.2-kb, nodD1 containing EcoRI-PstI fragment from S. sp. <sup>a</sup> NGR234 pNGR234a; in pRK7813, Tc <sup>r</sup> , IncP	Price et al. 1992
pMH682	9-kb, nodD3 and SyrM containing HindIII fragment from S. meliloti <sup>a</sup> SU47; in pWB5a, Tc <sup>r</sup> , IncP	Honma et al. 1990
pMH901	2.7-kb, nodD1 containing PvuII-HindIII fragment from S. meliloti <sup>a</sup> SU47; in pWB5a, Tc <sup>r</sup> , IncP	Honma et al. 1990
pVK31	3.5-kb, nodD2 containing BglII-XhoI fragment from S. sp. <sup>a</sup> BR816; in pVK100, Tc <sup>T</sup> , IncP	Van Rhijn et al. 1993
pGMI174	6.1-kb, nodGEFH containing EcoRI-PstI fragment from S. meliloti <sup>a</sup> RCR2011; in pBR322, Tc <sup>r</sup>	Debellé and Sharma 1986
pHBK401	3.8-kb, <i>nodSU</i> containing <i>Eco</i> RI fragment from <i>S. fredii</i> <sup>a</sup> USDA257; in pGEM7ZF(+), Amp <sup>r</sup>	Krishnan et al. 1992
pBBRNUZ9	4.8-kb, nodIJZnoeC containing ApaI-HindIII fragment from A. caulinodans ORS571 in pBBR1MCS, Cm <sup>r</sup>	Fernandez-Lopez et al. 1996

<sup>a</sup> The *R. meliloti-R fredii* branch including *Rhizobium* sp. NGR234 and BR816 (Martinez-Romero 1994) has been recently reclassified in the emended genus *Sinorhizobium* (de Lajudie et al. 1994).

### Plasmid profiles.

Strains from S. saheli (ORS609, ORS611), S. teranga (ORS1007, ORS1009, ORS1016, ORS1045, ORS1047, ORS1071, ORS1072, ORS1073, ORS19, ORS22, ORS51, ORS52, ORS53, ORS604, ORS613, and ORS1013) and genospecies Ua (ORS1001, ORS1004, ORS1010, ORS1014, ORS1015, ORS1018, ORS1020, ORS1024, and ORS1029) were analyzed for plasmid content using a modified Eckhardt (1978) technique as described by Bromfield et al. (1987). S. meliloti RCR2011 and R. leguminosarum bv. viciae VF39SM were used as standards for estimation of plasmid size from their electrophoretic mobility.

#### Hybridization of total DNA with nod gene probes.

The pellet of a 20-ml culture grown overnight at 30°C was rinsed with TE buffer (Tris 10 mM, EDTA 1 mM, pH 8), and resuspended in 10 ml of TE, SDS 0.5%, proteinase K 0.3 mg/ml, RNAse 0.3 mg/ml and incubated overnight at 37°C with slight agitation. Total DNA was then cut through a syringe needle and extracted twice with phenol/chloroform/isoamyl (25:24:1), and once with chloroform. After ethanol precipitation, pellet was resuspended in 5 ml of TE. One microgram of total DNA was digested overnight with EcoRI (Pharmacia), separated by agarose gel, and blotted to Biodyne membrane using Southern blot standard procedures (Sambrook et al. 1989). DNA was fixed on Biodyne membrane with 0.4 N NaOH. The hybridization probes were 0.65kb EcoRV fragment of pBBRNIJZ9 (nodZ of A. caulinodans), 2.1-kb PstI-EcoRI fragment of pHBK401 (nodSU of S. fredii), and 0.8-kb AvaII fragment of pGMI174 (nodH of S. meliloti). Twenty to fifty nanograms of desired fragment was probed with 25 µCi (<sup>32</sup>P) dCTP using the ready-to-go DNA labeling kit (Pharmacia), and purified on a microspin S200 HR column (Pharmacia). Prehybridization was performed at 60°C in 5× SSC, 0.5% SDS, 5x Denhardt (bovine serum albumin 0.02%, Ficoll 0.02%, polyvinylpyrrolidone 0.02%), 0.2 mg/ml denaturated salmon sperm DNA for 1 h. Hybridization was performed overnight in the same buffer without salmon sperm DNA, at the same temperature, using the whole labeled probe. The membrane was then rinsed using standard protocols (Sambrook et al. 1989) and exposed to Fuji RX film for 1 to 3 days.

#### ACKNOWLEDGEMENTS

We are grateful to Jean Dénarié for stimulating discussions and critical reading of the manuscript and to Clare Gough for English corrections. We acknowledge our colleagues for providing strains, especially G. Stacey and P. Mergaert for giving unpublished plasmids and F. Maillet for numerous sendings. We also thank Tidiane Badji and Paul Tendeng for technical assistance and ISRA, DRPF for providing plant seeds. Part of this work was supported by grant no. 9506126 from ACC-SV no. 6, Ministère de l'Education nationale, de l'Enseignement Supérieur, de la Recherche et de l'Insertion Professionnelle.

## LITERATURE CITED

- Applebaum, E. R., McLoughlin, T. J., Connell, M. O., and Chartrain, N. 1985. Expression of symbiotic genes of *Rhizobium japonicum* USDA 191 in other rhizobia. J. Bacteriol. 163:385-388.
- Bassam, J. B., Djordjevic, A. M., Redmond, W., Batley, M., and Rolfe, B. G. 1988. Identification of a *nodD*-dependent locus in the *Rhizo-bium* strain NGR234 activated by phenolic factors secreted by soy-

beans and other legumes. Mol. Plant-Microbe Interact. 1:161-168.

- Boivin, C., Barran, L. R., Malpica, C. A., and Rosenberg, C. 1991. Genetic analysis of a region of the *Rhizobium meliloti* pSym plasmid specifying catabolism of trigonelline, a secondary metabolite present in legumes. J. Bacteriol. 173:2809-2817.
- Bromfield, E. S. P., Thurman, N. P., Whitwill, S. T., and Barran, L. R. 1987. Plasmid and symbiotic effectiveness of representative phage types from two indigenous populations of *Rhizobium meliloti*. J. Gen. Microbiol. 133:3457-3466.
- Cardenas L., Dominguez, J., Quinto, C., Lopez-Lara, I. M., Lugtemberg, B. J. J., Spaink, H. P., Jan Rademaker G., Haverkamp J., and Thomas-Oates, J. E. 1995. Isolation, chemical structures and biological activity of the lipo-chitin oligosaccharide nodulation signals from *Rhizobium etli*. Plant. Mol. Biol. 29:453-464.
- Cedergren, R. A., Lee, J. G., Ross, K. L., and Hollingsworth, R. I. 1995. Common links in the structure and cellular localization of *Rhizobium* chitolipooligosaccharides and general *Rhizobium* membrane phospholipid and glycolipid components. Biochemistry 34:4467-4477.
- Debellé, F., and Sharma, S. B. 1986. Nucleotide sequence of *Rhizobium meliloti* RCR2011 genes involved in host specificity of nodulation. Nucleic Acids Res. 14:7453-7472.
- de Lajudie, P., Willems, A., Pot, B., Dewettinck, D., Maestrojuan, G., Neyra, M., Collins, M. D., Dreyfus, B. L., Kersters, K., and Gillis, M. 1994. Polyphasic taxonomy of rhizobia: Emendation of the genus Sinorhizobium and description of Sinorhizobium meliloti comb. nov., Sinorhizobium saheli sp. nov., and Sinorhizobium teranga sp. nov. Int. J. Syst. Bacteriol. 44:715-733.
- Demont, N., Ardourel, M., Maillet, F., Promé, D., Promé, J. C., and Dénarié, J. 1994. The *Rhizobium meliloti* regulatory *nodD3* and *syrM* genes control the synthesis of a particular class of nodulation factors N-acylated by (omega-1)-hydroxylated fatty acids. EMBO J. 13:2139-2149.
- Dénarié, J., Debellé, F., and Rosenberg C. 1992. Signaling and host range variation in nodulation. Annu. Rev. Microbiol. 46:497-531.
- Dénarié, J., and Cullimore, J. 1993. Lipo-oligosaccharide nodulation factors: A minireview. New class of signaling molecules mediating recognition and morphogenesis. Cell 74:951-954.
- Dénarié, J., Debellé, F., and Promé, J. C. 1996. *Rhizobium* lipochitooligosaccharide nodulation factors: Signaling molecules mediating recognition and morphogenesis. Annu. Rev. Biochem. 65:503-535.
- Dobert, R. C., Breil, B. T., and Triplett, E. W. 1994. DNA sequence of the common nodulation genes of *Bradyrhizobium elkanii* and their phylogenetic relationship to those of other nodulating bacteria. Mol. Plant-Microbe Interact. 7:564-572.
- Dreyfus, B. L., and Dommergues, Y. R. 1981. Nodulation of Acacia species by fast- and slow-growing tropical strains of *Rhizobium*. Appl. Environ. Microbiol. 41:97-99.
- Dreyfus, B. L., Garcia, J. L., and Gillis, M. 1988. Characterization of Azorhizobium caulinodans gen. nov., sp. nov., a stem nodulating nitrogen-fixing bacterium isolated from Sesbania rostrata. Int. J. Syst. Bacteriol. 38:89-98.
- Fellay, R., Rochepeau, P., Relic, B., and Broughton, W. J. 1995. Signals to and emanating from *Rhizobium* largely control symbiotic specificity. Pages 199-220 in: Pathogenesis and host specificity in plant diseases. Histopathological, biochemical, genetic and molecular bases, Vol. 1: Procaryotes. U. S. Singh, R. P. Singh and K. Kohmoto, eds. Pergamon/Elsevier Science Ltd., Oxford.
- Fernandez-Lopez, M., D'Haeze, W., Mergaert, P., Verplancke, C., Promé, J. C., Van Montagu, M., and Holsters, M. 1996. Role of nodI and nodJ in lipo-chitooligosaccharides secretion in Azorhizobium caulinodans and Eschericia coli. Mol. Microbiol. In press.
- Figurski, D. H., and Helinski, D. R. 1979. Replication of an origincontaining derivative of plasmid RK2 dependent on a plasmid function provided in trans. Proc. Natl. Acad. Sci. USA 76:1648-1652.
- Geelen, D., Leyman, B., Mergaert, P., Klarskov, K., van Montagu, M., Geremia, R., and Holsters, M. 1995. NodS is an S-adenosine-Lmethionine-dependent methyltransferase that methylates chitooligosaccharides deacetylated at the non-reducing end. Mol. Microbiol. 17:387-397.
- Goethals, K., Gao, M., Tomekpe, K., Van Montagu, M., and Holsters, M. 1989. Common nodABC genes in Nod locus 1 of Azorhizobium caulinodans: Nucleotide sequence and plant-inducible expression. Mol. Gen. Genet. 219:289-298.

- Graham, P. H., Sadowsky, M. J., Keyser, H. H., Barnet, Y. M., Bradley, R. S., Cooper, J. E., De Ley, D. J., Jarvis, B. D. W., Roslycky, E. B., Strijdom, B. W., and Young, J. P. W. 1991. Proposed minimal standards for the description of new genera and species of root and stemnodulating bacteria. Int. J. Syst. Bacteriol. 41:582-587.
- Honma, M. A., Asomaning, M., and Ausubel, F. 1990. Rhizobium meliloti nodD genes mediate host-specific activation of nodABC. J. Bacteriol. 172:901-911.
- Hynes, M. F., Brucksch, K., and Priefer, U. 1988. Melanin production encoded by a cryptic plasmid in a *Rhizobium leguminosarum* strain. Arch. Microbiol. 150:326-332.
- Jarvis, B. D. W., Downer, H. L., and Young, J. P. W. 1992. Phylogeny of fast growing soybean-nodulating rhizobia supports synonymy of Sinorhizobium and Rhizobium and assignment to Rhizobium fredii. Int. J. Syst. Bacteriol. 42:93-96.
- Jabbouri, S., Fellay, R., Talmont, F., Kamalaprija, P., Burger, U., Relic, B., Promé, J. C., and Broughton, W. J. 1995. Involvement of *nodS* in N-methylation and *nodU* in 6-O-carbamoylation of *Rhizobium* sp. NGR234 Nod factors. J. Biol. Chem. 270:22968-22973.
- Jordan, D. C. 1984. Family III. Rhizobiaceae Conn 1938. Pages 234-254 in: Bergey's Manual of Systematic Bacteriology, Vol. 1. N. R. Kreig and J. G. Holt, eds. Williams & Wilkins Co., Baltimore.
- Krishnan, H. B., Lewin, A., Fellay, R., Broughton, W. J., and Pueppke, S. G. 1992. Differential expression of nodS accounts for the varied abilities of *Rhizobium fredii* USDA257 and *Rhizobium* sp. strain NGR234 to nodulate *Leucaena* spp. Mol. Microbiol. 6:3321-3330.
- Kuykendall, L. D., Saxena, B., Devine, T. E, and Udell, S. E. 1988. Genetic diversity in *Bradyrhizobium japonicum* Jordan 1982 and a proposal for *Bradyrhizobium elkanii* sp. nov. Can. J. Microbiol. 38:501-505.
- Lerouge, P., Roche, P., Faucher, C., Maillet, F., Truchet, G., Promé, J. C., and Dénarié, J. 1990. Symbiotic host-specificity of *Rhizobium meliloti* is determined by a sulphated and acylated glucosamine oligosaccharide signal. Nature 344:781-784.
- Lewin, A., Cervantes, E., Chee-Hong, W., and Broughton, W. J. 1990. nodSU, two nod genes of the broad host range Rhizobium strain NGR234 encode host-specific nodulation of the tropical tree Leucaena leucocephala. Mol. Plant-Microbe Interact. 3:317-326.
- Lindström, K., Van Berkum, P., Gillis, M., Martinez, E., Novikova, N., and Jarvis, B. 1995. Report from the roundtable on *Rhizobium* taxonomy. Pages 807-810 in: Nitrogen fixation: Fundamentals and Applications, I. A. Tikhonovich, N. A. Provorov, V. I. Romonov, W. E. Newton, eds. Kluwer Academic Publishers, Dordrecht. The Netherlands.
- Lopez-Lara, I. M., van den Berg, J. D. J., Thomas-Oates, J. E., Glushka, J., Lugtenberg, B. J. J., and Spaink H. P. 1995a. Structural identification of the lipo-chitin oligosaccharide nodulation signals of *Rhizobium loti*. Mol. Microbiol. 15:627-638.
- Lopez-Lara, I. M., van der Drift, K. M. G. M., Van Brussel, A. A. N., Haverkamp, J., Lugtenberg, B. J. J., Thomas-Oates, J. E., and Spaink, H. P. 1995b. Induction of nodule primordia on *Phaseolus* and *Acacia* by lipo-chitin oligosaccharide nodulation signals from broad host range *Rhizobium* strain GRH2. Plant Mol. Biol. 29:465-477.
- Maillet, F., Debellé, F., and Dénarié, J. 1990. Role of the nodD and syrM genes in the activation of the regulatory gene nodD3, and the common and host-specific nod genes of *Rhizobium meliloti*. Mol. Microbiol. 4:1975-1984.
- Martinez-Romero, E., Segovia, L., Mercanre, F. M., Franco, A. A., Graham, P., and Pardo M. A. 1991. *Rhizobium tropici*, a novel species nodulating *Phaseolus vulgaris* L. beans and *Leucaena* sp. trees. Int. J. Syst. Bacteriol. 41:417-426.
- Martinez-Romero, E. 1994. Recent developments in *Rhizobium* taxonomy. Plant Soil 161:11-20.
- McIver, J., Djordjevic, M. A., Weinman, J. J., Bender, G. L., and Rolfe, B. G. 1989. Extension of host range of *Rhizobium leguminosarum* bv. *trifolii* caused by point mutations in *nodD* that result in alterations in regulatory function and recognition of inducer molecules. Mol. Plant-Microbe Interact. 2:97-106.
- Mergaert, P., Van Montagu, M., Promé, J. C., and Holsters, M. 1993. Three unusual modifications, a D-arabinosyl, an N-methyl, and a carbamoyl group, are present on the Nod factors of *Azorhizobium caulinodans* strain ORS571. Proc. Natl. Acad. Sci., USA 90: 1551-1555.
- Mergaert, P., D'Haeze, W., Geelen, D., Promé, D., van Montagu, M., Geremia, R., Promé, J. C., and Holsters, M. 1995a. Biosynthesis of

Azorhizobium caulinodans Nod factors: Study of the activity of the NodABCS proteins by expression of the genes in *Escherichia coli*. J. Biol. Chem. 270:29217-29223.

- Mergaert, P., Goormachtig, S., Geelen, D., Geremia, R., Valerio-Lepiniec, M., Fernandez-Lopez, M., Goethals, K., D'Haeze, W., Promé, J. C., de Bruijn, F. J., Van Montagu, M., and Holsters M. 1995b. Early events in the Azorhizobium caulinodans-Sesbania rostrata symbiosis. Pages 61-66 in: Nitrogen Fixation: Fundamentals and Applications, I. A. Tikhonovich, N. A. Provorov, V. I. Romonov, W. E. Newton, eds. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Mergaert, P., D'Haeze, W., Fernandez-Lopez, M., Geelen, D., Goethals, K., Promé, J. C., Van Montagu, M., and Holsters M. in press. Fucosylation and arabinosylation of Nod factors in *Azorhizobium caulinodans*: Involvement of *nolK*, *nodZ*, and *noeC*, and/or downstream genes. Mol. Microbiol.
- Peters, N. K., Frost, J. X., and Long, S. R. 1986. A plant flavone, luteolin, induces expression of *Rhizobium* nodulation genes. Science 233:977-980.
- Poupot, R., Martinez-Romero, E., and Promé, J. C. 1993. Nodulation factors from *Rhizobium tropici* are sulfated or non sulfated chitopentasaccharides containing an N-methyl-N-acylglucosaminyl terminus. Biochemistry 32:10430-10435.
- Poupot, R., Martinez-Romero, E., Gautier, N., and Promé, J. C. 1995. Wild-type *Rhizobium etli*, a bean symbiont, produces acetylfucosylated, N-methylated, and carbamoylated nodulation factors. J. Biol. Chem. 270:6050-6055.
- Price, N. P. J., Relic, B., Talmont, F., Lewin, A., Promé, D., Pueppke, S. G., Maillet, F., Dénarié, J., Promé, J. C., and Broughton W. J. 1992. Broad-host-range *Rhizobium* species strain NGR234 secretes a family of carbamoylated, and fucosylated, nodulation signals that are O-acetylated or sulphated. Mol. Microbiol. 6:3575-3584.
- Price, N. P. J., and Carlson, R. W. 1995. Rhizobial lipo-oligosaccharide nodulation factors: Multidimensional chromatographic analysis of symbiotic signals involved in the development of legume root nodules. Glycobiology. 5:233-242.
- Roche, P., Debellé, F., Maillet, F., Lerouge, P., Faucher, C., Truchet, G., Dénarié, J., and Promé, J. C. 1991. Molecular basis of symbiotic host specificity in *Rhizobium meliloti. nodH* and *nodPQ* genes encode the sulfation of lipooligosaccharide signals. Cell 67:1131-1143.
- Rosenberg, C., Boistard, P., Dénarié, J., and Casse-Delbart, F. 1981. Genes controlling early and late functions in symbiosis are located on a megaplasmid in *Rhizobium meliloti*. Mol. Gen. Genet. 184:326-333.
- Sambrook, J., Fritsch, E. F., and Maniatis, T. 1989. Molecular cloning: A Laboratory Manual. 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
- Schultze, M., Kondorosi, E., Ratet, P., Buiré, M., and Kondorosi, A. 1994. Cell and molecular biology of *Rhizobium*-plant interactions. Int. Rev. Cytol. 156:1-75.
- Schultze, M., Staehelin, C., Röhring, H. John, M., Schmidt, J., Kondorosi, E., Schell, J., and Kondorosi, A. 1995. In vitro sulfotransferase activity of *Rhizobium meliloti* NodH protein: Lipochitooligosaccharide nodulation signals are sulfated after synthesis of the core structure. Proc. Natl. Acad. Sci. USA 92:2706-2709.
- Segovia, L., Young, J. P. W., and Martinez-Romero, E. 1993. Reclassification of american *Rhizobium leguminosarum* biovar *phaseoli* type I strains as *Rhizobium etli* sp. nov. Int. J. Syst. Bacteriol. 43:374-377.
- Spaink, H. P., Wijffelman, C. A., Pees, E., Okker, R. J. H., and Lugtenberg, B. J. J. 1987. *Rhizobium* nodulation gene *nodD* as a determinant of host-specificity. Nature 328:337-339.
- Spaink, H. P., Aarts, A., Stacey, G., Bloemberg, G. V., Lugtenberg, B. J. J., and Kennedy, E. P. 1992. Detection and separation of *Rhizobium* and *Bradyrhizobium* Nod metabolites using thin-layer chromatography. Mol. Plant-Microbe Interact. 5:72-80.
- Spaink, H. P. 1994. The molecular basis of the host specificity of the *Rhizobium* bacteria. 1994. Antonie van Leeuwenhoek. 65:81-98.
- Stacey, G., Luka, S., Sanjuan, J., Banfalvi, Z., Nieuwkoop, A. J., Chun, J. Y., Forsberg, L. S., and Carlson R. 1994. nodZ, a unique hostspecific nodulation gene, is involved in the fucosylation of the lipooligosaccharide nodulation signal of *Bradyrhizobium japonicum*. J. Bacteriol. 176:620-633.
- van Rhijn, P. J. S., Feys, B., Verreth, C., and Vanderleyden, J. 1993. Multiple copies of *nodD* in *Rhizobium tropici* CIAT899 and BR816. J. Bacteriol. 175:438-447.

van Rhijn, P., Desair, J., Vlassak, K., and Vanderleyden, J. 1994. The NodD proteins of *Rhizobium* sp. BR816 differ in their interactions with coinducers and in their activities for nodulation of different host plants. Appl. Environ. Microbiol. 60:3615-3623.

WAI

- Vincent, J. 1970. A Manual for the Practical Study of Root Nodule Bacteria. I. B. P. Handbook. No. 15. Blackwell Scientific Publications, Oxford.
- Young, J. P. W. 1994. All those new names: An overview of the molecular phylogeny of plant-associated bacteria. Pages 73-80 in: Advances in Molecular Genetics of Plant-Microbe Interactions, M. J.

Daniels, J. A. Downie, and A. E. Osbourn, eds. Kluwer Academic Publishers, Dordrecht, The Netherlands.

- Zhang, X., Harper, R., Karsisto, M., and Lindström, K. 1991. Diversity of *Rhizobium* bacteria isolated from the root nodules of leguminous trees. Int. J. Syst. Bacteriol. 41:104-113.
- Zhang, Z., Scott, B., Stacey, G., Gresshoff, P. M. 1995. Identification of early nodulation signals in *Lotus japonicus*. Page 323 in: Nitrogen fixation: Fundamentals and Applications, I. A. Tikhonovich, N. A. Provorov, V. I. Romonov, W. E. Newton, eds. Kluwer Academic Publishers, Dordrecht, The Netherlands.