## GENETIC ANALYSIS OF LEISHMANIA PARASITES IN ECUADOR: ARE LEISHMANIA (VIANNIA) PANAMENSIS AND LEISHMANIA (V.) GUYANENSIS DISTINCT TAXA?

### ANNE-LAURE BAÑULS, RENAUD JONQUIERES, FRANÇOISE GUERRINI, FRANÇOIS/LE PONT, CESAR BARRERA, IVAN ESPINEL, RONALD GUDERIAN, RAMIRO ECHEVERRIA, AND MICHEL/TIBAYRENC

Centre d'Etudes sur le Polymorphisme des Microorganismes, Unité Mixte de Recherche no. 9926, Centre National de la Recherche Scientifique/Institut de Recherche pour le Développement (IRD), IRD, Montpellier, France; IRD/Instituto Boliviano de Biologia de Altura, La Paz, Bolivia; Department of Clinical Investigations, Vozandes Hospital, Quito, Ecuador; Fundación Eugenio Espejo, Quito, Ecuador

Abstract. In the course of an epidemiologic survey in Ecuador, the following collection of Leishmania stocks was isolated: 28 from patients with clinical signs of leishmaniasis, 2 from sloths, 1 from a dog, and 4 from sand flies. For genetic characterization of these stocks, multilocus enzyme electrophoresis (MLEE) and random amplified polymorphic DNA (RAPD) were used. Twenty six of the 35 stocks were identified as either Leishmania (V.) panamensis or L. (V.) guyanensis, 2 stocks were identified as L. (V.) braziliensis, the 2 stocks from sloths showed specific genotypes, and 5 stocks were characterized as hybrids between L. (V.) braziliensis and L. (V.) guyanensis. These data show that genetic diversity of Leishmania in Ecuador is high and that L. (V.) panamensis/guyanensis is the dominant group in this country. The genetic analysis questioned the distinctness between the two species L. (V.) guyanensis and L. (V.) guyanensis an

Cutaneous and mucocutaneous leishmaniases constituté a serious public health problem in Ecuador, since the disease is endemic in 17 of 20 provinces.<sup>1</sup> The most frequent clinical forms are the cutaneous and mucocutaneous ones, with a large spectrum of clinical variation.<sup>2</sup> This clinical variability of the disease is believed to be due to the Leishmania species diversity encountered in Ecuador.<sup>1</sup> Seven Leishmania species are known to be responsible for infections reported in the country: Leishmania (Viannia) braziliensis, L. (V.) panamensis, L. (V.) guyanensis, L. (Leishmania) mexicana, L. (L.) pifanoi, L. (L.) amazonensis, and L. (V.) equatorensis.<sup>3-6</sup> The most frequently sampled Leishmania species in Ecuador are L. (V.) guyanensis and L. (V.) panamensis.7 These two species belong to the guyanensis complex according to the World Health organization classification.8 Genetic and biochemical analysis demonstrated that they are genetically very close and that only one enzymatic system could be used as a diagnostic marker able to discriminate them.9

This study reports the genetic analysis of 35 Leishmania isolates collected in the course of extensive field studies in Ecuador.<sup>10</sup> This genetic epidemiology study improves our knowledge on the epidemiology of the disease in this country. Moreover, the results clearly raise the question of the distinctness between L. (V.) panamensis and L. (V.) guyanensis.

#### MATERIALS AND METHODS

Isolate collection. The study adhered to the ethical rules of the European Community in accordance with the Ecuadorian government. Twenty-eight stocks were isolated from patients presenting clinical signs, 2 isolates were from sloths, 1 from a dog, and 4 from sandfly vectors (Table 1). Twentysix of these stocks were obtained at 3 study stations selected in departments with a high transmission level of leishmaniasis: La Tablada, Paraiso Escundido, and Zumba (Figure 1). La Tablada and Paraiso Escundido on the Pacific coast were selected based on their geographic location (rural areas not easily accessible) and the endemic character of the disease. Paraiso Escundido is located in the interior part of the country in humid tropical forest, and La Tablada is a littoral village in a dry tropical forest. The third site, Zumba, is located in the southern part of the country in the Amazonian plain. Some parasites were isolated from 2 other sites: 1 stock came from Vozandes Hospital in Quito and 5 other stocks came from Augusto Egas Hospital in Santo Domingo (Table 1). The origin of 3 stocks is unknown.

Pole 2

To better ascertain the species attribution of the stocks surveyed, 11 reference stocks pertaining to different species were added to the study: 3 L. (V.) guyanensis (MHOM/ FG/84/H166, MHOM/BR/78/M5378, and MHOM/GF/85/ LEM669), 2 L. (V.) panamensis (MCHO/PA-755 and MHOM/PA/711/LS94), 3 L. (V.) braziliensis (MHOM/PE/ 89/LH754, MHOM/BO/84/Lpz595, and MHOM/BR/75/ M2903), 1 L. (V.) colombiensis (IHAR/CO/86/CL500), 1 L. (V.) equatorensis (MCHO/EC/82/LSP01), and 1 L. (L.) infantum (MHOM/MA(BE)/67/ITMAP63).

Isolation and culture of parasites. Human and canine stocks were isolated by classical techniques of biopsy and aspirated materials. Stocks PL2 and PL21 were taken from sloths by hepatic puncture. To identify the parasites from an isolate, the technique of Armijos and others was used. The parasites were bulk-cultured in RPMI 1640 medium with 10% fetal calf serum. They were harvested by centrifugation  $(8,000 \times g \text{ for 15 min at 4°C})$  and washed twice in phosphate-buffered saline, pH 7.3.

**Preparation of samples and isoenzyme electrophoresis.** Technical conditions for preparation of samples, electrophoresis, and staining procedures have been described by Ben Abderrazak and others.<sup>11</sup> Cellulose acetate electrophoresis was used. Fifteen enzyme systems were used: aconitase (ACON: EC 4.2.1.3), glucose-6-phosphate dehydrogenase (G6PD: EC 1.1.1.49), glucose phosphate isomerase (GPI: EC 5.3.1.9), glutamate oxaloacetate transaminase (GOT: EC 2.6.1.1), glutamate pyruvate transaminase (ALAT: EC 2.6.1.2), isocitrate dehydrogenase (IDH: EC 1.1.1.42), ma-

Fonds Documentaire ORSTOM Cote:  $B \times 1994^2$  Ex: 1



TABLE 1 World Health Organization codes of Leishmania stocks, hosts, clinical features, localities, isolation dates, and multilocus enzyme electrophoresis (MLEE) identification of Ecuadorian stocks

Stocks (WHO code)	Host	Clinical feature	Locality	Isolation date	MLEE identification
MHOM/EC/92/E72	Human	?	Santo Domingo	2/1/92	Leishmania (Viannia) braziliensis
MHOM/EC/92/E94*	Human	?	Santo Domingo	2/28/92	L. (V.) panamensis/guyanensis
MHOM/EC/92/E91*	Human	?	Santo Domingo	9/28/91	L. (V.) panamensis/guyanensis
MHOM/EC/91/E29	Human	Simple cutaneous lesion (arm)	Santo Domingo	11/30/91	L. (V.) panamensis/guyanensis
MHOM/EC/92/ST6	Human	? .	Santo Domingo	8/4/92	L. (V.) Panamensis/guyanensis
MCHO/EC/91/PL21	Sloth		La Tablada	11/27/91	?
MCHO/EC/92/PL2	Sloth	•	La Tablada	2/21/92	?
ITRA/EC/92/EK747	Lutzomyia trapidoi		La Tablada	8/17/92	L. (V.) panamensis/guyanensis
MHOM/EC/92/A8044*	Human	?	La Tablada	8/04/92	L. (V.) panamensis/guyanensis
MHOM/EC/91/E18	Human	Multiple cutaneous lesions (arms)	La Tablada	11/26/91	L. (V.) panamensis/guvanensis
MHOM/EC/91/E3*	Human	Simple cutaneous lesion (leg)	La Tablada	11/24/91	L. (V.) panamensis/guyanensis
MHOM/EC/91/E4*	Human	Simple cutaneous lesion (arm)	La Tablada	11/24/91	L. (V.) panamensis/guyanensis
MHOM/EC/91/E6	Human	Multiple cutaneous lesions (leg, arms)	La Tablada	11/25/91	L. (V.) panamensis/guyanensis
MHOM/EC/91/E19*	Human	Multiple cutaneous lesions (arm)	La Tablada	11/26/91	L. (V.) panamensis/guyanensis
MHOM/EC/92/E10	Human	Simple cutaneous lesion (leg)	La Tablada (Cacao)	11/25/91	L. (V.) panamensis/guyanensis
MHOM/EC/91/E11	Human	Multiple cutaneous lesions (face, arm, leg)	La Tablada (Cacao)	11/25/91	L. (V.) panamensis/guvanensis
MHOM/EC/91/E14	Human	Multiple cutaneous lesions (leg)	La Tablada (Crisanto)	11/26/91	L. (V.) panamensis/guyanensis
MHOM/EC/91/E12*	Human	Simple cutaneous lesion (shoulder)	La Tablada (Crisanto)	11/25/91	L. (V.) panamensis/guyanensis
MHOM/EC/91/E1* ·	Human	Multiple cutaneous lesions (leg, face)	La Tablada (Crisanto)	11/24/91	L. (V.) panamensis/guyanensis
MHOM/EC/91/E30*	Human	Simple cutaneous lesion (face)	Paraiso Escundido	11/30/91	L. (V.) panamensis/guyanensis
MCAN/EC/92/DOG1*	Dog		Paraiso Escundido	1/9/92	L. (V.) panamensis/guyanensis
ITRA/EC/92/EK112	Lutzomyia trapidoi	• • • • • • • • • • • • • • • • • • •	Paraiso Escundido	10/10/91	L. (V.) panamensis/guyanensis
ITRA/EC/92/EK665*	Lutzomyia trapidoi		Paraiso Escundido	8/11/92	L. (V.) panamensis/guyanensis
ITRA/EC/92/EK649*	Lutzomyia trapidoi		Paraiso Escundido	8/11/92	L. (V.) panamensis/guyanensis
MHOM/EC/92/E79	Human	· · ?	Vozandes (Quito)	2/10/92	L. (V.) braziliensis
MHOM/EC/92/Z9	Human	Cutaneous lesion	Zumba	7/17/92	Hybrid <sup>27</sup>
MHOM/EC/92/Z5	Human	Cutaneous lesion	Zumba	7/17/92	Hybrid <sup>27</sup>
MHOM/EC/92/PR1*	Human	Cutaneous lesion	Zumba	8/28/92	L. (V.) panamensis/guyanensis
MHOM/EC/92/P5	Human	Cutaneous lesion	Zumba (Palanda)	6/23/94	Hybrid <sup>27</sup>
HMOM/EC/91/E67*	Human	Cutaneous lesion	Zumba (Palanda)	1/20/92	L. (V.) panamensis/guyanensis
MHOM/EC/92/E102	Human	Cutaneous lesion	Zumba (Pucapamba)	4/16/92	Hybrid <sup>27</sup>
MHOM/EC/92/E107	Human	Cutaneous lesion	Zumba (Pucapamba)	4/16/92	Hybrid <sup>27</sup>
MHOM/EC/-/E73*	Human	Cutaneous lesion	?	?	L. (V.) panamensis/guyanensis
MHOM/EC-E50*	Human	Multiple cutaneous lesions (leg, arm)	?	?	L. (V.) panamensis/guyanensis
MHOM/EC-E49	Human	Simple cutaneous lesion (arm)	?	?	L. (V.) panamensis/guyanensis

\* Stocks used for the random amplified polymorphic DNA analysis.



FIGURE 1. Map of Ecuador showing the locations of *Leishmania* stock sampling areas.

late dehydrogenase NAD+ (MDH: EC 1.1.1.37), malate dehydrogenase NADP+ or malic enzyme (ME: EC 1.1.1.40), mannose phosphate isomerase (MPI: EC 5.3.1.8), nucleoside hydrolases, substrate inosine and substrate deoxyinosine, I and D, respectively (NHI and NHD: EC 2.4.2.\*), peptidases 1 and 2 (PEP1 and PEP2: EC 3.4.4.11 or 13), 6-phospho-gluconate dehydrogenase (6PGD: EC 1.1.1.44), and phosphoglucomutase (PGM: EC 2.7.5.1). The 15 enzyme systems used made it possible to study 16 different putative loci. Two different loci could be distinguished with the NHI system, *Nhi* 1 and *Nhi* 2.

**Random amplified polymorphic DNA (RAPD) protocol.** The RAPD fingerprinting method, first described by Welsh and McClelland<sup>12</sup> and Williams and others,<sup>13</sup> has been used in the characterization of various parasitic protozoa by Tibayrenc and others.<sup>14</sup>

A subset of 16 stocks of this Ecuadorian sample (marked by an asterisk in Table 1) plus 5 reference stocks, pertaining either to the *L*. (*V.*) *panamensis* or to *the L*. (*V.*) *guyanensis* species (3 *L*. (*V.*) *guyanensis*: MCHO/FG/83/CAY A116, MHOM/BR/78/M5378, and MHOM/FG/85/LEM669; 2 *L*. (*V.*) *panamensis*: MCHO/PA-M4039 and MHOM/CO/83/ REST417), was further analyzed by RAPD fingerprinting.

The DNA was isolated according to the protocol of Sambrook and others.<sup>15</sup>

Thirteen primers in a kit (Operon Technologies, Inc., Alameda, CA) were used: A10: GTGATCGCAG, B8: GTCCA-CACGG, F1: ACGGATCCTG, F13: GGCTGCAGAA, N13: AGCGTCACTC, N20: GGTGCTCCGT, R13: GGACGA-CAAG, R14: CAGGATTCCC, R15: GGACAACGAG, R16: CTCTGCGCGT, R20: CTCTGCGCGT, U15: ACGGGCCA-GT, and U16: CTGCGCTGGA.

Identification of stocks and estimation of genetic relationships. The electrophoretic profiles of Ecuadorian stocks were compared with those of stocks used as reference for the different species. On the basis of band similarity and known diagnostic loci, a species attribution was made for each stock.

The genetic relationships among the stocks were established by computing Jaccard distances from multilocus enzyme electrophoresis (MLEE) and RAPD data for all possible pairwise comparisons:<sup>16</sup> Dij = 1 - (a/a + b + c) where a = number of bands that are common to the i and j stocks, b = number of bands recorded for stock i and absent for stock j, and c = number of bands recorded for stock j and absent for stock i. From the distance matrix obtained, unweighted pair-group method with arithmetic averages (UPGMA) dendrograms were designed.<sup>17</sup>

Population genetic analysis. Leishmania population structure in this area was explored by the analysis of linkage disequilibrium or nonrandom association of genotypes occurring at different loci. The tests proposed by Tibayrenc and others<sup>18</sup> were used. They are based on the null hypothesis that the population is panmictic (recombination occurs at random). Any statistical departures from the panmictic expectations show that gene flow is inhibited in the population. The d1 test calculates the combinatorial probability of sampling the most common genotype as often as, or more often than, the observed frequency. The d2 test calculates the probability of observing any genotype as often as, or more often than the most common genotype in the sample. The e test calculates the probability of observing as few or fewer genotypes in the population than observed in the sample. The f test calculates the probability of observing a linkage disequilibrium in the population as high or higher than the one observed in the sample. The d2, e, and f tests are based on computer simulations (Montecarlo tests) with 10<sup>4</sup> runs. The biases generated by either spatial or temporal isolation, and the means to avoid them, have been reported by Tibayrenc and others.19

#### RESULTS

The species attribution of each stock was based on the diagnostic loci identified by Guerrini (Guerrini F, 1993. Génétique des Populations et Phylogénie des Leishmania du Nouveau-Monde. PhD Thesis. Université des Sciences et Techniques du Languedoc, Montpellier France) and on the comparison with the reference stocks. Table 1 shows the species identification of the isolates by MLEE. Of the 35 stocks examined, 2 were identified as L. (V.) braziliensis, the 2 isolated from sloths, PL2 and PL21, had a specific genotype different from any known Leishmania species, 26 were identified as either L. (V.) panamensis or L. (V.) guyanensis and 5 had complex isoenzyme patterns (Table 1). The stocks identified as L. (V.) panamensis/guyanensis were isolated either from humans, a dog, or sand flies (Lutzomyia trapidoi). The 6-Pgd locus described in the literature<sup>9</sup> as a diagnostic marker to distinguish L (V.) panamensis from L. (V.) guyanensis showed many different profiles in the L. (V.) panamensis/guyanensis group. In contrast, the Nhi 2 locus showed only 2 different patterns in the L. (V.) panamensis/ guyanensis group. In the L. (V.) panamensis/guyanensis group (26 stocks), 21 different isoenzyme profiles or zymodemes were recorded (genotypic variability = 0.807). Five stocks from the Zumba Area showed complex patterns in the Nhi 2 and Pgm loci.

The UPGMA dendrogram designed from the Jaccard's distance matrix separated the zymodemes (distinct isoenzyme profiles) into 4 distinct clusters, which are not linked to either host or geographic origin, except for the cluster 2 (Figure 2). The first cluster included all the strains identified as either L. (V.) panamensis or L. (V.) guyanensis, as well as the L. (V.) panamensis and L. (V.) guyanensis reference stocks. The average genetic distance within the L. (V.) panamensis/guyanensis group (without the reference stocks) was 0.264 with a standard deviation of 0.104. The second cluster is composed of only the 5 stocks from Zumba exhibiting complex Nhi 2 and Pgm profiles. The third cluster included the L. (V.) braziliensis reference stocks and the two stocks E79 and E72. The fourth cluster included the L. (V.) equatorensis and L. (V.) colombiensis reference stocks and the 2 stocks PL2 and PL21 isolated from sloths. These sloth stocks appeared to be equally distant (genetic distance = 0.7) from both reference stocks. Leishmania (L.) infantum was phytogenetically very distant from all other stocks included in the study, which is consistent with the assignment of this species to another subgenus (Leishmania).

The RAPD technique was performed only on a subset of 16 Ecuadorian stocks belonging to the L. (V.) panamensis/ guyanensis group (Table 1) and 5 reference stocks belonging to either L. (V.) panamensis species or L. (V.) guyanensis. We found no RAPD fragment specific to either L. (V.) panamensis or L. (V.) guyanensis. Furthermore, the UPGMA analysis did not show any structuring in 2 clear-cut clusters.

We performed the population genetic tests of recombination from isoenzyme data in the *L*. (*V.*) panamensis/guyanensis group, which is the largest part of the sample. The 5 stocks with complex *Nhi* 2 and *Pgm* patterns, which pose specific problems of interpretation, were excluded from the analysis, together with the reference stocks. Finally, the 26 Ecuadorian stocks of cluster 1 were used for the population genetic analysis. We used the recombination tests d1, d2, e, and f, which are all related to the linkage disequilibrium phenomenon, but explore different facets of it.<sup>17</sup> The results were as follows: d1:  $P = 5.9 \times 10^{-3}$  (for the genotype of the EK649 stock, represented 2 times),  $9 \times 10^{-3}$  (for the genotype of the E12 stock, represented 3 times); d2: *P* not significant; e: *P* not significant; f:  $P = 10^{-4}$ .

#### DISCUSSION

In Ecuador, the predominant clinical forms are the cutaneous and mucocutaneous leishmaniases with a large spectrum of clinical variation. It has been inferred that clinical variability of the disease was due to the diversity of *Leishmania* species encountered in Ecuador.<sup>1</sup>

Two of the 35 stocks analyzed were identified as L. (V.) *braziliensis*. These stocks were from a patient from in hospital in Santo Domingo, and from a patient in Vozandes Hospital in Quito (Figure 1). These results confirm previously published data showing the presence of L. (V.) *braziliensis* in Ecuador.<sup>3</sup> The identification of 2 L. (V.) *braziliensis* stocks is consistent with the presence of mucocutaneous clinical forms in Ecuador.<sup>20, 21</sup>

Two stocks, PL2 and PL21, isolated from sloths in the site at La Tablada (Figure 1) have specific isoenzyme multilocus genotypes. Their isoenzyme profile is distinct from any known Leishmania species. Nevertheless, they are clustered, although remotely, with the L. (V.) colombiensis and L. (V.) equatorensis reference stocks (Figure 2). These 2 sloth stocks have similar genotypes (only 1 allele difference). It is unexpected that these 2 sloth stocks are not identified as L. (V.) panamensis, since Edentata are known to be the major reservoir of this Leishmania species.<sup>22, 23</sup> It would be necessary to increase the sample of Leishmania stocks from sloths in Ecuador to better understand the epidemiology and taxonomic status of these specific genotypes.

The L. (V.) panamensis/guyanensis group is dominant in the present sample. Indeed, of 35 examined stocks, 26 stocks isolated from humans, a dog, or Lu. trapidoi pertain to this group. These results are consistent with the data of Hashigushi and others.<sup>5</sup> Only a few cases of dog infection by the species L. (V.) panamensis/guyanensis have been were reported.<sup>24</sup> Until now, the domestic dog has been previously demonstrated to be essentially the reservoir of L. (L.) infantum. One case of canine infection by L. (V.) tropica has been reported.25 It is also suspected to be a reservoir for the L. (V.) peruviana species.<sup>26</sup> Nevertheless, clinico-epidemiologic data suggest that the dog infection by the species L. (V.) panamensis/guyanensis recorded here is accidental, since this case was the only one in an extensive survey performed on dogs. The 4 stocks isolated from Lu. trapidoi were also identified as L. (V.) panamensis/guyanensis. These data are consistent with the notion that Lu. trapidoi is the major vector of L. (V.) panamensis in Ecuador.<sup>23</sup>

The 5 stocks from Zumba grouped in cluster 2 (Figure 2) could not be classified by the classical diagnostic markers in the *L*. (*V*.) panamensis/guyanensis group. The complex profiles of *Nhi* 2, *Pgm*, and *Mpi* loci for the stock P5 were interpreted as heterozygous patterns between the *L*. (*V*.) guyanensis and *L*. (*V*.) braziliensis species.<sup>27</sup>

On the MLEE UPGMA dendrogram, it is impossible to distinguish any clear additional subdivision within the cluster 1 composed of the L. (V.) panamensis and L. (V.) guyanensis reference stocks and the Ecuadorian stocks related to them. This lack of clear-cut subdivisions is fully confirmed by the RAPD data obtained from a subset of 16 Ecuadorian stocks and the reference stocks belonging to the L. (V.) guyanensis and L. (V.) panamensis species, which again show no tendency of clear clustering into 2 discrete groups that would correspond to each of these 2 species.

The 6-Pgd locus, which has been proposed as a diagnostic locus between these 2 species, shows considerable allelic diversity and does not permit any clear-cut subdivisions within the Ecuadorian sample of L. ( $\ddot{V}$ .) panamensis/guyanensis stocks (Figure 3).<sup>9</sup>

The only locus that could suggest a possible discrimination between the 2 species is Nhi 2 (Table 2). All L. (V.) guyanensis reference stocks show an allele with a strong enzyme activity, whereas the L. (V.) panamensis reference stocks show a weak or null allele (Figure 4). The Ecuadorian stocks have either one or the other of these 2 clearly distinct profiles. The 4 stocks isolated from Lu. trapidoi were also characterized by specific monoclonal antibodies, and were identified as L. (V.) guyanensis (Le Pont F, unpublished data). This species identification corresponds to the one obtained with the Nhi 2 locus (Table 2).

Nevertheless, based on only 1 of 15 isoenzyme loci and

842

Ì

BANULS AND OTHERS



FIGURE 2. Unweighted pair group method with arithmetic averages dendrogram built from Jaccard's genetic distances calculated from multilocus enzyme electrophoresis data. The genotypes that are specific for a given cluster are noted on the dendrogram. For definitions of enzyme loci, see Materials and Methods.

13 RAPD primers, it would be extremely tentative to infer that L. (V.) panamensis and L. (V.) guyanensis correspond to true distinct monophyletic lineages (discrete typing units [DTUs]).<sup>23</sup> If this were true, these two DTUs would be closely related, and the relevance of describing them as separate taxonomic entities would be doubtful. Table 2 summarizes the tentative species attributions that can be drawn from the 6-Pgd and Nhi 2 loci.

In the L. (V.) panamensis/guyanensis group considered as a whole, the genotype diversity is high (0.807), whereas the

phylogenetic variability is limited (mean  $\pm$  SD genetic distances = 0.264  $\pm$  0.104). Diagnostic characters (Tags)<sup>28</sup> can be proposed for the entire *L*. (*V*.) panamensis/guyanensis group (including the reference stocks, without the hybrid stocks) based on the *Nhi*, *Pgm*, and *Mpi* loci.

The 5 stocks from Zumba with complex isoenzyme patterns could not be classified as either L. (V.) panamensis/ guyanensis or L. (V.) braziliensis. Indeed, these stocks showed heterozygous profiles for the three loci (Nhi 2, Pgm, and Mpi) able to discriminate L. (V.) braziliensis from L.



FIGURE 3. Multilocus enzyme electrophoresis profiles obtained with the 6-phosphoglucontate dehydrogenase enzyme system. Lane 1, M2903 (*Leishmania (Viannia) braziliensis* reference stock); lane 2, E72; lane 3, E94; lane 4, EK649; lane 5, M4039 (*L. (V.) panamensis* reference stock); lane 6, EK665; lane 7, no profile because of enzyme degradation); lane 8, A8044.

#### TABLE 2

Tenative attribution to either *Leishmania* (Viannia) panamensis or L. (V.) guyanensis by 2 isoenzyme loci (6PGD and Nhi 2) of the *Leishmania* stocks isolated in the present study\*

Code	6PGD	Nhi 2
E73	L. (V.) panamensis	L. (V.) panamensis
E50	L. (V.) panamensis	L. (V.) panamensis
E49	L. (V.) panamensis	L. (V.) guyanensis
EK747	?	L. (V.) guyanensis
A8044	L. (V.) guyanensis	L. (V.) panamensis
E18	L. (V.) panamensis	L. (V.) guyanensis
E3	L. (V.) panamensis	L. (V.) guyanensis
E4	L. (V.) panamensis	L. (V.) guyanensis
E6	? `	L. (V.) guyanensis
E19	?	L. (V.) guyanensis
E10	L. (V.) guyanensis	L. (V.) panamensis
E11	L. (V.) panamensis	L. (V.) guyanensis
E14	?	L. (V.) guyanensis
E12	L. (V.) panamensis	L. (V.) guyanensis
El	L. (V.) panamensis	L. (V.) guyanensis
E30	L. (V.) panamensis	L. (V.) guyanensis
DOG1	L. (V.) panamensis	L. (V.) panamensis
EK112	L. (V.) panamensis	L. (V.) guyanensis
EK665	L. (V.) panamensis	L. (V.) guyanensis
EK649	`?	L. (V.) guyanensis
E94	?	L. (V.) guyanensis
E91	?	L. (V.) panamensis
E29	?	L. (V.) guyanensis
ST6	?	L. (V.) guyanensis
PR1	L. (V.) guyanensis	L. (V.) guyanensis
E67	?	L. (V.) panamensis
755	L. (V.) panamensis	L. (V.) panamensis
LS94	L. (V.) panamensis	L. (V.) panamensis
LEM669	L. (V.) guyanensis	L. (V.) guyanensis
M5378	L. (V.) guyanensis	L. (V.) guyanensis
H166	L. (V.) guyanensis	L. (V.) guyanensis

\* 6PGD = 6-phosphogluconate dehydrogenase, Nhi 2 = nucleoside hydrolase substrate inosine 2; ? = stocks that do not have the 6PGD band level of L (V) guyanensis reference stocks or the 6PGD band level of L (V) panamensis reference stocks. (V.) guyanensis/panamensis. For each stock, the other enzymatic systems presented profiles that were also observed either for the L. (V.) braziliensis reference strains or for L. (V.) panamensis/guyanensis strains. Based on the complete analysis of their complex genotypes and on already published RAPD data, we have proposed that these 5 stocks correspond to hybrid genotypes between L. (V.) panamensis/ guyanensis and L. (V.) braziliensis.27 If we accept as a working hypothesis that the Nhi 2 strong allele corresponds to L. (V.) guyanensis, these 5 stocks would be hybrids between this species and L. (V.) braziliensis. There heterozygous patterns corresponding to this locus are 5-banded (the nucleoside hydrolase inosine substrate 2 is a tetrameric enzyme). Conversely, the hybrids postulated by Belli and others<sup>29</sup> in Nicaragua would be hybrids between L. (V.) panamensis and L. (V.) braziliensis, since their heterozygous pattern for the Nhi 2 locus is asymmetrical 3-banded (this pattern would correspond to an heterozygous tetrameric enzyme with a null allele).

The d1 and f linkage disequilibrium tests showed highly significant results, which shows that gene flow is severely restricted in this set of stocks. The most parsimonious hypothesis to account for this result is a basically clonal population structure, which has already been inferred for other *Leishmania* species and other major human parasites.<sup>18</sup> Several comments can be made regarding this hypothesis. 1) It is not parsimonious to attribute departures from panmictic expectations to geographic distance and genetic drift. If this were the case, distribution of genotypes would be linked to geographic distance, which is not verified here.<sup>19</sup> 2) The 5 putative hybrid genotypes were excluded from the linkage disequilibrium analysis. Departures from panmictic expectations are not therefore attributable to the presence of genes



FIGURE 4. Multilocus enzyme electrophoresis profiles obtained with the nucleoside hydrolase substrate inosine (*Nhi*) enzyme system. Lane 1, M5378 (*Leishmania* (V.) guyanensis reference stock); lane 2, E72; lane 3, E94; lane 4, EK649; lane 5, M4039 (*L.* (V.) panamensis reference stock); lane 6, EK665; lane 7, EK747; lane 8, A8044.

from another species. 3) The existence of hybrids is not incompatible with the hypothesis of a basically clonal population structure. As a matter of fact, this hypothesis does not imply that sex is totally absent. It states only that recombination is not frequent enough to break the prevalent pattern of clonal population structure.

Financial support: This work was supported by European Community International Scientific Cooperation grant no CI1\*-CT91-0901, by a "Groupement de Recherche" CNRS/French Army grant, by an ACC-SV7 "Sciences du Vivant" French Ministry of Research grant, and by Fonds voor Wetenschappelijk Onderzoek (FWO) (National Loterij, grants FGWO 9.0024.90).

Authors' addresses: Anne-Laure Bañuls, Renaud Jonquieres, Françoise Guerrini, and Michel Tibayrenc, Centre d'Etudes sur le Polymorphisme des Microorganismes, Unité Mixte de Recherche no. 9926, Centre National de la Recherche Scientifique/Institut de Recherche pour le Développement (IRD), IRD, BP 5045, 34032 Montpellier, Cedex 1, France. François Le Pont, IRD/Instituto Boliviano de Biologia de Altura, Casilla, 9214 La Paz, Bolivia. Cesar Barrera, Ivan Espinel, and Ronald Guderian, Department of Clinical Investigations, Vozandes Hospital, CP 691 Quito, Ecuador. Ramiro Echeverria, Fundación Eugenio Espejo, Quito, Ecuador.

#### REFERENCES

- Chico ME, Guderian RH, 1989. Caracteristicas fotograficas de la leishmaniasis en el Ecuador. *Rev Med Vozandes 3:* 56-66.
   Hashiguchi Y, Gomez Landires EA, 1991. A review of leish-
- maniasis in Ecuador. Bull Pan Am Health Organ, 25: 64-76.
  Mimori T, Grimaldi Jr G, Kreutzer RD, Gomez EA, McMahon-Prätt D, Tesh RB, Hashiguchi Y, 1989. Identification, using isoenzyme electrophoresis and monoclonal antibodies, of Leishmania isolated from humans and wild animals of Ecuador. Am J Trop Med Hyg 40: 154-158.
- Armijos RX, Chico ME, Cruz ME, Guderian RH, Kreutzer RD, Berman JD, Rogers MD, Grögl M, 1990. Human cutaneous Leishmaniasis in Ecuador: identification of parasites by enzyme electrophoresis. Am J Trop Med Hyg 42: 424–428.
- 5. Hashigushi Y, Gomez EA, de Coronel VV, Mimori T, Kawabata M, Furuya M, Nonaka S, Takaoka H, Alexander JB, Quizhpe AM, Grimaldi Jr G, Kreutzer RD, Tesh RB, 1991. Andean leishmaniasis in Ecuador caused by infection with *Leishmania*

mexicana and L. major-like parasites. Am J Trop Med Hyg 44: 205-217.

- Grimaldi Jr G, Kreutzer RD, Hashiguchi Y, Gomez EA, Mimory T, Tesh RB, 1992. Description of *Leishmania equatorensis* sp.n. (Kinetoplastida: Trypanosomatidae), a new parasite infected arboreal mammals in Ecuador. *Mem Inst Oswaldo Cruz* 87: 221-228.
- Armijos RX, Weigel MM, Izurieta R, Racines J, Zurita C, Herrera W, Vega M, 1997. The epidemiology of cutaneous leishmaniasis in subtropical Ecuador. *Trop Med Int Health 2:* 140– 152.
- 8. World Health Organization, 1990. Control of the Leishmaniases. World Health Organ Tech Rep Ser 793.
- Lainson R, Shaw JJ, 1987. Evolution, classification and geographical distribution. Peters W, Killick-Hendrick R, eds. *The Leishmaniasis in Biology and Medicine*. Volume 1. London: Academic Press, 1-120.
- Barrera C, Herrera M, Martinez R, Leon R, Richard A, Guderian RH, Mouchet J, Echeverria R, Le Pont F, 1994. Leishmaniose en Equateur. 1. Incidence de la leishmaniose tegumentaire sur la façade pacifique. Ann Soc Belg Med Trop 74: 1-12.
- Ben Abderrazak S, Guerrini F, Mathieu-Daudé F, Truc P, Neubauer K, Lewicka K, Barnabé C, Tibayrenc M, 1993. Isoenzyme electrophoresis for parasite characterization. Hyde JE, ed. Protocols in Molecular Parasitology. Volume 21. Walker IM, series ed. Methods in Molecular Biology. Totowa, NJ: Humana Press, 361-382.
- 12. Welsh J, McClelland M, 1990. Fingerprinting genomes using PCR with arbitrary primers. Nucleic Acids Res 18: 7213-7218.
- Williams JG, Kubelik AR, Livak KJ, Rafalski JA, Tingey SV, 1990. DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Res* 18: 6531–6535.
- Tibayrenc M, Neubauer K, Barnabe C, Guerrini F, Skarecky D, Ayala FJ, 1993. Genetic characterization of six parasitic protozoa: parity between random-primer DNA typing and multilocus enzyme electrophoresis. *Proc Natl Acad Sci USA 90*: 1335-1339.
- Sambrook J, Fritch EF, Maniatis T, 1989. Molecular Cloning. A Laboratory Manual. Cold Spring Harbor, NY: Cold Spring Harbor Press.
- Jaccard P, 1908. Nouvelles recherches sur la distribution florale. Bull Soc Vaudoise Sci Nat 44: 223–270.
- 17. Sneath PHA, Sokal RR, 1973. Numerical Taxonomy. San Francisco: W. H. Freeman.
- 18. Tibayrenc M, Kjellberg F, Ayala FJ, 1990. A clonal theory of

parasitic protozoa: the population structures of Entamoeba, Giardia, Leishmania, Naegleria, Plasmodium, Trichomonas, and Trypanosoma and their medical and taxonomical consequences. Proc Natl Acad Sci USA 87: 2414-2418.

- Tibayrenc M, Kjellberg F, Arnaud J, Oury B, Breniere SF, Darde ML, Ayala FJ, 1991. Are eukaryotic microorganisms clonal or sexual? A population genetics vantage. *Proc Natl Acad Sci* USA, 88: 5129-5133.
- Wyler D, Marsden PD, 1984. Leishmaniasis. Warren K, Mahmoud A, eds. *Tropical and Geographical Medicine*. New York: McGraw-Hill, 28.
- Hashigushi Y, Gomez EA, 1987. A brief review of leishmaniasis in Ecuador. Hashigushi Y, ed. Studies on New World Leishmaniasis and Its Transmission, with Particular Reference to Ecuador. Research Reports Series No. 1. Kochi, Japan: Kyowa Printing Co., 5-32.
- Lainson R, Shaw JJ, 1979. The role of animals in the epidemiology of the South American leishmaniasis. Lumsden WH, Evans DA, eds. *Biology of the Kinetoplastida*. Volume 2. London: Academic Press, 1-116.
- Grimaldi Jr G, Tesh R, 1993. Leishmaniases of the New World: currents concepts and implications for future research. *Clin Microbiol Rev 3*: 230-250.

- Dedet JP, 1993. Leishmania et leishmanioses du continent américain. Ann Inst Pasteur 4: 3-25.
- 25. Guessons-Idrissi N, Berrag B, Riyad M, Sahibi H, Bichichi M, Rhalem A, 1997. Short report: *Leishmania tropica*: etiologic agent of a case of canine visceral leishmaniasis in northern Morocco. Am J Trop Med Hyg 57: 172–173.
- Herrer A, 1982. Lutzomyia peruensis Shannon, 1929, possible natural vector of uta (cutaneous leishmaniasis). Rev Inst Med Trop Sao Paulo 24: 168–172.
- 27. Bañuls AL, Guerrini F, Le Pont F, Barrera C, Espinel I, Guderian R, Echeverria R, Tibayrenc M, 1997. Evidence for hybridization by multilocus enzyme electrophoresis and random amplified polymorphic DNA between *Leishmania braziliensis* and *Leishmania panamensis/guyanensis* in Ecuador. J Eukaryot Microbiol 44: 408-411.
- Tibayrenc M, 1998. Genetic epidemiology of parasitic protozoa and other infectious agents: the need for an integrated approach. Int J Parasitol 28: 85-104.
- 29. Belli AA, Miles MA, Kelly JM, 1994. A putative Leishmania panamensis/Leishmania braziliensis hybrid is a causative agent of human cutaneous leishmaniasis in Nicaragua. Parasitology 109: 435-442.

ISSN: 0002-9637

**VOLUME 61** 

## NOVEMBER 1999

NUMBER 5

# The American Journal of TROPICAL MEDICINE HYGIENE



. .